

*executive committee of
the directing council*



**PAN AMERICAN
HEALTH
ORGANIZATION**

*working party of
the regional committee*

**WORLD
HEALTH
ORGANIZATION**



111th Meeting
Washington, D.C.
June-July 1993

Provisional Agenda Item 4.13

CE111/20 (Eng.)
25 May 1993
ORIGINAL: SPANISH

HEALTH OF INDIGENOUS PEOPLES

The Subcommittee on Planning and Programming, during its 18th Meeting, held in April 1992, recommended that a Hemispheric workshop on the health of indigenous peoples be held, in Canada, for the purpose of consulting with the indigenous peoples regarding their health situation and possible approaches to indigenous health. The Subcommittee also requested that the results of this event be communicated to the Governing Bodies of the Organization. The Canadian Society for International Health (CSIH), the technical representative of PAHO in Canada, took responsibility for organizing the workshop, which was held in Winnipeg, Canada, from 13 to 18 April 1993.

This document is composed of two main parts. The first part outlines the results of the Winnipeg Workshop, drawing from the presentations, discussions, and analysis that took place there. The second part proposes, in light of the recommendations made by the workshop participants, the basis and guidelines for the Organization's actions and those of its Member Governments.

The members of the Executive Committee are requested, after examining this document, to:

- Express their opinion regarding its content, with particular attention to the proposed lines of action.
- Propose strategies to be followed by the Organization and its Member Governments in order to move forward with the implementation of the Indigenous Peoples' Health Initiative.

CONTENTS

| | <u>Page</u> |
|--|-------------|
| 1. Introduction | 1 |
| 2. Indigenous Peoples and Health Workshop, Winnipeg, 1993 | 1 |
| 2.1 Indigenous Peoples and Health Workshop | 1 |
| 2.2 The Geopolitical and Social Context of Health in the Indigenous Populations of the Americas | 2 |
| 2.3 Health of Indigenous Peoples at the Beginning of the 1990s | 4 |
| 3. Basis and Guidelines for Action | 8 |
| 3.1 Basis for Action by PAHO | 8 |
| 3.2 General Guidelines for Action | 10 |
| 3.3 General Cooperation Activities | 15 |
| REFERENCES | 16 |

ANNEXES

Annex I. Indigenous Peoples and Health Workshop

Annex II. Estimated Indigenous Population in the Americas

Annex III. Principal Health and Disease Figures in the Indigenous Population in Selected Countries

1. INTRODUCTION

The Initiative on the Health of Indigenous Peoples of the Americas presented here has deep roots in the life of the countries, of the indigenous peoples, and of the Organization. Although it is recognized that in the different countries of the Region valuable initiatives have been launched with a view to improving the living conditions and health status of the indigenous peoples of the Hemisphere, these have usually been sporadic, short-term efforts. The Organization has provided its cooperation in several of these initiatives at various times.

In April 1992, in response to a proposal of the Delegation of Canada, which was echoed by the official delegations of Mexico, Ecuador, Peru, and Bolivia, the subject was included on the agenda of the 18th Meeting of the Subcommittee on Planning and Programming (SPP) of the Executive Committee of PAHO. In the presentation of this item, it was stated that the health of indigenous peoples was "perhaps the most technically complex and politically difficult health issue of the day" (PAHO, 1992). The members of the SPP, in response to this challenge and after their deliberations, were unanimous in underscoring the relevance and utmost importance of the issue, and approved the holding of a regional workshop in 1993, with the express recommendation that the meeting be participatory and allow for broad consultation with representatives of the indigenous peoples of the Americas. The Canadian Society for International Health (CSIH), the technical representative of PAHO in Canada, with the support of the Organization and other sponsoring agencies such as the International Development Research Center (IDRC), took responsibility for making the preparations and organizing the Indigenous Peoples and Health Workshop, which was held in the city of Winnipeg, Canada, from 13 to 18 April 1993. The results of this Workshop served as a basis for the formulation of the initiative on the health of the indigenous peoples of the Americas (Annex I).

2. INDIGENOUS PEOPLES AND HEALTH WORKSHOP, WINNIPEG 1993

2.1 *Indigenous Peoples and Health Workshop*

The Workshop was held in the city of Winnipeg, Manitoba, Canada, from 13 to 18 April 1993, and attended by 68 participants from 18 countries of the Region of the Americas. Also present were delegates from indigenous organizations, communities, and nations of the continent, official governmental delegations, and representatives of international agencies and nongovernmental organizations.

The Winnipeg Workshop was an opportunity for consultation at which a series of recommendations was prepared for the activities of PAHO, the Member Governments, and other organizations. These were then approved in plenary session by the participants. During the deliberations the following fundamental principles were

identified and adopted which formed the basis for subsequent discussions and for the recommendations:

- i) the need for a holistic approach to health;
- ii) the right to self-determination of indigenous peoples;
- iii) the right to systematic participation;
- iv) respect for and revitalization of indigenous cultures; and
- v) reciprocity in relations.

The Winnipeg Workshop also underscored the importance of approaching the health of indigenous peoples in its proper geopolitical and social context, taking into account the historical processes in progress.

2.2 The Geopolitical and Social Context of Health in the Indigenous Populations of the Americas

The term "indigenous population" refers to a diversity of peoples and cultures, without reflecting the special identity and idiosyncrasies of each people in particular. In order to understand the health of the indigenous peoples of the Americas, it is necessary to recognize their great ethnic and cultural diversity, as well as the complex interrelationships between peoples and cultures, identity and health.

It is significant that 500 years after the Europeans first set foot in America, there is persistent ambiguity and lack of agreement as to how to define the "indigenous" ethnic category (also called native, Indian, Amerindian, aboriginal, autochthonous, etc). The general tendency among the countries of the Americas has been to use changing definitions, or simply to omit ethnic classification from national population censuses, as well as from the ongoing registration of demographic information such as births, deaths, and migrations.¹

¹ The Workshop debated the acceptance of the term "indigenous," as well as the connotations of its usage, and agreed that "self-identification" as indigenous would be a fundamental criterion for defining the members of this group. ILO Convention 169 concerning Indigenous and Tribal Peoples in Independent Countries (1989) recognizes as "indigenous" that distinct section of the national community which is understood to consist of: "...peoples in independent countries who are regarded as indigenous on account of their descent from the populations which inhabited the country, or a geographical region to which the country belongs, at the time of conquest or colonization or the establishment of present state boundaries and who, irrespective of their legal status, retain some or all of their own social, economic, cultural, and political institutions" (Convention 169, Article 1).

Based on the estimates and sources available, the indigenous population of the Americas is approximately 42 million², distributed among some 400 ethnic groups. This represents around 6% of the total population of the Region of the Americas and slightly less than 10% of the population of Latin America and the Caribbean (Annex II). In socio-demographic terms, this population can be broken down into two main clusters. The first cluster numbers approximately 18 million and includes the indigenous peoples of Mesoamerica, consisting mainly of the Maya of Mexico and Guatemala, followed in decreasing order by other ethnic groups living in Mexico, Belize, Honduras, El Salvador, Panama, Nicaragua, and Costa Rica. The second cluster, estimated at around 20 million, is made up principally of Quechua and Aymara who are concentrated today in the Andean countries, mainly in Bolivia, Peru, and Ecuador, and to a lesser degree in Venezuela, Colombia and Chile. These two clusters, in Mesoamerica and the Andean region, together make up the bulk of the indigenous population: around 38 million people, or more than 80% of the indigenous population of the Americas.

A third, scattered cluster, estimated at approximately 3 million people, includes a heterogeneous conglomerate of Indian groups and nations who today live in the subarctic regions and in various settlements in Canada (more than 350,000) and the United States (approximately 1.6 million). Also included in this cluster are the peoples of the Caribbean islands, Guyana, and Suriname; Venezuela, Brazil (especially the Amazon region); Paraguay and the Southern Cone, with Argentina and Chile.

Finally, in recent years, the migrant indigenous population has been growing in importance. The migratory group can be broken down into seasonal migrants, who generally follow the harvests, and a broad sector of permanent migrants, who are part of the ongoing trend toward rural-urban migration recorded in recent decades. The members of this group live and work today in several of the medium-sized and large cities of Latin America and the United States. In Central America, a significant proportion of the migrant indigenous population is made up refugees and political exiles displaced by internal conflicts or "low intensity" wars. Despite the growing number of incidents involving inter-ethnic violence and conflict in various parts of the world, including Latin America, very little attention has been paid to the consequences for the physical and mental health of the indigenous population.

² Jordán Pardo, III-FAO, 1990.

The indigenous population is more severely affected by poverty than the rest of society throughout the continent. Living conditions, per capita income, employment, education, access to basic water, sanitation, and health services, housing conditions, and food availability for these people all fall below the national averages.

The indigenous population of the Region is passing through different stages of the acculturation process, in addition to feeling the impact of exogenous development models, the emergence of nation-states, and attempts at "national integration." All have been significant determinants in the disarticulation of the American indigenous cultures and the continuous devaluing of the indigenous identity, resulting in the marginalization of indigenous peoples. The issue of land ownership and use is a problem of utmost importance for the autochthonous groups of the Region. Land is central to life, culture, and history, and determines to a large extent the survival of indigenous peoples and nations, as well as their standards of living, health, and nutrition.

2.3 Health of Indigenous Peoples at the Beginning of the 1990s

The delegates of the indigenous peoples and organizations at the Indigenous Peoples and Health Workshop in Winnipeg emphasized the need to preserve the holistic approach inherent in the different health concepts of the indigenous peoples of the Americas. The common denominator in the most widely varying perspectives is the belief that health expresses dynamic relationships between inseparable components. These include the relationship between individual aspects (physical, mental, spiritual, and emotional) and group aspects (political, economic, cultural, and social), and between what is natural and what is social (Dion Stout, 1992; Rozental, 1988).

There is general agreement that Latin America and the Caribbean should be considered a multi-ethnic and culturally pluralistic region. However, the official sources in most of the countries of the Region have little or no information on the health status and living conditions of their indigenous peoples. As a rule, the epidemiological information available in the disease registries and the principal indicators of morbidity or mortality, birth rates, and life expectancy at birth are not disaggregated by ethnic or language group. However data and information are available from secondary sources that provide unmistakable evidence that the levels of health and nutrition of the autochthonous populations of America are several times below the national averages, reaching alarmingly low figures in some regions when compared with homologous reference populations.

The disease profile that characterizes the indigenous peoples of the Region features many of the same conditions that plague other socioeconomically disadvantaged groups (Annex III). Viral diseases (influenza, measles, dengue, poliomyelitis, arboviral respiratory diseases, hepatitis B, etc.) frequently explode into epidemics, particularly

among groups with low levels of immunity. The prevalence of diseases endemic to tropical and subtropical areas (e.g. leishmaniasis, onchocerciasis, cysticercosis, Chagas' disease, etc.) remains high, and especially affects those human settlements where indigenous people are a majority. Other communicable diseases such as tuberculosis and malaria are on the rise again. Primary health care workers frequently report high incidence rates and lethality of cholera in indigenous populations, as well as a considerable increase in the occurrence of sexually transmitted disease. The spread of AIDS poses an added and very grave risk for the indigenous people who live in areas with high rates of HIV infection.

The Winnipeg Workshop recognized that mental disorders and problems are a growing source of concern. Stress-related disorders, including violence toward others, depression, and suicide, as well as accidental and violent death, join abuse of alcohol, tobacco, and other substances as problems that show an increasingly high prevalence among young and adult indigenous people of both sexes. Protein-calorie malnutrition is a persistent problem, along with diseases stemming from deficiencies of micro-nutrients, especially iron, vitamin A, and iodine. Thyroid hyperplasia, cholelithiasis, obesity, and diabetes mellitus are frequently occurring conditions, particularly in the North American indigenous population.

The health profile of indigenous women is largely determined by the subordination they face on two fronts: in their couple relationship and in relation to the dominant sectors of local and national society. In addition to the illnesses described earlier, women suffer from problems related to reproduction (e.g. pregnancy at an early age; complications of pregnancy and delivery; iron-deficiency anemia, etc.) and others related to mental health (for example, sexual abuse and violence; alcoholism and drug abuse). And there are other more specific problems deriving from hazardous working conditions in agriculture, in the informal urban or service sector, and in industry.

Care during childbirth and the puerperium, disposal of the placenta, care of the umbilical cord, breast-feeding, and the care and feeding of indigenous infants are strongly influenced by the culture. It is in these areas that indigenous communities frequently experience difficulties in accessing the services that are available, and culturally rooted discrepancies appear between the medical services provided in the hospital and the home care that is administered by family members and traditional midwives.

From this perspective, health, health-disease processes, and health systems can be seen as cultural systems. The purely medical approach in health services delivery, in addition to failing to meet demand, is inadequate to deal with an epidemiological profile that is so complex and difficult to resolve. Moreover, traditional healing practices, while known to be efficient for the management of various culturally-bound

afflictions and syndromes, still fall short when it comes to articulating an effective response to the "new" profile of diseases and health problems arising from the new context (for example, the AIDS threat, degenerative diseases, etc.). When indigenous peoples are exposed to these problems, they are unable either to produce the appropriate biological response (e.g. immunity) or to make the necessary sociocultural adjustment (for example, in traditional medical practices).

Finally, most of the countries do not have sustained initiatives and adequate financing to support the development of specific policies and programs on traditional medicine and indigenous therapeutic resources (for example, medicinal plants), or research and development of alternative models of care for indigenous populations or particular ethnic groups. With some exceptions, measures have not been taken at the official level to address this situation. A few of the Ministries of Health have established ad hoc groups or offices to oversee the health of indigenous communities in the national territory, usually as part of projects financed by bilateral agencies, philanthropic organizations or religious missions. The nongovernmental sector is promoting local initiatives in almost all the countries and many nongovernmental organizations (NGOs) and private foundations are actively committed to working with organized indigenous communities on development projects and efforts to improve living conditions in marginalized urban and rural areas. However, these initiatives are generally of limited coverage and duration.

The ethnic and cultural heterogeneity of the indigenous peoples makes it difficult or impossible to adopt single programs or universal health care models. This diversity, in which ethnic and cultural differences are accentuated, means that each indigenous people must be considered individually and that the emphasis must therefore shift toward the development of local health care strategies. Given that the morbidity profile of the indigenous population is different from that of other ethnic groups, proposals to provide differential care and actions appear to be the most valid (SSA-INI, 1992). If local health systems are that set of processes that comprises all social activities in health at the local level, including but not restricted to health services delivery (PAHO 1993), the strategy of development of local health systems is a valid response to this health situation, particularly in areas with a diverse ethnic population or a significant proportion of indigenous inhabitants.

The Environmental Health Program's evaluation of the International Drinking Water Supply and Sanitation Decade in the late 1980s showed that although progress has been made, it has mainly benefited the urban populations of Latin America and the Caribbean. Moreover, most of the water supply systems have operational problems that hinder the process of continuous disinfection, while only 5% of the wastewater from sewerage systems receives adequate treatment. In the rural areas, almost half the population does not have access to clean water, and two thirds do not have services for

the disposal of excreta and refuse. The countries with the highest proportions of indigenous inhabitants also have the lowest percentages of their population covered by water supply and excreta disposal services (PAHO, 1990).

Water resources are particularly affected by the activities associated with the mining of metals (for example, copper, aluminum, tin, lead, etc.). Since mining activities frequently take place in high mountainous areas--such as the Andes--they pose their greatest risk to indigenous communities, which are the most directly exposed to the contamination. In the rural environment, contamination results from the use of chemical fertilizers, pesticides, and organophosphate insecticides, as well as from the disposal of toxic or radioactive waste. The problem has grown to the point where significant traces of substances such as DDT, as well as toxic levels of mercury, have been detected in surface water, food, and other basic nutrients necessary for survival, such as breast milk.³

In many countries of the Amazon Pact, especially in Brazil and in the Amazon basin of the Andean countries, the most important challenges to human and environmental health for many indigenous communities of the Amazon region are: persistent and uninterrupted over-exploitation of natural resources (wood, gold, oil, and other resources such as rubber in the past) in the "low" jungle by companies, independent miners, or "garimpeiros" [prospectors], together with successive waves of migration toward the "high" jungle, the presence of evangelizing missions, military incursions (associated with border conflicts, repression of subversive activities, etc.) or civil incursions to engage in illegal activities such as smuggling and drug traffic. In addition, the colonization of land, the construction of roads, dams, and hydroelectric plants, and other development projects have considerably increased the proliferation of vectors or intermediary hosts, with a consequent rise in the transmission of certain diseases among indigenous populations.⁴

Both the discussions and the final recommendations of the Workshop emphasized the interdependence between indigenous peoples and the natural environment, in particular stressing the importance of access to those natural resources that are considered essential for their health and survival (food and nutrition, shelter, water and energy sources, medicinal plants, etc.). In the conceptual framework of the indigenous peoples of the Americas, human beings are at one with the natural environment. Thus their

³ It is interesting to note that suicide by insecticide poisoning is reported to be frequent among indigenous groups, to the point that in some countries, the majority of deaths are from deliberate poisoning by organophosphate insecticides and not accidental exposure of agricultural workers (PAHO, 1990: 225).

⁴ According to official sources in Brazil, for example, malaria prevalence is from two to ten times higher in communities near the Trans-Amazon highway, than in those located in more remote regions (de Koning, 1992).

cosmic view and their practices are constructed from nature, in which they live and from which they are inseparable.

In short, it is recognized that: (a) The health situation of the indigenous peoples is the end result of a historical process that has fostered dependency, loss of identity, and marginalization. (b) The indigenous peoples of the Americas therefore have a shorter life expectancy than homologous national groups, along with higher mortality and a distinct and changing morbidity profile that reflects their lower standard of living, social status, and level of acculturation, as well as their higher risk of disease and death. The health of indigenous peoples is largely determined by their environmental conditions and the new challenges posed by impoverishment and the modernization process. (c) Health, health-disease processes, and health systems for the indigenous peoples are cultural systems.

3. BASIS AND GUIDELINES FOR ACTION

The recommendations made by the delegates to the Winnipeg Workshop in response to the consultation launched by the Organization were in line with the stated principles and had the goal of contributing to joint resolution of the problems identified and analyzed in the description of the health situation. A synthesis of the recommendations is attached which tries to capture their original sense. Although the original text and the order of priority assigned by the Workshop participants have been modified in the drafting of this annex to the document, there have been no substantive changes in the content.

In the original text the recommendations were grouped into several sections: health policies, human resources, research, health programs and projects in indigenous regions and communities, and others covering the international and intergovernmental sphere (Annex I).

3.1 *Basis for Action by PAHO*

Taking into account the subjects covered in Annex I and the previous sections, and after an analysis of the Organization's background, strategies, and programs, several proposals for the future have been developed which comprise the policy foundations and guidelines for action in the area of health of indigenous peoples in the Region of the Americas.

Indigenous peoples must regain control over their own lives, of which health is only one aspect. A fundamental means of reaching this goal is to progressively restore power to the indigenous populations to decide on policies and strategies for their own

development and to look for consensus with groups in power so that they can recover and have access to land and means of production, as well as to the resources necessary to satisfy their basic needs. They must regain control over the production of goods in general, and basic social services in particular, since this is critical to their being able to break the cycle of dependency that has perpetuated poverty, racial discrimination, and marginalization among most of the Region's indigenous peoples.

In trying to attain these political goals it is important to reassess the value of indigenous wisdom and to strengthen the unique elements of indigenous cultures, recognizing that it is the members of these cultures who have the best understanding of their own people, their health and development needs, and the responses that need to be implemented. However, this should not be used to justify further isolation; rather, joint efforts should be promoted to overcome common obstacles (economic, political, social, cultural) through the forging of interethnic and intercultural ties that are based on reciprocity, mutual respect, and coexistence.

These premises set the course for the Indigenous Peoples' Health Initiative and present challenges of a complexity that is not addressed by the current mandates of PAHO/WHO and which is beyond the capacity of the health sector at the country level. These challenges also orient the policy foundations presented here.

The first challenge to the Indigenous Peoples' Health Initiative is to promote joint efforts and shared responsibility by PAHO and its Member Governments and the indigenous organizations and communities, in conjunction with national and international agencies and organizations (governmental and nongovernmental). In this regard, PAHO is in a position to offer its experience and resources as a catalyst to stimulate numerous efforts at different levels by a wide variety of actors.

The challenge posed by the lack of sufficient and adequate knowledge and information on the health of indigenous peoples in the face of a need for immediate action and impact must be met through the establishment of strategies that allow adequate knowledge and information to be generated during the action ("learning by doing"), as well as making it possible to systematically store up the knowledge and information gained through experience ("learning from both past and present experience"). The resulting knowledge should serve as a navigational chart to guide the Initiative and its proponents.

A third challenge arises out of the Initiative's inherent multidimensionality and diversity. The various dimensions of the Initiative reflect the concrete political, economic, social, and cultural realities that manifest themselves at every level, from individual and community to national and international. Recognizing diversity means formulating proposals that respond to particular situations and contexts which vary from

country to country and region to region, and from one people to another. Accordingly, it is indispensable to simultaneously address all facets of the Initiative at all levels, and to involve the entire Organization in this process, from the Headquarters to the Representative Offices to the countries themselves, placing special emphasis on local experiences and processes where actions of proven impact and concrete viability are required. In this way it will be possible to generate responses as varied and diverse as the situations and peoples involved, so that these experiences can build up a store of knowledge and conclusions that will form the basis for and help to stimulate numerous other processes and actions.

Finally, the underlying premise for the articulation of the general guidelines for action is the recognition that the indigenous peoples are not the problem, but rather those most severely affected by the urgent demands of the prevailing context which, in one way or another, impacts on all the peoples of the Region. In addition, these peoples are the possessors of a rich and diverse culture which is essential to our survival as a species, and to the preservation of life. Thus the ultimate goal of working together to solve this set of problems is not to help the indigenous peoples, but rather to help each other in order to achieve health for all.

Another aspect to be considered is the need to prioritize the Organization's work during this initial phase. For this purpose it is proposed that the efforts and guidelines for action be focused on those countries where the greatest need exists, as well as on those which show the strongest interest in and commitment to the Indigenous Peoples' Health Initiative.

3.2 *General Guidelines for Action*

Seven guidelines for action are presented below:⁵

3.2.1 *Health Promotion*

The Indigenous Peoples' Health Initiative is in principle a health promotion initiative. This is apparent when reviewing the doctrine behind this strategy as defined in the Ottawa Charter for Health Promotion, adopted by the First International Conference

⁵ These are consistent with Strategic Orientations and Program Priorities of the Organization.

on Health Promotion, in November 1986 (WHO, 1986). The Ottawa Charter for Health Promotion states that "Health promotion is the process of enabling people to increase control over, and to improve, their health."^{6, 7}

Health promotion was adopted at the XXIII Pan American Sanitary Conference in 1990 as a strategic orientation for the work of PAHO during the quadrennium that began in 1991. It was defined as "the sum activity of the population, the health services, the health authorities, and other productive and social services, aimed at improving the status of individual and collective health." Thus health promotion is a strategy for making the concept of health in development a reality (PAHO, 1991). Accordingly, the health promotion strategy is a fundamental orientation to support the Indigenous Peoples' Health Initiative.

3.2.2 Transformation of the Health Sector: Local Health Systems, Equity and Access to Health Care Services, Decentralization, and Community Participation

The transformation of national health systems and the development of local health systems (Paganini et al., 1990) are valuable tactical resources in efforts to overcome current problems of deficient coverage, lack of access, poor acceptance and low health impact in the health systems and services for autochthonous populations. Community participation is one of the fundamental aspects of the local health system strategy, growing out of the recognition that it is essential to develop horizontal and symmetrical ties with indigenous organizations and communities, and to thus open the door to consensus among the different community actors at the local level. The effort to increase equity, through decentralization, intersectoral action, and participatory research, is extremely important to the implementation of the Indigenous Peoples' Health Initiative (PAHO/WHO, Document CD33/14; PAHO, 1990).

It is in the context of local health systems that the traditional wisdom of autochthonous peoples can be preserved and in turn articulated with institutional

⁶ This same document underscores the importance of peace, shelter, education, food, access to basic services, drinking water and sanitation, income, a stable ecosystem, and sustainable resources. The Ottawa Charter, in stating the principles of health promotion, issues a call for equity as the sole means of closing the health gaps between countries and peoples (WHO, 1986).

⁷ The Bogota Declaration on health promotion, known as "Promotion of Health and Equity" (Santa Fe, Bogota 1992), underscores the importance of the relationship that exists among equity, community participation, culture, development, and health in Latin America. The strategies and commitments adopted in Bogota by the representatives of the 21 participating countries are, as a group, of utmost importance to the Indigenous Peoples' Health Initiative. It is worth noting, for example, that in the Declaration, the countries commit themselves to stimulate a dialogue about different knowledge bases so that the health development process can incorporate all elements of the Region's cultural heritage (Promotion of Health and Equity, Commitment 8, 1992).

knowledge and efforts. Local health systems also provide a place where it is possible to measure the impact of different activities carried out in line with mutually agreed upon goals for concerted health and well-being. The development and strengthening of local health systems is the fundamental practical strategy to achieve health in the face of a diversity of situations and specific needs at the local level. The obvious choice for technical cooperation aimed at consolidating the Indigenous Peoples' Health Initiative is to provide support for these experiences, for exchanges within and between countries, for the mobilization of resources, and for efforts by other sectors and entities to support and strengthen the impact and viability of these efforts. This is also the most convincing argument for working to sensitize institutions at all levels so that they commit themselves to support greater equity in and access to health services and community participation.

3.2.3 Regional Plan for Investment in the Environment and Health: Environmental Health, Preservation of Habitat, and Protection of Traditional Lifestyles

PAHO's Regional Plan for Investment in the Environment and Health (PIAS) is a crucial initiative for indigenous peoples. It supports the mobilization of resources and efforts to improve their marginal status and provide basic services that guarantee a minimum level of well-being, as well as recognizing the traditional knowledge about the natural environment that these people possess.

Innovative strategies and culturally appropriate technologies also need to be developed in order to adapt water supply and basic sanitation programs to indigenous populations. It is also indispensable to draw upon, document, evaluate, and revitalize traditional knowledge and technologies that have to do with preservation of the habitat and adequate management of natural resources, as well as to document and evaluate the impact that development projects and activities to extract natural resources have on the environment and health of indigenous populations at the local level.

3.2.4 Human Resources

In line with the recommendations adopted in this area at the Winnipeg Workshop, activities are needed that are fundamentally oriented toward: (i) formulating educational strategies to train health professionals and health workers as part of the Indigenous Peoples' Health Initiative; (ii) providing support to training centers so that incentives are created to give indigenous people more access to professional careers, and curricula are adapted to help these people preserve their identity and maintain their commitment to their community of origin; and (iii) to draw upon experiences and initiatives in the Region to help the Member Countries generate strategies and incentives that will encourage health workers to work in indigenous communities.

3.2.5 Evaluation and Monitoring of the Health Status and Living Conditions of Indigenous Peoples

The Organization has accumulated substantial experience in documenting and evaluating health status and living conditions in almost all the countries of the Region. However, appropriate mechanisms are needed for the collection and utilization of information that encompass the perspective and support of indigenous organizations and communities.

It is therefore imperative to have basic information along with adequate and low-cost monitoring systems (e.g. using sentinel populations or selected samples) that will make it possible to periodically evaluate progress and impact in projects and interventions in autochthonous communities. Strategies and instruments for rapid assessment are also available which could provide a short-term solution to the lack of information on certain sectors and problems.

3.2.6 Formulation of Health Policies for Indigenous Peoples and Legal and Ethical Issues

The Member Countries need help in revising the macroscopic and sectoral policies that address or should address the specific health problems of the indigenous peoples in the country or in a specific region, taking into account the strong presence of indigenous identities and the reiterated proposal calling for national multicultural and multilingual states that welcome diversity and pluralism (PAHO/HSP, 1993). At the heart of these new policies is support for the preservation and reevaluation of indigenous peoples, and for attempts to promote new types of interrelationship between these peoples, the state, and the national society, including the so-called solidarity sectors (América Indígena, 1990).

It is important to move ahead in establishing cooperative arrangements with the Member Countries to support their identification of immediate selective actions for the development of health policies that can exercise a stronger impact on the health status of this population. The areas of policy that merit special consideration and action have to do with traditional practices and the utilization and preservation of medicinal plants.

In recognition of the effect, importance, and cultural value of traditional medical practices, legal questions are pursued with the specific goal of revising the pertinent legislation and legal codes in order to limit or reduce the provisions that discount or proscribe such practices.

With regard to the ethical principles that should guide research on human populations, gaps and ambiguities persist that are urgently in need of being addressed by the Organization, particularly in the case of research⁸ involving indigenous populations.

3.2.7 *Health Programs, Areas, and Problems of Particular Importance*

The participants at the Winnipeg Workshop cited the importance of participatory research as a fundamental and important aspect of all the guidelines listed here. PAHO and the research institutions or centers at the country level should give priority to and promote research on the field of indigenous health and traditional medical practices.

Some of the problems that have a serious impact on indigenous peoples and thus require special attention include: i) various types of violence; ii) alcohol and substance abuse; iii) infectious diseases; iv) sexually transmitted diseases; v) problems related to contamination of the habitat and the work environment; and vi) malnutrition and micro-nutrient deficiencies. The programs that the Workshop participants mentioned as being of special note, in addition to those already mentioned, were: i) mental health; ii) health, indigenous women, and development; iii) communication for the health of indigenous peoples;⁹ iv) food and nutrition;¹⁰ and v) health of migrant populations, particularly transitory migrants in border areas.

The Organization, in coordination with public institutions, nongovernmental agencies, educational and research centers, and indigenous organizations, can serve as the catalyst for successful experiences carried out in the countries as part of the Indigenous Peoples' Health Initiative. It is important to note the central role played by the Representative Offices, which, working in coordination with the services and organizations in each country, can draw on experiences in progress, promoting and coordinating new experiences between institutions, while at the same time promoting activities under the Organization's Regional programs.

⁸ For example, ethnopharmacology, genetics, epidemiology, population studies, etc.

⁹ It is important to develop, with broad participation by indigenous communities and experts at all levels, communication strategies in the areas of health and education, at all levels of action of the initiative on the health of indigenous peoples: from heightening of awareness among governments, institutions, and society at large to the dissemination of culturally appropriate information to, from, and between indigenous communities and their organizations and the professional and technical health sectors.

¹⁰ This has to do not just with eating habits and traditional agricultural practices, but rather, and above all, with sources of income, land-holding and use, preservation and exploitation of the natural habitat, and the development of highly nutritious traditional food products.

3.3 *General Cooperation Activities*

Based on the analysis of the context and health status of the indigenous peoples in the Americas, the policy foundations and guidelines for action presented, and the recommendations of the Winnipeg Workshop, several general cooperation strategies and activities are proposed for the implementation of the Indigenous Peoples' Health Initiative 1993-1995. The Governing Bodies of the Organization will be asked to recommend strategies, lines of action, and new lines of cooperation for the countries.

It is felt that cooperation activities should be organized along two main lines: (i) participation by indigenous peoples in the leadership and management of the Indigenous Peoples' Health Initiative from the earliest stages; and (ii) a coordinated program of cooperation activities under the Indigenous Peoples' Health Initiative, to be carried out through the Organization's regular divisions and programs.

In principle, it is proposed to establish operations for interprogram coordination, resource mobilization, and technical support for the initiatives at the Regional level and in selected countries. Once these operations are established, support will be provided for meetings and workshops in the countries to promote networking so that basic information can be exchanged and ties established between official institutions and NGOs, collaborating centers dealing with related matters or Regional institutions, indigenous institutions, and indigenous organizations. There is an obvious need to collect basic information on the living conditions and health status of indigenous peoples in selected localities, so that, on the one hand, it will be possible to document and evaluate the current situation, and, on the other, to promote the exchange and circulation of information about indigenous health, traditional medicine, active projects and studies underway, etc.

The implementation of projects under the Indigenous Peoples' Health Initiative, at the local health system level in the selected countries, will involve indigenous organizations as participants, as well as human resources from the official and informal health sector. These are the levels where it is expected to formulate policies and strategies under the Indigenous Peoples' Health Initiative, as well as to experiment with different primary health care models that are socially relevant and culturally sensitive, and which promote the articulation of local resources, based on the principles of more equitable distribution and full community participation, which are at the heart of the local health system strategy.

REFERENCES

Coloma, C. Health Situation of the Indigenous Peoples of Latin America. (Background document. Workshop '93, Winnipeg, Canada).

Dion Stout, M. Indigenous Peoples and Health in North America. (Background document. Workshop '93, Winnipeg, Canada).

de Koning, H.W. (Editor). La Salud Ambiental y la Gestión de los Recursos de Agua Dulce en las Américas. Programa de Salud Ambiental, OPS. Washington, D.C., enero de 1992.

SSA-INI (Secretaría de Salud-Instituto Nacional Indigenista). La Salud de los Pueblos Indígenas en México. México, D.F., 1992.

PAHO/WHO. Resolution XV of the XXXIII Meeting of the Directing Council. Progress Report. Development and Strengthening of Local Health Systems in the Transformation of National Health Systems. HSD/SILOS-10. Washington, D.C., 1991.

Paganini, J.M. y Capote Mir, R. (Eds.). Los Sistemas Locales de Salud. Publicación Científica No. 519. OPS, Washington, D.C., 1990.

OPS/OMS. Participación Social en la Producción de la Salud. Desarrollo y Fortalecimiento de los Sistemas Locales de Salud. HSS/SILOS-26. Washington, D.C., 1993.

PAHO. Strategic Orientations and Program Priorities, 1991-1994. Washington, D.C., 1991.

PAHO. 18th Meeting of the Subcommittee on Planning and Programming, Provisional Agenda Item 8, Document SPP 18/7, Washington, D.C., 8-10 April 1992.

OPS. Ministerio de Salud República de Colombia, Promoción de la Salud y Equidad. Declaración de la Conferencia Internacional de Promoción de la Salud. Santa Fé de Bogotá, Colombia, 9-12 November 1992.

PAHO/HSP. Health of Indigenous Peoples in the Region of the Americas. Working Document. Washington, D.C., April 1993.

WHO. Health and Welfare Canada. Canadian Public Health Association. Ottawa Charter for Health Promotion. International Conference on Health Promotion, Ottawa, Canada, November 1986.

PAHO. Health and Development of Women of Indigenous Peoples and Ethnic Groups (Subregional Project). Central American Health Initiative, Washington, D.C., September 1992.

PAHO. Health Conditions in the Americas. Scientific Publication No. 524, (Vol. I), Washington, D.C., 1990.

Rozental, M. Community Health and Development Process. Northwest Saskatchewan. Saskatchewan Indian Federated College Journal. Volume 4, No. 2, pp. 115-136, 1988.

Wolf, E.R. Europe and the People without History. U. of California Press. Berkeley, CA., 1990.

**WORKSHOP ON INDIGENOUS PEOPLES
AND HEALTH**

(Winnipeg, 13-18 April 1993)

RECOMMENDATIONS ¹

The recommendations of the Winnipeg meeting are as follows:

1. In response to the alarming health situation confronting some Indigenous peoples of the continent, the workshop recommends that PAHO and the Member Governments take immediate action to identify the priority areas and the neediest populations, in order to then declare a state of emergency in specific areas or communities. This should lead to special and priority attention to health problems and the improvement of living conditions at the local level. It is necessary to define at the country level, in consultation with the affected populations, strategies and programs for intervention designed to address the most urgent health problems, as well as to plan medium- and long-term actions.
2. PAHO and the Ministries of Health should establish a surveillance system to track the living conditions and health status of Indigenous peoples. Specific methods and indicators, as well as epidemiological instruments, should be developed for assessing this situation continuously and systematically. Mechanisms should be established that will enable Indigenous peoples to participate in deciding the kind of information to be gathered and the use it will be given.
3. The development of health projects and programs in Indigenous communities should be based on maximum and appropriate use of local resources and on active and systematic community participation in the process of planning, execution, and evaluation of these activities. PAHO and the Member Governments should ensure that in the execution of health programs and projects in Indigenous communities direct coordination is established with these communities and with grass-roots organizations. These programs should help to strengthen the self-government and autonomy of the Indigenous peoples.

¹ It should be noted that these recommendations were drafted at the conclusion of the workshop in Winnipeg. Although the original wording and the order assigned by the participants may have been changed here, the basic content has not been modified. At the time this document was drafted, the final version of the recommendations had not been ratified as the six-week period the delegates had been given to examine them and submit comments or corrections had not yet elapsed.

4. Special priority should be given to women's health programs. Migrant Indigenous populations and those in border areas, particularly those subject to forced migration, should be the target of special health policies and programs.

5. PAHO and the countries will promote dialogue and exchange between programs of primary health care with the local health system strategy in areas with Indigenous populations for the purpose of adjusting the content of the programs to local cultures and adapting the activities to the real needs of the communities at the local level. This adjustment and adaptation should be done with the direct participation of the Indigenous peoples.

6. All health projects and programs in Indigenous areas should be built on respect for the culture, values, and traditions of the Indigenous peoples involved and should acknowledge geographic and social distinctions between communities.

7. In the definition of a new health care model, PAHO and the Member Governments should recognize that culture and intercultural factors are key determinants of the living conditions and the health of individuals and communities.

8. The countries should make an effort to train non-Indigenous health workers who serve in Indigenous areas to develop favorable attitudes of understanding and respect for the local culture, beliefs, and traditional medical practices. Health workers in remote and hard-to-reach areas should receive incentives and opportunities for adequate training and education.

9. Universities, centers of study and training, and the public sector should ensure quotas so that members of Indigenous communities have access to training programs for health workers. The criteria and the educational profiles of the candidates will be established with the participation of the Indigenous communities. In the training of professionals and health auxiliaries, emphasis should be placed on intercultural studies (cross-cultural health and medicine) in the curriculum.

10. International institutions and governmental agencies should recognize and utilize the experience of Indigenous peoples in the conservation and management of the environment and natural resources. A joint effort should be made to defend Indigenous lands, improve housing and nutrition, protect the environment, and prevent environmental pollution.

11. PAHO and the Member Governments should strengthen policies for natural resource conservation and control at the country level, establishing the necessary regulations in order to prevent exploitation and destruction of medicinal plants and substances by pharmaceutical corporations and other interests, and to preserve biodiversity.

12. PAHO should actively promote regional and local initiatives that seek to articulate traditional and western medicine, promoting the exchange of experiences, as well as broader dissemination of information on the implementation and results of such initiatives. These exchanges should involve not only investigators, but also primary care workers and traditional healers or therapists and Indigenous organizations.

13. It is of fundamental importance that the countries, with the technical assistance and support of PAHO, initiate the revision of health legislation with a view to recognizing Indigenous values and traditional medical practices, seeking at the same time to limit or eliminate repressive or punitive measures against traditional healers or therapists.

14. PAHO should provide support and technical cooperation for the generation of knowledge and its dissemination in the field of Indigenous medicine and health, through collaborative research projects between countries, promoting and supporting meetings, networking, symposia, and special publications.

15. It is important that PAHO promote and support participatory research related to the health of Indigenous peoples. Research topics should be determined jointly with the local populations. Both the process and the results of research should be shared with Indigenous peoples at the local, regional, and national levels.

16. No research or action related to the health of Indigenous peoples should be promoted unless it is planned and conducted with the participation of Indigenous communities at all stages of the process. Research on Indigenous health should adhere to the international codes of ethics in force. In research on Indigenous peoples, a mechanism should be provided that ensures control by the affected populations.

17. The participants desire to emphasize the need to develop legislation that recognizes Indigenous health needs and entitlements and provides for adequate enforcement. It must also be recognized that without political will, legislation is useless.

18. Finally, the participants in the Workshop call on the international and intergovernmental community to do the following:

- i) PAHO is requested to include the issue of Indigenous health on the agenda of the next meeting of the Ministers of Health of the Americas, as well as to propose its inclusion on the agenda of the next World Health Assembly.
- ii) The United Nations task force on Indigenous populations is encouraged to promote the International Declaration of Rights of Indigenous Peoples.

- iii) The Governments of the Region are urged to ratify and implement Convention 169 of the ILO.
- iv) The Governments are also encouraged to implement the recommendations contained in Chapter 26 of Agenda 21, which emanated from the United Nations Conference on Environment and Development, held in Rio de Janeiro.
- v) The Member Governments are urged to adopt the resolutions emanating from the UNICEF World Summit for Children, in particular those relating to the health of Indigenous children.
- vi) The countries and the concerned agencies (e.g. UNESCO) are encouraged to take the necessary measures to restore, protect, and preserve the sacred Indigenous sites in order to prevent the loss of the cultural identity of Indigenous peoples.
- vii) Finally, the participants at this Workshop strongly urge the international agencies, nongovernmental organizations, institutions, and governments to mobilize the economic resources necessary for the implementation of the recommendations emanating from the Workshop.

FOLLOW-UP

Several recommendations were made with a view to following up the resolutions adopted by the Workshop. Noteworthy among them are the following:

- a) A commission of Indigenous delegates (from North, Central, and South America) to the Workshop should be formed to follow up on the actions agreed upon and support PAHO's efforts and those of the Governing Bodies of the Organization and to ensure that the recommendations are presented at the next Meeting of Ministers of Health of the Americas.
- b) Workshop delegates from each country should present the recommendations to the Ministers of Health in their respective countries in order to support PAHO's work in this regard. In addition they will be responsible for presenting these recommendations at regional and world forums (e.g., the United Nations Conference on Human Rights).
- c) A task force should be established to support and follow up on the actions taken in different countries on Indigenous health.

- d) **Indigenous lawmakers around the Region should be called on to put forward legislation in support of Indigenous health.**
- e) **An information network should be established to help maintain contact between delegates (and other interested parties), with representation in every country.**

ESTIMATED INDIGENOUS POPULATION IN THE AMERICAS
Countries and Selected Territories
(by millions of inhabitants)

| | COUNTRY | NATIONAL POPULATION | INDIGENOUS POPULATION | % |
|----------------|--------------------|---------------------|-----------------------|------|
| More than 40% | Bolivia | 6.9 | 4.9 | 71 |
| | Guatemala | 8.0 | 5.3 | 66 |
| | Peru | 20.0 | 9.3 | 47 |
| | Ecuador | 9.5 | 4.1 | 43 |
| From 5% to 20% | Belize | 0.15 | 0.029 | 19 |
| | Honduras | 4.8 | 0.70 | 15 |
| | Mexico | 85.0 | 12.0 | 14 |
| | Chile | 12.0 | 1.0 | 8 |
| | El Salvador | 5.5 | 0.4 | 7 |
| | Guyana | 0.8 | 0.045 | 6 |
| | Panama | 2.2 | 0.14 | 6 |
| | Suriname | 0.5 | 0.03 | 6 |
| | Nicaragua | 3.5 | 0.16 | 5 |
| From 1% to 4% | French Guiana | 0.1 | 0.004 | 4 |
| | Paraguay | 3.5 | 0.10 | 3 |
| | Colombia | 30.0 | 0.60 | 2 |
| | Venezuela | 18.0 | 0.40 | 2 |
| | Jamaica | 2.4 | 0.048 | 2 |
| | Puerto Rico | 3.6 | 0.072 | 2 |
| | Canada | 25.0 | 0.35 | 1 |
| | Costa Rica | 2.9 | 0.03 | 1 |
| | Argentina | 31.9 | 0.35 | 1 |
| | From 0.01% to 0.9% | Brazil | 140.0 | 0.3 |
| United States | | 245.0 | 1.6 | 0.65 |

Inter-American Development Bank, Preliminary Project for the Creation of the Development Fund for the Indigenous Peoples of Latin America and the Caribbean, Washington, D.C., 1991;
Inter-American Indigenous Institute, Data Bank, Mexico, 1992.

Principal Health and Disease Figures in the Indigenous
Population in Selected Countries of the Southern Cone,
According to Various Studies

Table 1. Principal data of health and diseases of indigenous peoples in selected countries of Cone South.

| Country | Region or indigenous people | Date | Pathology or indicator | Rates | Type | Total Country |
|--------------|--|--------------------------|---------------------------|--------------|---------------------|----------------------|
| ARGENTINE(1) | Matacos, Chorotes, Chunupis and Tobas | 1979-1980 | Respiratory diseases | 37.90% | | |
| | | | Gastroenteric diseases | 15.00% | | |
| | | | Skin infections | 8.70% | | |
| | | | Gynaecological and | 14.80% | | |
| | | | Muscle-skeletal disorders | 5.90% | | |
| | | | Other diseases | 17.70% | | |
| CHILE (2) | Aymara | end of 1970 to early '80 | Chagas | 14.10% | Index house infect. | |
| | | | | 12.50% | Serology + | |
| | Atacameños | early '80 | | 33.80% | Index house infect. | |
| | | | | 10.00% | Serology + | |
| | Aymara | early '80 | Hydatidosis | 100 | per 100,000 | 7 |
| | Pehuenche (Community of Lonquimay) | | | 48 | per 100,000 | 7 |
| | Pehuenche (women's population) | 1968 | Bocio | 50% | | |
| | Aymara (women's population of Parinacota) | 1990 | Syphilis | 5% | | per 100,000 inhab. 7 |
| | | 1990 | Candidiasis | 26.30% | | |
| | | 1990 | Trichomoniasis | 3.70% | | |
| | Rapa-nui | 1992 | Leprosy | 26.10% | Serology + | |
| | Mapuche (n=405) (3) | 1991 | HTLV-1 | 0.70% | Serology + WB or | |
| | | 1991 | HIV-1 | 0,00% | Serology + ELISA | |
| | Mapuche (Indigenous communities selection) | 1975-1980 | Life expectancy | 59 | years | |
| 1988 | | | 63 | years | 71 | |
| 1985 | | Infant mortality | 45 | per thousand | 20 | |

1 Brett, Mary T. "Primary care and the Pattern of Disease in a Rural Area of the Argentina Chaco" PAHO Bulletin, vol 18, n° 2, 1984 Washington

2 Medina Cárdenas, Eduardo "Pueblos Indígenas y Salud en Chile" Organización Panamericana de la Salud, Santiago de Chile 1992

3 Inostroza, J., Díaz, P and Saunier, C. "Prevalence of Antibodies to HTLV-1 in South American Indians (Mapuches) from Chile" in Scandinavian Journal of Infectology Diseases 23 507 508, 1991

Table 2. Principal data of health and diseases of indigenous peoples in selected countries of the Andean region.

| Country | Region or indigenous people | Date | Pathology or indicator | Number | Rates | Type | Tot. Country | |
|-------------|--|--------------------|------------------------|----------------|------------|---------------------|------------------|--------|
| BOLIVIA (1) | | 1986 | TBC | | | New cases | 5941 | |
| | Dept. Santa Cruz and Chuquisaca | 1980-1983 | Chagas | | 100% | Index house infect. | 26.30% | |
| | (Population in risk) | | | | 30 to 45% | Serology + | | |
| | Dept. Beni, Sta.Cruz, Tarija and Chuquisaca | 1981-1984 | Malaria | 47,222 | | Registered cases | | |
| | Dept. Beni, Sta.Cruz, Cochabamba, and La Paz | 1981-1984 | Yellow fever | 213 | | Registered cases | | |
| | Dept. La Paz | 1981-1984 | Bubonic | 78 | | Registered cases | | |
| | La Paz (city), Cochabamba and Santa Cruz | | 1982-1984 | Human rabies | 26 | | Registered cases | |
| | | | 1970-1980 | Silicosis | | | Cases/year | 12,000 |
| | | | | | | | Deaths/year | 363 |
| | | | 1976 | Fertility rate | | | | 6.7 |
| | 1980-1985 | Maternal mortality | | | per 10,000 | 48 | | |

1. Pedersen, D., Betts C., Mariscal, J., Torres Goitfa, J., "Supervivencia Infantil en Bolivia, Situación actual y prioridades para la acción", Informe de Consultoría, USAID, La Paz, 1987.

(continued)

| | | | | | | | |
|--|---------|-----------|---|-------|--------|-------------------|--------|
| | | 1982 | Births unregistered | | | | 57.60% |
| | | 1982 | Deaths unregistered | | | | 77.40% |
| | | 1970-1975 | Infant mortality (projection of CELADE) | | | per 1,000 | 151% |
| | | 1985-1990 | | | | | |
| | Quechua | 1976 | Infant mortality according to maternal language | | 277.7% | Quechua | 153% |
| | Aymara | 1976 | | | 238.8% | Aymara | |
| | | | | | 186.8% | Spanish and other | |
| | | | | | 125.6% | Spanish | |
| | | 1981 | Causes of death in < 5 years: | | | | |
| | | | Gastroenteric diseases | 2,257 | | | 19% |
| | | | Respiratory diseases | 2,211 | | | 36.70% |
| | | | Affections originated in the neonatal period | 1,664 | | | 14% |
| | | | Others bacterial diseases | 1,020 | | | 8.60% |
| | | | Viral diseases | 743 | | | 6.20% |
| | | | Undefined signs and symptoms | 1,223 | | | 10.30% |

Table 2. Principal data of health and diseases of indigenous peoples in selected countries of the Andean region. (continued)

| Country | Region or indigenous people | Date | Pathology or indicator | Number | Rates | Type | | Total Country | |
|---------|--------------------------------|--|------------------------------|--|-------|--|------------------------------|---------------|--|
| | | | | | | | | | |
| ECUADOR | Quichua (Community of Ilumán) | 1978 | Undefined signs and symptoms | | 58% | Percentage of total morbidity reported 2 weeks before survey | Rates per 10,000 inhabitants | 78.42 | |
| | | | woman's population | Diseases of S.N.C. and of sensory organs | | | | 18% | |
| | | | | Gastroenteric diseases | | | | 7.50% | |
| | | | children's population | Infectious and parasitic diseases | | | | 42% | |
| | | | | Respiratory diseases | | | | 18% | |
| | | | | Undefined signs and symptoms | | | | 24% | |
| | Quichua (Community of Imantag) | woman's population | | Undefined signs and symptoms | | 41% | | | |
| | | | | S.N.C. and org. of sense diseases | | 15% | | | |
| | | | | Mental troubles | | 10% | | | |
| | | children's population | | Infectious and parasitic diseases | | 48% | | | |
| | | | | Respiratory diseases | | 22% | | | |
| | | | | Undefined signs and symptoms | | 15% | | | |
| | | (1) Quichuas of Chimborazo and Pichincha | 1978-1980 | Intestinal parasites | | | | | |
| | | | | Nematodes: | | | | | |
| | | | | Trichuris trichiura | 51 | 23% | | | |
| | | | Ascaris lumbricoides | 129 | 58% | | | | |

(continued)

| | | | | | | | |
|--|--|--|----------------------------------|----|-----|--|--|
| | | | Strongyloides stercoralis | 3 | 1% | | |
| | | | Cestodes: | | | | |
| | | | Hymenolepis spp. | 2 | <1% | | |
| | | | Protozoos: | | | | |
| | | | Entamoeba histolytic | 69 | 31% | | |
| | | | Entamoeba coli | 63 | 28% | | |
| | | | Balantidium coli | 5 | 2% | | |

I. Peplow, Daniel "Parásitos intestinales en la población de varias regiones del Ecuador: estudio estadístico" Boletín de la Oficina Sanitaria Panamericana, 93 (3) 1982.

Table 3. Principal pathologies diagnosed in Yanomami area of Amazon region (Brazil), for indigenous peoples.

| Population of study | Paapiú 846 | Catrimani 1728 | Demini 925 | Ajuricaba 59 | Tootobi 1373 | Marari 483 | Marauá 127 | Surucucu 3210 | Xidéa 2029 | Homoxi 2501 | Parafuri 366 | Auarais 2764 | Waikás 159 | Ericó 239 | Palmiú 494 | Mujacai 837 | Total 18 140 |
|------------------------|---------------|-------------------|---------------|-----------------|-----------------|---------------|---------------|------------------|---------------|----------------|-----------------|-----------------|---------------|--------------|---------------|----------------|-----------------|
| I R A.: | | | | | | | | | | | | | | | | | |
| Light | 69 | 349 | 112 | - | 107 | 50 | - | 157 | 119 | 128 | 16 | 153 | 1 | 39 | 68 | 159 | 1 527 |
| Moderate | 2 | 51 | 15 | 1 | 1 | 8 | 50 | 6 | 15 | 38 | - | 78 | - | 1 | 2 | 7 | 275 |
| Grave | 9 | 18 | 2 | - | 1 | 3 | - | 32 | - | 12 | - | 10 | - | - | - | 1 | 88 |
| Gastroenteritis | 26 | 154 | 118 | 3 | 6 | 87 | 2 | 61 | 225 | 80 | 116 | 13 | 20 | 36 | 2 | 61 | 1 010 |
| Intestinal | 85 | 227 | 59 | 13 | 88 | 33 | 12 | 577 | 221 | 253 | 73 | 211 | 7 | 38 | 6 | 229 | 2 132 |
| Parasites | | | | | | | | | | | | | | | | | |
| Skin. | | | | | | | | | | | | | | | | | |
| Scabies | 121 | 6 | 58 | 24 | 29 | 46 | 8 | 57 | 184 | 120 | 59 | 12 | 13 | 13 | 9 | 8 | 767 |
| Infections | 24 | 55 | 36 | - | 13 | 14 | 1 | 130 | 39 | 53 | 9 | 3 | 3 | 9 | 5 | 25 | 419 |
| Others | 32 | 11 | 68 | 4 | 69 | 25 | 5 | 16 | 41 | 8 | - | 28 | - | 10 | 10 | 41 | 368 |
| Conjunctivitis | 12 | 66 | 175 | - | 104 | 138 | 17 | 121 | 173 | 65 | 18 | 70 | 6 | 31 | 2 | 101 | 1 099 |
| Injured/ traumatism | - | 104 | 58 | - | 13 | 2 | - | 27 | 29 | 54 | 25 | 8 | 6 | 2 | 2 | 8 | 338 |
| Leishmaniosis | 2 | - | - | - | 7 | - | - | 10 | - | 5 | - | 6 | - | 3 | - | 2 | 35 |
| TBC | 1 | 2 | 1 | - | 6 | 1 | 18 | - | 1 | 18 | - | 2 | - | 4 | 6 | 6 | 66 |
| Onchocercosis | - | 1 | - | - | 7 | - | - | 2 | - | 7 | - | 3 | 8 | - | - | - | 28 |
| Others | 6 | 22 | 259 | 13 | 134 | 85 | 28 | 30 | 152 | 109 | 7 | 13 | 8 | 15 | 2 | 81 | 964 |
| Malaria | 308 | 228 | 34 | 15 | 531 | 54 | 44 | 886 | 1002 | 1447 | 362 | 987 | 124 | 48 | 180 | 243 | 6 493 |
| Anemia | 237 | 4 | 2 | 4 | 172 | - | - | 344 | 132 | 250 | - | 112 | - | 9 | 4 | 10 | 1 280 |
| Esplenomegal | 311 | - | - | 16 | 510 | - | 9 | 50 | 23 | - | - | 2 | - | - | 58 | - | 979 |
| Malnutrition. | | | | | | | | | | | | | | | | | |
| Moderate | 96 | 20 | - | 2 | 27 | 4 | 1 | 288 | 35 | 159 | 71 | 110 | 1 | 2 | 8 | 2 | 826 |
| Grave | 6 | - | - | - | - | - | - | 36 | 3 | 34 | 8 | 2 | - | - | - | - | 89 |

Source "Primeiro Relatório do Distrito Sanitário Yanomami. Avaliação das Atividades e Diagnóstico de Saúde" Fundação Nacional de Saúde, Brasília 1991

161

Table 4. Principal pathologies diagnosed in indigenous peoples of the Amazon region or tropical forest in South-American countries.

| Country | Region or indigenous peoples | Date | Pathology or indicator | Number | Rates | Type | Total Country |
|-----------------|---|-----------|---|--------|---|--|--------------------------------|
| PARAGUAY | Chamococos | 1984 | Tuberculosis (n=201) Parasitosis (n= 650) Diarrhoea (n=656) Anemia (n= 655) Syphilis | | 2% 15.69% 10.36% 8.70% 4.50% | Baciloscop. + Cases + VDRL + | 0.52%o(3) 0.37%o(3) |
| | Angaité, Lengua, Sanapaná, Tobas-Maskoy | | Tuberculosis (n=104) Parasitosis (n= 592) Diarrhoea (n=596) Anemia (n=621) Syphilis | | 6.50% 31.25% 6.54% 19.64% 1.60% | Baciloscop. + Cases + VDRL + | 0.52%o(3) 0.37%o(3) |
| ECUADOR (2) | Communities of Napo and Pastaza (n= 711 included settler population) | 1978-1980 | Intestinal Parasites | | | | |
| | | | Nematodes: | | | | |
| | | | Trichuris trichiura | 639 | 90% | | |
| | | | Ascaris lumbricoides | 445 | 63% | | |
| | | | Ancilostoma duodenalis | | | | |
| | | | Necator americanus | 246 | 35% | | |
| | | | Strongyloides stercoralis | 57 | 8% | | |
| | | | Enterobius vermicularis | 5 | < 1% | | |
| Capillaris spp. | 1 | < 1% | | | | | |
| Cestodes: | | | | | | | |

(continued)

| | | | | | | | |
|--|--|--|----------------------|-----|------|--|--|
| | | | Taenia spp. | 5 | < 1% | | |
| | | | Hymenolepis spp. | 2 | < 1% | | |
| | | | Protozoa: | | | | |
| | | | Entamoeba histolytic | 135 | 19% | | |
| | | | Entamoeba coli | 52 | 7% | | |
| | | | Balantidium coli | 61 | 9% | | |
| | | | Chilomastix mesnili | 22 | 3% | | |
| | | | Giardia lamblia | 113 | 16% | | |
| | | | Trichomonas hominis | 2 | < 1% | | |

1 Benítez, P., Oddone, H., Rivarola, D., Cabral, R. "La sociedad nacional y las comunidades indígenas. estudio de la educación y la salud en cuatro comunidades del Alto Paraguay y actitudes de la población nacional hacia los indígenas" Instituto Paraguayo del Indígena, Asunción 1985.

2 Peplow, Daniel "Parásitos intestinales en la población de varias regiones del Ecuador: estudio estadístico" Boletín de la Oficina Sanitaria Panamericana, 93 (3) 1982

3 Rate of 1986. "Las Condiciones de Salud de las Américas", Publicación Científica N° 524, vol. II, Organización Panamericana de la Salud, Washington 1990.

(continued)

Table 4. Principal pathologies diagnosed in indigenous peoples of the Amazon region or tropical forest in South-American countries.
(continuation)

| Country | Region or indigenous peoples | Date | Pathology or indicator | Number | Rates | Type | Total Country | |
|-----------|--|------|--|--------|-------|--------------|---------------|--|
| VENEZUELA | Territorio Federal de Amazonas: Yanomami (28.49%), Guajibo Piaroa (20.62%), Yequana (8.91%), Curripaco (4.76%), Bare (3.71%), Baniva (3.42%), Piapoco (1.88%), Saxema (6.94%), Puinave (1.44%), Warekena (0.93%), Hotis (1.17%), Yaberana (0.45%), Panare (6.98%), Others (1.04%), Non specified (2.24%) | 1986 | Birth rate | 1357 | 18.4 | per thousand | 28.3 | |
| | | 1987 | General mortality | | 3.4 ‰ | | 4.4‰ | |
| | | 1984 | Diagnostics in Hospital and "Medicaturas Rurales" | | | | | |
| | | | Catarrh | 2155 | | | | |
| | | | Tonsillitis | 950 | | | | |
| | | | Helminthiasis | 713 | | | | |
| | | | Anemias | 666 | | | | |
| | | | Gastroenteritis | 593 | | | | |
| | | | Gastroduodenal | 540 | | | | |
| | | | Injuries | 323 | | | | |
| | | | Mycosis | 110 | | | | |
| | | | Piodermatitis | 108 | | | | |
| | | | Dysentery | 92 | | | | |
| | | | Bronchitis | 65 | | | | |
| | | | Diarrhoea | 61 | | | | |
| | | | Allergies | 54 | | | | |
| | Accidents | 48 | | | | | | |

1. "Informe General de Pasantías: Territorio Federal de Amazonas" Escuela de Nutrición y Dietética, Facultad de Medicina, Universidad Central de Venezuela, abril/julio 1989

Table 5. Principal data of health and diseases of indigenous peoples in selected countries of the Mesoamerica region.

| Country | Region or indigenous peoples | Date | Pathology or indicator | Rates | Type | Total Country | |
|-----------|------------------------------|----------------------|------------------------|-----------|--------------|---------------|--------|
| Guatemala | Indigenous peoples | 1981 | Infant mortality (1) | 106 | per thousand | 87 (2) | |
| | | 1986 | | 77 | per thousand | 70 (2) | |
| | | 1985 | Infant mortality (3) | 100 - 150 | per thousand | | |
| | | 1990 | Fertility rate (1) | 6.8 | | 4.4 (2) | |
| | | 1989 | Maternal mortality (1) | | estimated | 20.2 | |
| | | 1987 | General mortality(4): | | | | |
| | | | Household | | | | 75.60% |
| | | | Hospital | | | | 19.10% |
| | | | Street | | | | 4% |
| | | | Institutional homes | | | | 1.30% |
| | | | Without medical care | | | | 67.20% |
| 1987 | With medical care | | | | 32.80% | | |
| | With certification (5) | | | | 92.90% | | |
| 1987 | Without certification | | | | 7.10% | | |
| | 1985-1990 | Life expectancy (1): | | . (6) | | | |

(continued)

| | | | | | |
|--|--------------------|-------|--|--------|---------------------------|
| | | Men | | years | 59.7 |
| | | Women | | years | 64.4 |
| | Santa María Cauqué | 1974 | Citomegalovirus (n= 109 children) (7) | 46.80% | in the first year of life |
| | north zone | | Malaria (4) | 63.20% | Percentage of positivity |
| | south zone | | | 22.90% | |
| | | 1988 | Causes of infant mortality(4): | | |
| | | | Diarrhoea | | 23.80% |
| | | | Respiratory diseases | | 23.60% |
| | | | Malnutrition | | 4.30% |
| | | | Affections originated in the neonatal period | | 4.10% |
| | | | Hydroelectrolitic imbalance | | 3.20% |

1 "Análisis de la situación del niño y la mujer", UNICEF / SEGEPLAN, Guatemala 1991.

2. Totals in non indigenous population

3. Minkowski, W. "Mayan Indian Health in Guatemala" West Journal Medicine 148: 474-476, 1988.

4. "Las Condiciones de Salud en las Américas" Vol II: 162, Organización Panamericana de la Salud 1990.

5. Nearly 50% had medical certification, 34,1% from municipal authorities and 10,3% for personnel outside the health sector.

6 Minkowski thinks the life expectancy for the indigenous people is 10 to 15 years less than the national average.

7. Cruz, J ; Mata, L. y Urrutia, J. "Citomegaloviruria durante el primer año de vida: estudio prospectivo en una población indígena de Guatemala" Boletín Oficina Sanitaria Panamericana 83 (3). 218-222, 1977

(Continued)

Table 5. Principal data of health and diseases of indigenous peoples in selected countries of the Mesoamerica region. (continuation)

| Country | Region or indigenous people | Date | Pathology or indicator | Rates | Type | Total Country |
|-----------------------------------|-----------------------------|------|--|--------|--|---------------|
| Mexico | | 1988 | Growth of population | | Percentage / year | 1.80% |
| | | 1990 | Life expectancy (1) | | years | 70 |
| | | 1984 | Infant mortality | | | 29.16% |
| | | 1985 | Maternal mortality (2) | | | 0.6% |
| | | 1983 | General mortality | | | 5.1 |
| | Rural marginal zones (3) | 1986 | Digestive system diseases (enteritis, amebiasis, helminthiasis) | 38.20% | Percentage of total communicable diseases notified | 22.47 |
| | | | Respiratory system diseases (IRA, influenza and pneumonia) | 49.70% | | |
| | | | Tuberculosis | 25 | Cases per 100,000 inhab. | |
| | | | Malaria | 69,7 | | |
| | | | Dengue | 10,26 | | |
| | | | Trachoma | 3,44 | | |
| | | | Onchocercosis | 0,29 | | |
| | | | High pressure blood | 275 | | |
| | | | Diabetes | 148 | | |
| | | | Rheumatic fever | 3,4 | | |
| | | | Cirrhosis hepatic | 16,2 | | |
| | | | Scabies | 560 | | |
| Whooping-cough | 1,39 | | | | | |
| Neonatal tetanus | 0,36 | | | | | |
| Tetanus | 0,21 | | | | | |
| Measles | 8,28 | | | | | |
| Accident, poisoning, and violence | 1.582 | | | | | |

1 "Estado Mundial de la Infancia 1992". UNICEF

2 "La Situación de la Mujer pobre en México" UNICEF México 1990

3 "Diagnóstico de salud en las zonas marginadas rurales de México" Instituto Mexicano del Seguro Social, México 1986

4 Rates per total inhabitants registered for Social Security in the "Programa de Solidaridad Social por Cooperación Comunitaria IMSS-COPLAMAR"

Table 6. Prenatal and childbirth care for regions or indigenous people, in countries of the Region.

| Country | Region or indigenous people | Date | Pathology or indicator | Number | Rates | Type | Total Country |
|-------------|--|------|--|--------|--------|--|---------------|
| Chile (1) | Mapuche | 1988 | Childbirth care: | | | | |
| | | | Medical and traditional birth attendants | | 69% | | |
| | | | Machi or healers | | 23% | | |
| | | | Person without qualification | | 7% | | |
| | | | Rural health auxiliary | | 1% | | |
| Bolivia (2) | Total Country | 1982 | Institutional childbirth | 47.760 | | Percentage | 18.90% |
| | | | (Health Ministry and Social Security) Home childbirth | | | distribution of childbirth care | 80% |
| Peru(3) | Womens in rural area (12 to 49 years old) . Highland sierra | 1984 | Prenatal care: | | | Percentage | |
| | | | Hospital | | 6.20% | distribution of prenatal care by place | 41.90% |
| | | | Health center | | 1.60% | | 1.70% |
| | | | Consulting room, clinic | | 0.70% | | 8.10% |
| | | | Household | | 90.80% | | 47.80% |
| | Others | | 0.70% | | 0.60% | | |
| | Forest | | Hospital | | 7.90% | | 41.90% |
| | | | Health center | | 1.40% | | 1.70% |

(continued)

| | | | | | | |
|-------------|--|------|---|---------|--|---------|
| | | | Consulting room, clinic | 1.60 % | | 8.10 % |
| | | | Household | 88.20 % | | 47.80 % |
| | | | Others | 1 % | | 0.60 % |
| Ecuador (4) | Indigenous communities quichuas. Highland sierra | 1981 | Home birth: solo, husband, family or traditional birth attendants (TBA's) | 83 % | Percentage distribution of childbirth care | |
| | :Imbabura | | Institutional: professional care | 7 % | | |
| | Bolivar | | Home birth: solo, husband, family or traditional birth attendants (TBA's) | 98 % | | |
| | | | Institutional: professional care | 2 % | | |
| | Indigenous communities quichuas. Forest: | | Home birth: solo, husband, family or traditional birth attendants (TBA's) | 96 % | | |
| | Pastaza | | Institutional: professional care | 4 % | | |

1. "Condiciones de vida de los indígenas people: Estudio realizado en reducciones mapuches seleccionadas". Universidad de la Frontera / INE / Fundación Instituto Indígena / CELADE / Programa de Apoyo y Extensión en Salud Materno Infantil, Santiago de Chile 1991.

2. Pedersen, D., Betts, C., Mariscal, J., Torres Goitia, J. "Supervivencia Infantil en Bolivia. Situación actual y prioridades para la acción" Informe de Consultoría, USAID, La Paz 1987

3. Fuente. Instituto Nacional de Estadística, e Informática, "Encuesta Nacional de Nutrición y Salud (ENNSA), Lima 1984.

4. Coloma, C.; Baruffati, V.; Chela, T. "Maternidad e infancia en los Quichuas del Ecuador" CIEI, Universidad Católica del Ecuador, Quito 1981.

(continued)

Table 6. Prenatal and childbirth care for regions or indigenous people, in countries of the Region.

| Country | Region or indigenous people | Date | Pathology or indicator | Number | Rates | Type | Total Country |
|---------------|--|------|--|--------|--------|---------------|---------------|
| Guatemala (1) | Rural areas | 1989 | Childbirth care: Traditional birth attendants | | 77% | | |
| | Maya area | | Traditional birth attendants | | 90% | | |
| | Rural areas | | Prenatal care: | | | | 38% |
| Mexico | Total Country (2) | 1987 | Prenatal care | | | | 62% |
| | | | Childbirth care (institutional) | | | | 38% |
| | | | Childbirth care (traditional birth) | | | | 14% |
| | Rural population (3) | 1986 | Childbirth care in rural hospital | 39,949 | 22.70% | Percentage of | |
| | | | Consultation for pregnancy in women under 15 years old | 2,015 | 15.70% | | |
| | | | Perinatal deaths during pregnancy | | | 87.40% | |
| | | | After childbirth care | | | 3.70% | |
| | Maternal mortality: | | | | | | |
| | After childbirth hemorrhages | | | 16.99% | | | |
| | Other complications of childbirth | | | 15.44% | | | |
| | High pressure blood | | | 8.11% | | | |
| | Puerperal sepsis | | | 6.95% | | | |
| | Other foetal and placenta problems | | | 6.56% | | | |
| | Retention of placenta and/or membranes without hemorrhages | | | 5.41% | | | |
| | Other maternal causes | | | 40.54% | | | |

1 "Análisis de la situación del niño y la mujer", UNICEF / SEGEPLAN, Guatemala 1991.

2. "La situación de la mujer pobre en México" UNICEF, México 1990.

3. "Diagnóstico de salud en las zonas marginadas rurales de México" Instituto Mexicano del Seguro Social, México 1986.

Table 7. Nutritional situation in region or indigenous population, in country studies.

| Country | Region or indigenous people | Date | Indicator | Rates | Type | Total |
|------------|---|------|---|--------------------------------------|--|---------------|
| Guatemala | Rural area | 1989 | Feminine goitre | 22.60% | Prevalence in pregnant women in non-pregnant women | |
| | | | Iron deficit | 48 % 21 % | | |
| | Sololá Totonicapán | 1978 | Weight deficit | 40.60% | indigenous children | 37.40% |
| | | | Height deficit | 71.60% | indigenous children | |
| | Totonicapán, Jalapa, Zacapa, Escuintla, Amatitlán, Santa Rosa and Baja Veracruz | 1990 | Malnutrition moderate and severe. under 5 years old (n= 74,000) Moderate and severe | 64.60% 60.50% 76.10% 41.20% | in school children 6 to 9 years According to Waterlow classification | |
| Mexico (2) | Rural area | 1989 | Malnutrition in children under 5 years old | 41.50% | | |
| | Zonas Mixteca and Cañada | | Malnutrition children <5 years old Malnutrition chronic | 66.40% 5.40% | | |
| | Total country | | Status nutritional (perimeter mesobraquial-height-age <5 years | | Percentage difference between areas well and poorly nourished | 358.80% |
| | Rural area | | Average consumption per capita Kilocalories Proteins | | kcal./day grams/day | 1,880 59.2 |
| | | | Proteins of animal origin | | Percentage difference between areas well and poorly nourished | 111% |

1. "Análisis de la situación del niño y la mujer", UNICEF/ SEGEPLAN, Guatemala 1991

2. "México. Diagnóstico de la situación alimentaria y nutricional" Consejo Nacional de Alimentación, 1992.

Table 8. Water supply and sanitation in indigenous population or región of selected countries.

| Country | Region or indigenous people | Date | Indicator | Rates | Type | Total Country | | |
|-----------------|-----------------------------|----------|----------------------|-------------------------------|--------------------------|--------------------------|-----|-------|
| Chile (1) | Mapuche | 1988 | Water supply: | | Percentage of population | 85.97% (2) | | |
| | | | Open well | 36% | | | | |
| | | | Protected well | 53.40% | | | | |
| | | | Slope, river, stream | 9.30% | | | | |
| | | | Other | 0.30% | | | | |
| | | | Sewage disposal: | | Percentage of population | 88.95% (2) | | |
| | | | Boxed well | 90.20% | | | | |
| | | | Latrine | 0.50% | | | | |
| Toilet in house | 0.20% | | | | | | | |
| None | 9.10% | | | | | | | |
| Bolivia (3) | Urban (Total country) | 1976 (4) | Sewage disposal: | | Percentage of population | | | |
| | | | Sewer system | | | | 30% | |
| | | | Septic bed | | | | 4% | |
| | | | Well | | | | 14% | |
| | None | | | 52% | | | | |
| | Rural (Total country) | | | Sewage disposal: | | | | |
| | | | | Sewer system | | | | 0.30% |
| | | | | Septic bed | | | | 0.50% |
| | | | | Well | | | | 3.50% |
| | None | | | 95.70% | | | | |
| | Population total country | | | Water supply: | | Percentage of population | | |
| | | | | Access to public sewer system | | | | 39% |
| Well | | | | | 24% | | | |
| River, lake | | | | | 33% | | | |

(continued)

| | | | | | | |
|-------------|------------------|------|----------------------|--|--------------------------|-----------|
| | | | Other | | | 4% |
| Peru (5) | Rural population | 1988 | Potable water supply | | Percentage of population | 22.31% |
| | Urban population | | | | | 77.60% |
| | Rural population | | Sewage disposal | | Percentage of population | 16.60% |
| | Urban population | | | | | 55.0% (6) |
| Ecuador (5) | Rural population | 1988 | Potable water supply | | Percentage of population | 36.97% |
| | Urban population | | | | | 75.11% |
| | Rural population | | Sewage disposal | | Percentage of population | 34.19% |
| | Urban population | | | | | 75.24% |

1 "Condiciones de vida de los pueblos indígenas Estudio realizado en reducciones mapuches seleccionadas". Universidad de la Frontera/INE/Fundación Instituto Indígena/CELADE, 1991.

2 "Las Condiciones de Salud en las Américas" vol.1: 232-233, Organización Panamericana de la Salud, Washington 1990.

3. Pedersen, D , Betts, C., Mariscal, J. ,Torres Goitia, J. "Supervivencia Infantil en Bolivia. Situación actual y prioridades para la acción" Informe de Consultoría, USAID, La Paz 1987.

4. Source: Morales y Rocabado, 1987.

5. "Las Condiciones de Salud en las Américas" vol 1: 232-233, Organización Panamericana de la Salud, Washington 1990.

6. Data insufficient

(continued)

Table 8. Water supply and sanitation in indigenous population or región of selected countries. (continuation)

| Country | Region or indigenous people | Date | Indicator | Rates | Type | Total Country |
|---------------|-----------------------------|------|---------------------------------|--------|---|---------------|
| Paraguay (1) | Indigenous population total | 1981 | Water supply: | | Percentage of homes | |
| | | | Stream, river, lake | 68.50% | | |
| | | | Well, cistern | 25.40% | | |
| | | | Public tap | 5.90% | | |
| | | | Other | 0.30% | | |
| | | | Sewage disposal: | | Percentage of homes | |
| | | | Available | 62.90% | | |
| | | | Not available | 37.10% | | |
| Brazil (2) | Rural population | 1984 | Water supply: | | Percentage of population, total country | 6.30% |
| | | | Access to public sewer system | | | 18.75% |
| | | | Wells, rivers not protected | | | 8.60% |
| | | | Others | | | |
| | Rural population | | Excreta disposal | | Percentage of population, total country | 1.20% |
| | | | Access to public sewer system | | | 1.66% |
| | | | Septic bed | | | 10.56% |
| | | | Latrine | | | 1.35% |
| | | | Others (in rivers, lakes, etc.) | | | 15.07% |
| | | | Unspecified | | | |
| Guatemala (3) | Rural population | 1990 | Potable water supply | | Percentage of population, total country | 42.60% |
| | Petén | | | 15.50% | | |
| | North-west | | | 43% | | |
| | North | | | 25.10% | | |
| | Rural population | | Sanitation | | | 51.60% |

(continued)

| | | | | | | |
|------------|---------------------|------|--------------------------|---------|---------------------|---------|
| | North-west | | | 54.60 % | | |
| | South-west | | | 40 % | | |
| | North | | | 53.90 % | | |
| Mexico (4) | Marginal rural zone | 1988 | Water connection in home | | Percentage of homes | 39.80 % |
| | | | W.C. english type | | | 5.90 % |
| | | | Septic bed | | | 6.60 % |
| | | | Latrine | | | 24.70 % |

1. "Censo y Estudio de la Population Indígena del Paraguay 1981" Instituto Paraguayo del Indígena, Asunción 1982

2 Source: IBGE, "Pesquisa Nacional por Amostra de Domicílios 1984" citado en IPEA "Bases para Formulação de Políticas e Programas em Saneamento Rural" N°2, Brasília 1989.

3. "Análisis de la situación del niño y la mujer", UNICEF / SEGEPLAN, Guatemala 1991.

4. "Diagnóstico de salud en las zonas marginadas rurales de México" IMSS - Coplamar, México 1988