



LIVING SYSTEMATIC REVIEW OF THERAPEUTIC OPTIONS FOR POST-ACUTE AND POST-COVID19 CONDITION

June 16 2023

Living Systematic Review of Therapeutic Options for Post Acute or Post COVID-19
Condition

Washington, D.C.

16 June 2023

© Pan American Health Organization, 2023

Some rights reserved. This work is available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO license ([CC BY-NC-SA 3.0 IGO](https://creativecommons.org/licenses/by-nc-sa/3.0/igo/)).

Under the terms of this license, this work may be copied, redistributed, and adapted for non-commercial purposes, provided the new work is issued using the same or equivalent Creative Commons license and it is appropriately cited. In any use of this work, there should be no suggestion that the Pan American Health Organization (PAHO) endorses any specific organization, product, or service. Use of the PAHO logo is not permitted. All reasonable precautions have been taken by PAHO to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall PAHO be liable for damages arising from its use.

This document includes the results of a rapid systematic review of current available literature. The information included in this review reflects the evidence as of the date posted in the document. In recognition of the fact that there are numerous ongoing clinical studies, PAHO will periodically update this review and corresponding recommendations as new evidence becomes available.

Contents

Acknowledgments	6
Funding	6
Executive summary	7
Summary of evidence	7
P-ACC-related asthenia or fatigue	8
P-ACC-related dyspnea	12
P-ACC-related neurocognitive symptoms or sleep disturbances	14
P-ACC-related olfactory and/or gustatory dysfunction	16
P-ACC-related cardiovascular system symptoms	18
P-ACC-related psychological distress	19
P-ACC-related thromboembolic risk	20
Pediatric inflammatory multisystem syndrome associated with SARS-CoV-2 (PIMS-TS)	21
P-ACC prophylaxis	22
Changes since previous edition	24
Concluding remarks	25
Systematic review of therapeutic options for post acute or post COVID-19 condition (P-ACC)	26
Background	26
Methods	27
Search strategy	27
Study selection	27
Inclusion criteria	27
Living evidence synthesis	28
Results	31
Studies identified and included	31
Risk of bias	32
Main findings	34
P-ACC-related asthenia or fatigue	34
P-ACC-related dyspnea	39
P-ACC-related neurocognitive symptoms or sleep disturbances	42
P-ACC-related olfactory and/or gustatory dysfunction	43

P-ACC-related cardiovascular system symptoms.....	45
P-ACC-related psychological distress.....	46
P-ACC-related thromboembolic risk	47
Full description of included studies	51
References.....	102
Annex 1. Summary of findings tables	111

Acknowledgments

This document was developed by Ariel Izcovich, Martin Ragusa, Fernando Tortosa, Sasha Peiris, and Ludovic Reveiz from the knowledge Translation Program, Department of Evidence and Intelligence for Action in Health and the Incident Management System for the response to COVID-19, Pan American Health Organization (PAHO). It was strengthened with the valuable contributions of: Pedro Ordunez, Antony Duttine and Carmen Martinez from the Noncommunicable Diseases and Mental Health department, PAHO, and Sylvain Aldighieri, Deputy Director of the Health Emergencies and Incident Management System for the response to COVID-19.

Funding

This work was partially funded by the Government of the United States of America.

Executive summary

Background

Post COVID-19 condition (PCC), also known as long COVID or post-acute sequelae of SARS-CoV-2 infection (PASC), is the continuation or development of new symptoms in the period after acute infection with SARS-CoV-2. The World Health Organization (WHO) definition of PCC states that these symptoms should be present after three months of the initial SARS-CoV-2 infection and last for at least two months with no other explanation. While PASC definitions state that persistent or new symptoms need to be present 30 days after a documented SARS-COV-2 infection or the onset of COVID-19 symptoms, post COVID-19 condition or post-acute sequelae of SARS-CoV-2 infection (P-ACC) can affect anyone exposed to SARS-CoV-2, regardless of age or severity of acute infection. Many of the reported symptoms are debilitating and have a strong negative impact on mental health and quality of life. While most patients recover, some may experience multiple outcomes, with multiple organ systems affected simultaneously, including cardiovascular, mental, metabolic, renal, and others.

This review compiles the following evidence on potential therapeutic options for P-ACC. It includes all the identified clinical forms, symptoms, and manifestations of P-ACC for which an intervention was assessed in at least one randomized controlled trial (RCT). It is hoped this information will support investigators, policymakers, and prescribers navigate the flood of relevant data to ensure that management of P-ACC, at both the individual and population levels, is based on the best available knowledge. This resource will be continually updated as more research is released into the public space.

Summary of evidence

All odd numbered tables (Table ES1 to ES15) present RCTs according to the reported P-ACC related organ/system affected and indicate the primary outcome measures used for each investigation and the level of certainty. The even numbered tables (Table ES2 to

ES16) summarize the status of evidence for the 37 potential therapeutic options for P-ACC for which studies were identified through this systematic review.

P-ACC-related asthenia or fatigue

Table ES1. List of RCTs on interventions for P-ACC-related asthenia or fatigue with primary outcome measures and certainty (n=27)

Intervention	Overall number of studies including the intervention, n=27	HRQL improvement (n of studies)	Overall symptom improvement (n of studies)	Fatigue improvement (n of studies)	Functional capacity improvement (n of studies)	Strength improvement (n of studies)	Adverse events (n of studies)	Severe adverse events (n of studies)
Physical training	NEW	4	2	2	2	1		
Telerehabilitation	NEW	4	3	3	3			
Fermented food supplements		2	2					
tDCS		2	1	2			1	
1_MNA		1		1	1			
Actovegin		1		1				
ADAPT_232 (adaptogens)		1		1				
Arginine_Vitamin C		1		1		1		
Aromatherapy		1	1	1				
AXA1125		1		1	1		1	
Cognitive behavioral therapy	NEW	1		1	1			
CQ10		1	1	1				
Cytoflavin		1		1				
Enzymes_Probiotics		1		1				
Hydrogen (nasal)		1		1	1			
Immunodaa		1	1	1				
Mindfulness		1		1				
Phytochemicals		1	1	1				
Leronlimab		1		1				

	GRADE High- Moderate certainty	GRADE Low certainty
Beneficial effect		
No significant effect		
Harmful effect		
Uncertain effect		
No evidence or no estimable effect		

Table ES2. Summary of findings on potential therapeutic options for P-ACC-related asthenia or fatigue (n=19), as of 16 June 2023

	Intervention	Summary of findings
1	1-MNA	Uncertainty in potential benefits and harms. Further research is needed.
2	Actovegin	Actovegin may improve fatigue. However, certainty of the evidence was low. Further research is needed.
3	ADAPT-232 (adaptogens)	ADAPT-232 may not improve fatigue. However, certainty of the evidence was low. Further research is needed.
4	Arginine + Vitamin C	Uncertainty in potential benefits and harms. Further research is needed.
5	Aromatherapy	Uncertainty in potential benefits and harms. Further research is needed.
6	AXA1125	AXA1125 may increase fatigue improvement but may not increase functional capacity improvement. However, certainty of the evidence was low. Further research is needed.
7	Coenzyme Q10	Uncertainty in potential benefits and harms. Further research is needed.
8	Cognitive behavioral therapy	Cognitive behavioral therapy may increase fatigue improvement and functional capacity improvement. Further research is needed.
9	Cytoflavin	Cytoflavin may not improve fatigue. However, certainty of the evidence was low. Further research is needed.
10	Enzymes + probiotics	Enzymes + probiotics may improve fatigue. However, certainty of the evidence was low. Further research is needed.
11	Fermented food supplements	Uncertainty in potential benefits and harms. Further research is needed.
12	Hydrogen (nasal)	Uncertainty in potential benefits and harms. Further research is needed.
13	Immunodaat	Uncertainty in potential benefits and harms. Further research is needed.
14	Leronlimab	Uncertainty in potential benefits and harms. Further research is needed.

	Intervention	Summary of findings
15	Mindfulness	Uncertainty in potential benefits and harms. Further research is needed.
16	Phytochemicals	Phytochemicals may improve fatigue and HRQL. However, certainty of the evidence was low. Further research is needed.
17	Physical training	Physical training may improve HRQL, fatigue and functional capacity. However, certainty of the evidence was low. Further research is needed.
18	Transcranial direct current stimulation (tDCS)	tDCS may improve fatigue and HRQL and may not increase adverse events. However, certainty of the evidence was low. Further research is needed.
19	Telerehabilitation	Telerehabilitation may improve fatigue and functional capacity. However, certainty of the evidence was low. Further research is needed.

Key findings

• **Therapeutic options:** According to the WHO International Clinical Trials Registry Platform (ICTRP), multiple potential interventions are being assessed in hundreds of clinical trials and observational studies. In this review, we identified and examined 18 therapeutic options for P-ACC-related asthenia or fatigue.

• **Actovegin:** The results of one RCT suggest that actovegin may improve fatigue. However, certainty of the evidence was low because of imprecision. Further research is needed.

• **ADAPT-232 (adaptogens):** The results of one RCT suggest that ADAPT-232 may not improve fatigue. However, certainty of the evidence was low because of imprecision. Further research is needed.

• **AXA1125 (amino acids + N-acetylcysteine):** AXA1125 may increase fatigue improvement but may not increase functional capacity improvement. However, certainty of the evidence was low because of imprecision. Further research is needed.

- **Cytoflavin:** The results of one RCT suggest that cytoflavin may not improve fatigue. However, certainty of the evidence was low because of imprecision and risk of bias. Further research is needed.
- **Enzymes + probiotics:** The results of one RCT suggest that enzymes + probiotics may not improve fatigue. However, certainty of the evidence was low because of imprecision and risk of bias. Further research is needed.
- **Physical training:** The results of four RCTs suggest that physical training may improve HRQL, fatigue and functional capacity. However, certainty of the evidence was low because of imprecision and risk of bias. Further research is needed.
- **Transcranial direct current stimulation (tDCS):** The results of two RCTs suggest that tDCS may improve fatigue and HRQL and may not increase adverse events. However, certainty of the evidence was low because of imprecision. Further research is needed.
- **Telerehabilitation:** The results of four RCTs suggest that telerehabilitation may improve fatigue and functional capacity. However, certainty of the evidence was low because of imprecision and risk of bias. Further research is needed.

P-ACC-related dyspnea

Table ES3. List of RCTs of interventions for P-ACC-related dyspnea with primary outcome measures and certainty (n=15)

Intervention	Overall number of studies including the intervention, n=15	HRQL improvement (n of studies)	Dyspnea improvement (n of studies)	Functional capacity improvement (n of studies)	Pulmonary function improvement (n of studies)	Radiological response (n of studies)	Adverse events (n of studies)	Severe adverse events (n of studies)
Respiratory training	NEW	9	5	8	2	1		
ADAPT_232 (adaptogens)		1		1				
Antifibrotics		1			1			
Diaphragm release	NEW	1		1				
High dose steroids		1		1		1	1	1
Nebivolol		1		1				
Treamid		1		1	1		1	

	GRADE High- Moderate certainty	GRADE Low certainty
Beneficial effect		
No significant effect		
Harmful effect		
Uncertain effect		
No evidence or no estimable effect		

Table ES4. Summary of findings on potential therapeutic options for P-ACC-related dyspnea (n=7), as of 16 June 2023

	Intervention	Summary of findings
1	Antifibrotics	Uncertainty in potential benefits and harms. Further research is needed.
2	ADAPT-232 (adaptogens)	ADAPT-232 may not improve dyspnea. However, certainty of the evidence was low. Further research is needed.
3	Diaphragm release	Uncertainty in potential benefits and harms. Further research is needed.
4	High dose steroids	High dose steroids, compared to standard dose steroids, may not improve dyspnea and may not increase adverse events. However, certainty of the evidence was low. Further research is needed.
5	Nebivolol	Uncertainty in potential benefits and harms. Further research is needed.
6	Respiratory training/rehabilitation	Respiratory training/rehabilitation probably improves HRQL and dyspnea, and may improve functional capacity.

	Intervention	Summary of findings
7	Treamid	Treamid may improve dyspnea and pulmonary function but may not improve functional capacity. Treamid may increase adverse events. However, certainty of the evidence was low. Further research is needed.

Key findings

- **Therapeutic options:** According to the WHO International Clinical Trials Registry Platform (ICTRP), multiple potential interventions are being assessed in hundreds of clinical trials and observational studies. In this review, we identified and examined five therapeutic options for P-ACC-related dyspnea.
- **ADAPT-232 (adaptogens):** The results of one RCT suggest that ADAPT-232 may not improve dyspnea. However, certainty of the evidence was low because of imprecision. Further research is needed.
- **High dose steroids:** The results of one RCT suggest that high dose steroids (prednisone 40 mg a day) may not improve dyspnea compared to standard dose steroids (prednisone 10 mg a day). However, certainty of the evidence was low because of risk of bias and imprecision. Further research is needed.
- **Respiratory training/rehabilitation:** The results of nine RCTs suggest that respiratory training probably improves HRQL and dyspnea, and may improve functional capacity.
- **Treamid:** The results of one RCT suggest that treamid may improve dyspnea and pulmonary function but may not improve functional capacity. However, certainty of the evidence was low because of imprecision. Further research is needed.

P-ACC-related neurocognitive symptoms or sleep disturbances

Table ES5. List of RCTs of interventions for P-ACC-related neurocognitive symptoms or sleep disturbances with primary outcome measures and certainty (n=8)

Intervention	Overall number of studies including the intervention, n=8	HRQL improvement (n of studies)	Overall symptom improvement (n of studies)	Cognitive improvement (n of studies)	Depression improvement (n of studies)	Sleep quality improvement (n of studies)	Adverse events (n of studies)	Severe adverse events (n of studies)
Actovegin	1			1				
Electrical stimulation	1							
HBO	1	1		1	1			
Mindfulness	1				1			
Palmitoylethanolamide_Luteolin	1			1				
Physical training	1	1				1		
taVNS	1							
tDCS	1			1				

	GRADE High- Moderate certainty	GRADE Low certainty
Beneficial effect		
No significant effect		
Harmful effect		
Uncertain effect		
No evidence or no estimable effect		

Table ES6. Summary of findings on potential therapeutic options for P-ACC-related neurocognitive symptoms or sleep disturbances (n=8), as of 16 June 2023

	Intervention	Summary of findings
1	Actovegin	Actovegin may improve cognition. However, certainty of the evidence was low. Further research is needed.
2	Electric stimulation	Uncertainty in potential benefits and harms. Further research is needed.
3	Hyperbaric oxygen (HBO)	HBO may improve HRQL. However, certainty of the evidence was low. Further research is needed.
4	Mindfulness	Uncertainty in potential benefits and harms. Further research is needed.
5	Palmitoylethanolamide + Luteolin	Uncertainty in potential benefits and harms. Further research is needed.
6	Physical training	Uncertainty in potential benefits and harms. Further research is needed.

	Intervention	Summary of findings
7	Transcutaneous auricular vagus nerve stimulation (taVNS)	Uncertainty in potential benefits and harms. Further research is needed.
8	Transcranial direct current stimulation (tDCS)	tCDS may not improve cognition. However, certainty of the evidence was low. Further research is needed.

Key findings

- **Therapeutic options:** According to the WHO International Clinical Trials Registry Platform (ICTRP), multiple potential interventions are being assessed in hundreds of clinical trials and observational studies. In this review, we identified and examined three therapeutic options for PCC neurocognitive symptoms or sleep disturbances.
- **Actovegin:** The results of one RCT suggest that actovegin may improve cognition. However, certainty of the evidence was low because of risk of bias. Further research is needed.
- **Hyperbaric oxygen (HBO):** The results of one RCT suggest that HBO may improve HRQL. However, certainty of the evidence was low because of imprecision. Further research is needed.
- **Transcranial direct current stimulation (tDCS):** The results of one RCT suggest that tDCS may not improve cognition. However, certainty of the evidence was low because of imprecision. Further research is needed.

P-ACC-related olfactory and/or gustatory dysfunction

Table ES7. List of RCTs of interventions for P-ACC-related olfactory and/or gustatory dysfunction with primary outcome measures and certainty (n=11)

Intervention	Overall number of studies including the intervention, n=11	HRQL improvement (n of studies)	Overall symptom improvement (n of studies)	Olfactory symptoms improvement (n of studies)	Gustatory symptoms improvement (n of studies)	Adverse events (n of studies)	Severe adverse events (n of studies)
Olfactory training	3			3			
Steroids	2			2	1		
ADAPT_232 (adaptogens)	1			1			
Diode Laser NEW	1				1		
DTPA NEW	1			1			
Palmitoylethanolamide_Luteolin NEW	1			1			
Steroids (nasal)	1			1			
Theophylline_nasal	1	1		1		1	

	GRADE High- Moderate certainty	GRADE Low certainty
Beneficial effect		
No significant effect		
Harmful effect		
Uncertain effect		
No evidence or no estimable effect		

Table ES8. Summary of findings on potential therapeutic options for P-ACC-related olfactory and/or gustatory dysfunction (n=8), as of 16 June 2023

	Intervention	Summary of findings
1	ADAPT-232 (adaptogens)	ADAPT-232 may not improve olfactory symptoms. However, certainty of the evidence was low. Further research is needed.
2	Diode laser	Uncertainty in potential benefits and harms. Further research is needed.
3	Olfactory training	Uncertainty in potential benefits and harms. Further research is needed.
4	Palmitoylethanolamide + Luteolin	Palmitoylethanolamide + Luteolin may improve olfactory symptoms. However, certainty of the evidence was low. Further research is needed.
5	Pentasodium diethylenetriamine pentaacetate (DTPA)	Uncertainty in potential benefits and harms. Further research is needed.

	Intervention	Summary of findings
6	Steroids (nasal)	Uncertainty in potential benefits and harms. Further research is needed.
7	Steroids	Steroids may not improve olfactory nor gustatory symptoms. Further research is needed.
8	Theophylline (nasal)	Uncertainty in potential benefits and harms. Further research is needed.

Key findings

- **Therapeutic options:** According to the WHO International Clinical Trials Registry Platform (ICTRP), multiple potential interventions are being assessed in hundreds of clinical trials and observational studies. In this review, we identified and examined five therapeutic options for PCC olfactory and/or gustatory dysfunction.
- **ADAPT-232 (adaptogens):** The results of one RCT suggest that ADAPT-232 may improve olfactory symptoms. However, certainty of the evidence was low because of imprecision. Further research is needed.
- **Palmitoylethanolamide + Luteolin:** The results of one RCT suggest that Palmitoylethanolamide + Luteolin may improve olfactory symptoms. However, certainty of the evidence was low because of imprecision. Further research is needed.
- **Steroids:** The results of two RCTs suggest that steroids may not improve olfactory nor gustatory symptoms. However, certainty of the evidence was low because of imprecision. Further research is needed.

P-ACC-related cardiovascular system symptoms

Table ES9. List of RCTs of interventions for P-ACC-related cardiovascular system symptoms with primary outcome measures and certainty (n=1)

Intervention	Overall number of studies including the intervention, n=1	HRQL improvement (n of studies)	Overall symptom improvement (n of studies)	Tachycardia improvement (n of studies)	Adverse events (n of studies)	Severe adverse events (n of studies)
Ivabradine	1			1		

	GRADE High- Moderate certainty	GRADE Low certainty
Beneficial effect		
No significant effect		
Harmful effect		
Uncertain effect		
No evidence or no estimable effect		

Table ES10. Summary of findings on potential therapeutic options for P-ACC-related cardiovascular system symptoms (n=1), as of 16 June 2023

	Intervention	Summary of findings
1	Ivabradine	Uncertainty in potential benefits and harms. Further research is needed.

Key findings

- **Therapeutic options:** According to the WHO International Clinical Trials Registry Platform (ICTRP), multiple potential interventions are being assessed in hundreds of clinical trials and observational studies. In this review, we identified and examined one therapeutic option for P-ACC- related cardiovascular system symptoms.
- **The effects of assessed interventions are uncertain.**

P-ACC-related psychological distress

Table ES11. List of RCTs of interventions for P-ACC-related psychological distress with primary outcome measures and certainty (n=1)

Intervention	Overall number of studies including the intervention, n=1	HRQL improvement (n of studies)	Depression improvement (n of studies)	Post traumatic stress disorder improvement (n of studies)	Psychologic distress improvement (n of studies)	Adverse events (n of studies)	Severe adverse events (n of studies)
Virtual reality info	1		1	1	1		

	GRADE High- Moderate certainty	GRADE Low certainty
Beneficial effect		
No significant effect		
Harmful effect		
Uncertain effect		
No evidence or no estimable effect		

Table ES12. Summary of findings on potential therapeutic options for PCC psychological distress (n=1), as of 16 June 2023

	Intervention	Summary of findings
1	Virtual reality informational video	Virtual reality informational video may improve depression, post-traumatic stress, and psychological distress. However, certainty of the evidence was low. Further research is needed.

Key findings

- **Therapeutic options:** According to the WHO International Clinical Trials Registry Platform (ICTRP), multiple potential interventions are being assessed in hundreds of clinical trials and observational studies. In this review, we identified and examined one therapeutic option for PCC psychological distress.
- **Virtual reality informational video:** The results of one RCT suggest that Virtual reality informational video may improve depression, post-traumatic stress, and psychological distress. However, certainty of the evidence was low because of imprecision. Further research is needed.

P-ACC-related thromboembolic risk

Table ES13. List of RCTs of interventions for P-ACC-related thromboembolic risk with primary outcome measures and certainty (n=2)

Intervention	Overall number of studies including the intervention, n=2	Mortality (n of studies)	HRQL improvement (n of studies)	VTE (symptomatic) (n of studies)	Major bleeding (n of studies)	Severe adverse events (n of studies)
Anticoagulants_Proph	2	2	1	2	2	

	GRADE High- Moderate certainty	GRADE Low certainty
Beneficial effect		
No significant effect		
Harmful effect		
Uncertain effect		
No evidence or no estimable effect		

Table ES14. Summary of findings on potential therapeutic options for PCC thromboembolic risk (n=1), as of 16 June 2023

	Intervention	Summary of findings
1	Anticoagulants (prophylactic dose)	Anticoagulants may not have an important effect on mortality, VTE, major bleeding and HRQL. However, certainty of the evidence was low because of imprecision. Further research is needed.

Key findings

- **Therapeutic options:** According to the WHO International Clinical Trials Registry Platform (ICTRP), multiple potential interventions are being assessed in hundreds of clinical trials and observational studies. In this review, we identified and examined one therapeutic option for PCC olfactory and/or gustatory dysfunction.

- **Anticoagulants:** The results of two RCTs suggest that anticoagulants (rivaroxaban and apixaban) may not have an important effect on mortality, HRQL, VTE or major bleeding. However, certainty of the evidence was low because of risk of imprecision. Further research is needed.

Pediatric inflammatory multisystem syndrome associated with SARS-CoV-2 (PIMS-TS)

Table ES13. List of RCTs of interventions for PIMS-TS with primary outcome measures and certainty (n=1)

Intervention	Overall number of studies including the intervention, n=1	Mortality (n of studies)	Time to discharge reduction (n of studies)	Respiratory support (n of studies)	Inotropics requirement (n of studies)	LVEF < 55% (n of studies)	Arrhythmia (n of studies)	VTE (symptomatic) (n of studies)	Major bleeding (n of studies)
Steroids	1		1	1	1	1	1	1	1

	GRADE High- Moderate certainty	GRADE Low certainty
Beneficial effect		
No significant effect		
Harmful effect		
Uncertain effect		
No evidence or no estimable effect		

Table ES14. Summary of findings on potential therapeutic options for PCC thromboembolic risk (n=1), as of 16 June 2023

	Intervention	Summary of findings
1	Steroids	Steroids may reduce time to discharge and respiratory support requirements. However, certainty of the evidence was low for risk of bias and imprecision. Further research is needed.

Key findings

- **Therapeutic options:** According to the WHO International Clinical Trials Registry Platform (ICTRP), multiple potential interventions are being assessed in hundreds of clinical trials and observational studies. In this review, we identified and examined one therapeutic option for PCC olfactory and/or gustatory dysfunction.
- **Steroids:** The results of one RCT suggest that steroids may reduce time to discharge and respiratory support requirements. However, certainty of the evidence was low because of risk of bias and imprecision. Further research is needed.

P-ACC prophylaxis

Table ES15. List of RCTs of interventions for P-ACC prophylaxis with primary outcome measures and certainty (n=5)

Intervention	Overall number of studies including the intervention, n=5	Mortality (n of studies)	HRQL improvement (n of studies)	P-ACC (n of studies)	Adverse events (n of studies)	Severe adverse events (n of studies)
Fluvoxamine	2			2		
CP	1			1		
Ivermectin	1			1		
Leflunomide	1			1		1
Metformin	1			1		
Remdesivir	1	1		1		

	GRADE High- Moderate certainty	GRADE Low certainty
Beneficial effect		
No significant effect		
Harmful effect		
Uncertain effect		
No evidence or no estimable effect		

Table ES16. Summary of findings on potential therapeutic options for P-ACC prophylaxis (n=6), as of 9 May 2023

	Intervention	Summary of findings
1	Convalescent plasma	Convalescent plasma may not reduce P-ACC. However, certainty of the evidence was low. Further research is needed.
2	Fluvoxamine	Fluvoxamine may not reduce P-ACC. However, certainty of the evidence was low. Further research is needed.
3	Ivermectine	Ivermectin may not reduce P-ACC. However, certainty of the evidence was low. Further research is needed.
4	Leflunomide	Leflunomide may not reduce P-ACC. However, certainty of the evidence was low. Further research is needed.
5	Metformin	Metformin may reduce P-ACC. However, certainty of the evidence was low. Further research is needed.

	Intervention	Summary of findings
6	Remdesivir	Remdesivir may not reduce P-ACC. However, certainty of the evidence was low. Further research is needed.

Key findings

• **Therapeutic options:** According to the WHO International Clinical Trials Registry Platform (ICTRP), multiple potential interventions are being assessed in hundreds of clinical trials and observational studies. In this review, we identified and examined one therapeutic option for PCC olfactory and/or gustatory dysfunction.

• **Metformin:** The results of one RCT suggest that metformin may reduce P-ACC. However, certainty of the evidence was low because of risk of bias and imprecision. Further research is needed.

• **Ivermectin:** The results of one RCT suggest that ivermectin may reduce P-ACC. However, certainty of the evidence was low because of risk of bias and imprecision. Further research is needed.

• **Convalescent plasma:** The results of one RCT suggest that convalescent plasma may not reduce P-ACC. However, certainty of the evidence was low because of risk of bias and imprecision. Further research is needed.

• **Remdesivir:** The results of one RCT suggest that remdesivir may not reduce P-ACC. However, certainty of the evidence was low because of risk of bias and imprecision. Further research is needed.

• **Leflunomide:** The results of one RCT suggest that leflunomide may not reduce P-ACC. However, certainty of the evidence was low because of risk of bias and imprecision. Further research is needed.

- **Fluvoxamine:** The results of two RCTs suggest that fluvoxamine may not reduce P-ACC. However, certainty of the evidence was low because of imprecision. Further research is needed.

Changes since previous edition

- **Telerehabilitation for P-ACC related asthenia or fatigue:** New evidence included affecting results interpretation and/or certainty of the evidence judgments.

- **Cognitive behavioral therapy for P-ACC related asthenia or fatigue:** New evidence included affecting results interpretation and/or certainty of the evidence judgments.

- **Diode laser for P-ACC related olfactory and/or gustatory dysfunction:** New evidence included affecting results interpretation and/or certainty of the evidence judgments.

- **Respiratory training/rehabilitation for P-ACC related dyspnea:** New evidence included affecting results interpretation and/or certainty of the evidence judgments.

- **Physical training for P-ACC related asthenia or fatigue:** New evidence included affecting results interpretation and/or certainty of the evidence judgments.

- **Pentasodium diethylenetriamine pentaacetate (DTPA) for P-ACC related olfactory and/or gustatory dysfunction:** New evidence included affecting results interpretation and/or certainty of the evidence judgments.

- **Diaphragm release for P-ACC related dyspnea:** New evidence included affecting results interpretation and/or certainty of the evidence judgments.

- **Palmitoylethanolamide + Luteolin:** New evidence included affecting results interpretation and/or certainty of the evidence judgments.

Concluding remarks

- The Pan American Health Organization (PAHO) is continually monitoring ongoing research on any possible therapeutic options. As evidence emerges, PAHO will immediately assess and update its position, particularly as it applies to any special population subgroups such as children, expectant mothers, and those with immune conditions.
- PAHO is also mindful of the emerging differential impact of PCC on ethnic and minority groups and is continuously seeking data that could help in mitigating excess risk of severe illness or death in minority subgroups. These groups are plagued by social and structural inequities that bring to bear a disproportionate burden of COVID-19 illness.
- The safety of the patient suffering from COVID-19 is a key priority to improve the quality of care in the provision of health services.
- Adequately designed and reported clinical trials are crucial for the practice of evidence-based medicine. Most of the research to date on PCC has very poor methodology that is hidden and very difficult to validate. Greater transparency and better designed studies are urgently needed.

Systematic review of therapeutic options for post acute or post COVID-19 condition (P-ACC)

Background

Post COVID-19 condition (PCC), also known as long COVID or post-acute sequelae of SARS-CoV-2 infection (PASC), is the continuation or development of new symptoms in the period after acute infection with SARS-CoV-2 (1–4). The World Health Organization (WHO) definition of PCC states that these symptoms should be present after three months of the initial SARS-CoV-2 infection and last for at least two months with no other explanation (1, 2). While PASC definitions state persistent or new symptoms need to be present 30 days after a documented SARS-COV-2 infection or the onset of COVID-19 symptoms (3, 4). Post COVID-19 condition or post-acute sequelae of SARS-CoV-2 infection (P-ACC) can affect anyone exposed to SARS-CoV-2, regardless of age or severity of acute infection. Many of the reported symptoms are debilitating and have a strong negative impact on mental health and quality of life (5). While most patients recover, some may experience multiple outcomes, with multiple organ systems affected simultaneously, including cardiovascular, mental, metabolic, renal, and others (3, 6). Recommendations for the management of patients with PCC are continuously being developed and need to evolve as evidence of interventions effects becomes available (7).

In this review, we compiled the following evidence on potential therapeutic options for P-ACC. We included all the identified clinical forms, symptoms, and manifestations of P-ACC for which an intervention was assessed in at least one randomized controlled trial (RCT). We hope this information will support investigators, policymakers, and prescribers navigate the flood of relevant data to ensure that management of P-ACC, at both the individual and population levels, is based on the best available knowledge. We will endeavor to continually update this resource as more research is released into the public space.

Methods

We used the Living Overview of Evidence (L-OVE; available from: <https://iloveevidence.com>) platform to identify studies for inclusion in this review. This platform is a system that maps PICO (Patient–Intervention–Comparison–Outcome) questions to a repository developed by Epistemonikos Foundation. This repository is continuously updated through searches in electronic databases, preprint servers, trial registries, and other resources relevant to COVID-19. The latest version of the methods, the total number of sources screened, and a living flow diagram and report of the project is updated regularly on the L-OVE website (8).

Search strategy

We systematically searched in L-OVE for COVID-19. The search terms and databases covered are described on the L-OVE search strategy methods page (available from: https://app.iloveevidence.com/loves/5e6fdb9669c00e4ac072701d?question_domain=undefined§ion=methods). The repository is continuously updated, and the information is transmitted in real time to the L-OVE platform. It was last checked for this review on 16 June 2023. The searches covered the period from the inception date of each database, and no study design, publication status, or language restriction was applied.

Study selection

The results of the searches in the individual sources were de-duplicated by an algorithm that compares unique identifiers (database identification number, digital object identifier [DOI], trial registry identification number), and citation details (i.e., author names, journal, year of publication, volume, number, pages, article title, and article abstract). Then, the information matching the search strategy was sent in real time to the L-OVE platform, where at least two authors independently screened the titles and abstracts yielded against the inclusion criteria. We obtained the full reports for all titles that appeared to meet the inclusion criteria or required further analysis and then decided about their inclusion.

Inclusion criteria

We aimed to find all available RCTs for potential therapeutic interventions for P-ACC with study designs that included head-to-head comparisons, or control groups with no intervention or a placebo. Target patient populations included both adults and children with persistent, or new, symptoms or clinical manifestations after acute COVID-19. We used the term Post Acute or Post COVID-19 condition (P-ACC) to refer to the population included in our review (studies reporting on patients with persistent or new symptoms after acute COVID-19 independently of the time of onset of those symptoms) (1–4). We focused on comparative effectiveness studies that provide evidence on outcomes of crucial importance to patients (mortality, health-related quality of life [HRQL], and disease-specific symptoms).

Living evidence synthesis

An artificial intelligence algorithm deployed in the Coronavirus/COVID-19 topic of the L-OVE platform provides instant notification of articles with a high likelihood of being eligible. The authors review them, decide upon inclusion, and update the living web version of the review accordingly. If meta-analytical pooling is possible from retrieved evidence, we will do this to derive more precise estimates of effect and derive additional statistical power. No electronic database search restrictions were imposed.

For any meta-analytical pooling, if and when data allow, we pool all studies and present the combined analysis with relative and absolute effect sizes. To assess interventions' absolute effects, we applied relative effects to baseline risks (risks with no intervention). For baseline risks we used the mean risk in the control groups from included RCTs. For continuous outcomes, when possible, we calculated relative and absolute effects by estimating the proportion of patients with important improvement or deterioration following published guidance (9).

For result interpretations and imprecision assessment we used a minimally contextualized approach that considers whether the 95% confidence interval (CI) includes the null effect, or, when the point estimate is close to the null effect, whether the 95% CI lies within the

boundaries of small but important benefit and harm that corresponds to every outcome assessed (10, 11).

We used the following absolute effects thresholds to define important benefits and harms: Mortality, +/-1%; HRQL improvement, +/-2%; Overall symptom improvement, +/-5%; Functional capacity improvement, +/-5%; Strength improvement, +/-5%; Fatigue improvement, +/-5%; Pulmonary function improvement, +/-10%; Radiological response, +/-10%; Cognitive improvement, +/-5%; Depression improvement, +/-5%; Olfactory symptoms improvement, +/-5%; Gustatory symptoms improvement, +/-5%; Tachycardia improvement, +/-5%; Venous thromboembolism (VTE) (symptomatic), +/-3%; Post-traumatic stress disorder improvement, +/-5%; Psychological distress improvement, +/-5%; Major bleeding, +/-3%; Severe adverse events, +/-3%; Adverse events, +/-5%; Time to discharge reduction, +/-4%; Respiratory support requirement +/-2%; Inotropic requirement +/-2%; Left ventricular ejection fraction deterioration (LVEF <55%) +/-5%; Arrhythmia +/-5%; P-ACC, +/-3%.

For some interventions when we found significant heterogeneity, we performed subgroup analysis considering: 1) risk of bias (high/moderate vs low risk of bias); and 2) intervention characteristics (e.g., different doses or administration schemes). When we observed significant differences between subgroups, we presented individual subgroups' estimates of effect and certainty of the evidence assessment.

A risk of bias assessment was applied to RCTs focusing on randomization, allocation concealment, blinding, attrition, or other biases relevant to the estimates of effect (Table 1) (12). The GRADE approach was used to assess the certainty of the body of evidence for every comparison on an outcome basis (13).

Study selection, data extraction, and risk of bias assessment were performed, independently and in parallel, by two reviewers. Discrepancies were resolved by discussion.


We used MAGIC authoring and publication platform (available from: <https://app.magicapp.org/>) to generate the tables summarizing our findings, which are included in Annex 1.

Results

Studies identified and included

The study identification and selection process is shown in Figure 1. A total of 65 RCTs were selected for inclusion. A list of excluded studies is available upon request.

Figure 1. Study identification and selection process



Risk of bias

Overall, our risk of bias assessment for the limited reported RCTs found high risk of bias due to suboptimal randomization, allocation concealment, and blinding (as well as other methodological and reporting concerns). Most RCTs were also very small in size and had small event numbers. The methods were very poor overall, and the reporting was suboptimal. In general, follow-up was short. The risk of bias assessment of each RCT is presented in Table 1.

Table 1. Risk of bias of included RCTs

Study	Risk-of-bias arising from randomization process	Risk-of-bias due to deviations from the intended interventions	Risk-of-bias due to missing outcome data	Risk-of-bias in measurement of the outcome	Risk-of-bias in selection of the reported result	Overall Risk-of-bias judgement	
						Mortality	HRQL symptom specific outcomes
Vaira LA et al	High	Some Concerns	Low	Some Concerns	Low	High	High
RC 4-7-2020 (Abdelalim AA et al)	High	Some Concerns	Low	Some Concerns	Low	High	High
Di Stadio	Low	Low	Low	Low	Low	Low	Low
Chudzik M et al	High	Some Concerns	Low	Some Concerns	Low	High	High
CITADEL	High	Some Concerns	Low	Some Concerns	Low	High	High
MICHELLE	Low	Some Concerns	Low	Some Concerns	Low	Low	High
Zilberman-Itskovich	Low	Low	Low	Low	Low	Low	Low
Botek M et al	High	Some Concerns	Low	Some Concerns	Low	High	High
Jadhav KP et al	High	Some Concerns	Low	Some Concerns	Low	High	High
COLDSTER	Low	Some Concerns	Low	Some Concerns	Low	Low	High
Oliver-Mas	Low	Low	Low	Low	Low	Low	Low
Nambi	Low	Some Concerns	Low	Some Concerns	Low	Low	High
Di Stadio_2	Low	Some Concerns	Low	Some Concerns	Low	Low	High
Hansen	Low	Low	Low	Low	Low	Low	Low
Tosato	High	Some Concerns	Low	Some Concerns	Low	High	High
Rathi	High	Some Concerns	Low	Some Concerns	Low	High	High
Bazdyrev	Low	Low	Low	Low	Low	Low	Low
King	Low	Some Concerns	Low	Some Concerns	Low	Low	High
ICU-VR	High	Some Concerns	Low	Some Concerns	Low	High	High
ENO Breathe	Low	Some Concerns	Low	Some Concerns	Low	Low	High
Pires	High	Some Concerns	Low	Some Concerns	Low	High	High
McNarry	High	Some Concerns	Low	Some Concerns	Low	High	High
Srinivasan	High	Some Concerns	Low	Some Concerns	Low	High	High
Kharaeva_Moderate	High	Some Concerns	Low	Some Concerns	Low	High	High
Kharaeva_Severe	High	Some Concerns	Low	Some Concerns	Low	High	High
Gaylis	High	Some Concerns	Low	Some Concerns	Low	High	High
Karosanidze	Low	Low	Low	Low	Low	Low	Low
Badran	Low	Low	Low	Low	Low	Low	Low
COVANOS	Low	Some Concerns	Low	Some Concerns	Low	Low	High
RECOVER	High	Some Concerns	Low	Some Concerns	Low	High	High
Kutashov	High	Some Concerns	Low	Some Concerns	Low	High	High
Vallier	High	Some Concerns	Low	Some Concerns	Low	High	High
Swissped RECOVERY	Low	Some Concerns	Low	Some Concerns	Low	Low	High
UK Phyto-V	High	Some Concerns	Low	Some Concerns	Low	High	High
Rodriguez-Blanco	High	Some Concerns	Low	Some Concerns	Low	High	High
COVID-OUT - Metformin	Low	Low	High	Low	Low	High	High
COVID-OUT - Ivermectin	Low	Low	High	Low	Low	High	High
COVID-OUT - Fluvoxamine	Low	Low	High	Low	Low	High	High
Santana	Low	Low	Low	Low	Low	Low	Low
CSSC-004	Low	Low	Some Concerns	Low	Low	Low	Some Concerns
Deshpande	High	Some Concerns	Low	Some Concerns	Low	High	High
Dal Negro	High	Some Concerns	Low	Some Concerns	Low	High	High
InsCOVID	Low	Some Concerns	Low	Some Concerns	Low	Low	High
Rutkowski	Low	Some Concerns	Low	Some Concerns	Low	Low	High

SOLIDARITY - Finland	Low	Low	Low	Low	Low	Low	Low
Hawkins	Low	Low	Low	Low	Low	Low	Low
Schepens	Low	Low	Low	Low	Low	Low	Low
Kusumawardani	High	Some Concerns	Low	Some Concerns	Low	High	High
SCENT2	Low	Low	Low	Low	Low	Low	Low
Hauswirth	High	Some Concerns	Low	Some Concerns	Low	High	High
Versace	High	Low	Low	Low	Low	High	High
ACTIV-4C	Low	Low	Low	Low	Low	Low	Low
Simpson	High	Some Concerns	Low	Some Concerns	Low	High	High
Stavrou	High	Some Concerns	Low	Some Concerns	Low	High	High
Zulbaran-Rojas	High	Some Concerns	Low	Some Concerns	Low	High	High
Kalayeh	High	Some Concerns	Low	Some Concerns	Low	High	High
DEFEAT-COVID	Low	Some Concerns	Low	Some Concerns	Low	Low	High
Kerget	High	Some Concerns	Low	Some Concerns	Low	High	High
Farahani	Low	Low	Low	Low	Low	Low	Low
Finnigan	Low	Low	Low	Low	Low	Low	Low
Longobardi	Low	Some Concerns	Low	Some Concerns	Low	Low	High
Kuut	Low	Some Concerns	Low	Some Concerns	Low	Low	High
Shabaan	Low	Low	Low	Low	Low	Low	Low
RECOVE	High	Some Concerns	Low	Some Concerns	Low	High	High
Del Corral	Low	Low	Low	Low	Low	Low	Low
Imam	High	Some Concerns	Low	Some Concerns	Low	High	High
Nagy	High	Some Concerns	Low	Some Concerns	Low	High	High
Ibrahim	High	Some Concerns	Low	Some Concerns	Low	High	High
Samper-Pardo	Low	Some Concerns	Low	Some Concerns	Low	Low	High

Main findings

P-ACC-related asthenia or fatigue

Actovegin

See Summary of findings Table A1, Annex 1

We identified one RCT including 444 participants in which Actovegin was compared against standard of care. Our results showed:

- Actovegin may improve fatigue, relative risk (RR) 1.84 (95% CI 1.59 to 2.14); risk difference (RD) 39.7% (95% CI 27.7% to 56.3%); Low certainty ⊕⊕○○

ADAPT-232 (adaptogens)

See Summary of findings Table A2, Annex 1

We identified one RCT including 99 participants in which ADAPT-232 was compared against standard of care. Our results showed:

- ADAPT-232 may not improve fatigue, relative risk (RR) 1.02 (95% CI 0.84 to 1.24); risk difference (RD) 1.6% (95% CI –12.6% to 18.9%); Low certainty ⊕⊕○○

Cytoflavin

See Summary of findings Table A3, Annex 1

We identified one RCT including 200 patients in which cytoflavin was compared against standard of care. Our results showed:

- Cytoflavin may not improve fatigue, RR 1.02 (95% CI 0.98 to 1.06); RD 2.1% (95% CI –1.9% to 6.2%); Low certainty ⊕⊕○○

Enzymes + probiotics

See Summary of findings Table A4, Annex 1

We identified one RCT including 200 patients in which enzymes + probiotics were compared against standard of care. Our results showed:

- Enzymes + probiotics may improve fatigue, RR 6.07 (95% CI 3.79 to 9.71); RD 76% (95% CI 41.8% to 85%); Low certainty ⊕⊕○○

Physical training

See Summary of findings Table A30, Annex 1

We identified four RCTs including 217 patients in which physical training was compared against standard of care. Our results showed:

- Physical training may improve HRQL, RR 1.41 (95% CI 0.81 to 2.58); RD 10% (95% CI -4.3% to 35.8%); Low certainty ⊕⊕○○
- Physical training may improve fatigue, RR 4.66 (95% CI 1.96 to 11.09); RD 44.4% (95% CI 11.7% to 87.9%); Low certainty ⊕⊕○○
- Physical training may improve functional capacity, RR 4.66 (95% CI 1.96 to 11.09); RD 44.4% (95% CI 11.7% to 87.9%); Low certainty ⊕⊕○○

Phytochemicals

See Summary of findings Table A5, Annex 1

We identified one RCT including 147 patients in which phytochemicals were compared against standard of care. Our results showed:

- Phytochemicals may improve HRQL, RR 1.33 (95% CI 1.03 to 1.71); RD 18% (95% CI 1.8% to 39%); Low certainty ⊕⊕○○
- Phytochemicals may improve fatigue, RR 1.24 (95% CI 0.95 to 1.62); RD 13.1% (95% CI -2.5% to 33.5%); Low certainty ⊕⊕○○


Transcranial direct current stimulation (tDCS)

See Summary of findings Table A6, Annex 1

We identified two RCTs including 117 patients in which tDCS was compared against standard of care. Our results showed:

- tDCS may improve fatigue, RR 1.36 (95% CI 0.76 to 2.45); RD -16.9% (95% CI -11.2% to 53%); Low certainty ⊕⊕○○ (see figure 2.)
- tDCS may improve HRQL, RR 1.37 (95% CI 1.09 to 1.71); RD -26% (95% CI -6.7% to 30%); Low certainty ⊕⊕○○
- tDCS may not increase adverse events, RR 0.83 (95% CI 0.26 to 2.73); RD -3.4% (95% CI -15.5% to 36%); Low certainty ⊕⊕○○
-

Figure 2. Fatigue in RCTs comparing tDCS with standard of care for treatment of patients with P-ACC-related asthenia/fatigue



AXA1125

See Summary of findings Table A27, Annex 1

We identified one RCT including 41 participants in which AXA1125 was compared against standard of care. Our results showed:

- AXA1125 may improve fatigue, RR 1.07 (95% CI 0.79 to 1.44); RD 5.1% (95% CI –16.6% to 34.5%); Low certainty ⊕⊕○○
- AXA1125 may not improve functional capacity, RR 0.87 (95% CI 0.51 to 1.48); RD –8.1% (95% CI –30% to 29.3%); Low certainty ⊕⊕○○

Telerehabilitation


See Summary of findings Table A28, Annex 1

We identified four RCTs including 148 participants in which telerehabilitation was compared against standard of care. Our results showed:

- Telerehabilitation may improve fatigue, RR 1.63 (95% CI 1.04 to 2.54); RD 19.1% (95% CI 1.2% to 46.8%); Low certainty ⊕⊕○○ (see figure 3.)

- Telerehabilitation may improve functional capacity, RR 1.48 (95% CI 1.07 to 2.04); RD 13.8% (95% CI 2% to 29.8%); Low certainty ⊕⊕○○

Figure 3. Fatigue in RCTs comparing telerehabilitation with standard of care for treatment of patients with P-ACC-related asthenia/fatigue



Cognitive behavioral therapy

See Summary of findings Table A29, Annex 1

We identified one RCT including 114 participants in which cognitive behavioral therapy was compared against standard of care. Our results showed:

- Telerehabilitation may improve fatigue, RR 2.2 (95% CI 1.35 to 3.58); RD 31.6% (95% CI 9.2% to 68%); Low certainty ⊕⊕○○
- Telerehabilitation may improve functional capacity, RR 1.37 (95% CI 1.08 to 1.73); RD 22.4% (95% CI 4.8% to 44.7%); Low certainty ⊕⊕○○

P-ACC-related dyspnea

ADAPT-232 (adaptogens)

[See summary of findings Table A7 in Annex 1](#)

We identified one RCT including 99 patients in which ADAPT-232 was compared against standard of care. Our results showed:

- ADAPT-232 may not improve dyspnea, RR 1 (95% CI 0.94 to 1.06); RD 0% (95% CI –5.4% to 5.7%); Low certainty ⊕⊕○○

High dose steroids

[See Summary of findings Table A9, Annex 1](#)

We identified one RCT including 130 patients in which high dose steroids (prednisone 40 mg a day) was compared against standard dose steroids (prednisone 10 mg a day). Our results showed:

- High dose steroids may not improve dyspnea, RR 1 (95% CI 0.87 to 1.15); RD 0% (95% CI –11% to 13%); Low certainty ⊕⊕○○
- High dose steroids may not increase adverse events, RR 0.92 (95% CI 0.75 to 1.13); RD –6.2% (95% CI –19.3% to 10%); Low certainty ⊕⊕○○


Respiratory training/rehabilitation

[See Summary of findings Table A10, Annex 1](#)

We identified nine RCTs including 480 patients in which different modalities of respiratory training/rehabilitation were compared with standard of care. In addition, we identified one study that compared home based respiratory training vs. inpatient respiratory training, two studies comparing VR respiratory training vs. conventional respiratory training and one study that compares incentive spirometry vs. conventional respiratory training. Our results showed:

- Respiratory training/rehabilitation probably improves HRQL, RR 1.47 (95% CI 1.14 to 1.89); RD 20.7% (95% CI 6.2% to 39.3%); Moderate certainty ⊕⊕⊕○ (see Figure 4)
- Respiratory training/rehabilitation probably improves dyspnea, RR 1.81 (95% CI 1.47 to 2.22); RD 23.3% (95% CI 13.5% to 35.1%); Moderate certainty ⊕⊕⊕○
- Respiratory training/rehabilitation may improve functional capacity, RR 1.65 (95% CI 1.11 to 2.46); RD 22.7% (95% CI 3.8% to 51%); Low certainty ⊕⊕○○

Figure 4. HRQL in RCTs comparing respiratory training/rehabilitation with standard of care for treatment of patients with P-ACC-related dyspnea.



Treamid

See Summary of findings Table A11, Annex 1

We identified one RCT including 59 patients in which treamid was compared with standard of care. Our results showed:

- Treamid may improve dyspnea, RR 1.96 (95% CI 0.9 to 4.25); RD 21.7% (95% CI –2.3% to 73.7%); Low certainty ⊕⊕○○
- Treamid may improve functional capacity, RR 1.1 (95% CI 0.64 to 1.9); RD 0.4% (95% CI 16.2% to 39.8%); Low certainty ⊕⊕○○
- Treamid may increase adverse events, RR 1.19 (95% CI 0.56 to 2.5); RD 5.5% (95% CI –12.7% to 43.6%); Low certainty ⊕⊕○○

P-ACC-related neurocognitive symptoms or sleep disturbances

Actovegin

See Summary of findings Table A12, Annex 1

We identified one RCT including 44 patients in which actovegin was compared with standard of care. Our results showed:

- Actovegin may improve cognition, RR 1.19 (95% CI 1.06 to 1.33); RD 12.7% (95% CI 4.2% to 22.3%); Low certainty ⊕⊕○○

Hyperbaric oxygen (HBO)

See Summary of findings Table A13, Annex 1

We identified one RCT including 73 patients in which HBO was compared with standard of care. Our results showed:

- HBO may improve HRQL, RR 1.3 (95% CI 0.84 to 2); RD 13.9% (95% CI –7.4% to 46.9%); Low certainty ⊕⊕○○

Transcranial direct current stimulation (tDCS)

See Summary of findings Table A14, Annex 1

We identified one RCT including 47 patients in which tDCS was compared with standard of care. Our results showed:

- tDCS may not improve HRQL, RR 0.59 (95% CI 0.33 to 1.05); RD –27.5% (95% CI –44.8% to 3.4%); Low certainty ⊕⊕○○

P-ACC-related olfactory and/or gustatory dysfunction

ADAPT-232 (adaptogens)

See Summary of findings Table A15, Annex 1

We identified one RCT including 99 patients in which ADAPT-232 was compared with standard of care. Our results showed:

- ADAPT-232 may not improve olfactory symptoms, RR 0.89 (95% CI 0.79 to 1.01); RD –10.3% (95% CI –20.5% to 1.4%); Low certainty ⊕⊕○○

Palmitoylethanolamide + Luteolin

See Summary of findings Table A16, Annex 1

We identified one RCT including 130 patients in which palmitoylethanolamide + luteolin was compared with standard of care. Our results showed:

- Palmitoylethanolamide + luteolin may improve olfactory symptoms, RR 3.11 (95% CI 1.47 to 6.66); RD 35.5% (95% CI 7.8% to 83.3%); Low certainty ⊕⊕○○


Steroids

See Summary of findings Table A17, Annex 1

We identified two RCTs including 131 patients in which steroids were compared with standard of care. Our results showed:

- Steroids may not improve olfactory symptoms, RR 1.09 (95% CI 0.83 to 1.44); RD 3.3% (95% CI –6.2% to 16.1%); Low certainty ⊕⊕○○ (figure 5)
- Steroids may not improve gustatory symptoms, RR 1.01 (95% CI 0.67 to 1.53); RD 0.5% (95% CI –14.6% to 23.3%); Low certainty ⊕⊕○○

Figure 5. Olfactory symptoms in RCTs comparing steroids with standard of care for treatment of patients with P-ACC-related olfactory and/or gustatory dysfunction.



P-ACC-related cardiovascular system symptoms

The effects of the assessed interventions are uncertain.

P-ACC-related psychological distress

Virtual reality (VR) informational video

See Summary of findings Table A18, Annex 1

We identified one RCT including 89 patients in which a virtual reality-based (VR) intervention was compared with standard of care. Our results showed:

- VR informational video may improve depression, RR 1.21 (95% CI 0.95 to 1.54); RD 14% (95% CI –3.7% to 36.7%); Low certainty ⊕⊕○○
- VR informational video may improve post-traumatic stress, RR 1.18 (95% CI 0.98 to 1.42); RD 13.8% (95% CI –1.5% to 32.3%); Low certainty ⊕⊕○○
- VR informational video may improve psychological distress, RR 1.49 (95% CI 1.08 to 2.05); RD 25.5% (95% CI 4.1% to 55.1%); Low certainty ⊕⊕○○

P-ACC-related thromboembolic risk


Anticoagulants

See Summary of findings Table A19, Annex 1

We identified two RCTs including 1535 patients with COVID-19 after hospitalization in which anticoagulants in prophylactic dose were compared with standard of care. Our results showed:

- Anticoagulants may not have an important effect on mortality, RR 0.85 (95% CI 0.38 to 0.88); RD -0.3% (95% CI -1.2% to 1.8%); Low certainty ⊕⊕○○ (Figure 6)
- Anticoagulants may not have an important effect on RR 0.99 (95% CI 0.78 to 1.24); Low certainty ⊕⊕○○
- Anticoagulants may not have an important effect on VTE, RR 1 (95% CI 0.29 to 3.45); RD 0% (95% CI -2.3% to 7.9%); Low certainty ⊕⊕○○ (based on low RoB studies)
- Anticoagulants may not have an important effect on VTE, RR 2.01 (95% CI 0.18 to 22.1); RD 0.1% (95% CI -0.1% to 1.2%); Low certainty ⊕⊕○○

Figure 6. Mortality in RCTs comparing anticoagulants with standard of care for treatment of patients with COVID-19 after hospitalization.



Pediatric inflammatory multisystem syndrome associated with SARS-CoV-2 (PIMS-TS)

Steroids

[See Summary of findings Table A20, Annex 1](#)

We identified one RCT including 75 patients in which systemic steroids were compared with intravenous immunoglobulins (IVIG). Our results showed:

- Steroids may reduce time to discharge, RR 1.09 (95% CI 0.88 to 1.39); RD 4.5% (95% CI -6% to 19.5%); Low certainty ⊕⊕○○
- Steroids may reduce respiratory support requirements, RR 0.49 (95% CI 0.27 to 0.89); RD -28.2% (95% CI -40.5% to -5.9%); Low certainty ⊕⊕○○

P-ACC prophylaxis

Metformin

[See Summary of findings Table A21, Annex 1](#)

We identified one RCT including 1125 patients in which metformin was compared with standard of care. Our results showed:

- Metformin may reduce P-ACC, RR 0.59 (95% CI 0.39 to 0.88); RD -4.3% (95% CI -6.4% to -1.2%); Low certainty ⊕⊕○○

Ivermectin

[See Summary of findings Table A22, Annex 1](#)

We identified one RCT including 739 patients in which metformin was compared with standard of care. Our results showed:

- Metformin may reduce P-ACC, RR 0.99 (95% CI 0.61 to 1.62); RD 0% (95% CI -1.7% to 2.6%); Low certainty ⊕⊕○○

Convalescent plasma

[See Summary of findings Table A23, Annex 1](#)

We identified one RCT including 882 patients in which metformin was compared with standard of care. Our results showed:

- Convalescent plasma may not reduce P-ACC, RR 0.93 (95% CI 0.77 to 1.12); RD -2.4% (95% CI -7.9% to -4.2%); Low certainty ⊕⊕○○

Remdesivir

[See Summary of findings Table A24, Annex 1](#)

We identified one RCT including 181 patients in which metformin was compared with standard of care. Our results showed:

- Remdesivir may not reduce P-ACC, RR 1.06 (95% CI 0.53 to 2.13); RD 0.8% (95% CI -6.9% to -16.4%); Low certainty ⊕⊕○○

Leflunomide

[See Summary of findings Table A25, Annex 1](#)

We identified one RCT including 172 patients in which leflunomide was compared with standard of care. Our results showed:

- Leflunomide may not reduce P-ACC, RR 1.28 (95% CI 0.92 to 1.77); RD 11.2% (95% CI -3.2% to 31.1%); Low certainty ⊕⊕○○

Fluvoxamine

[See Summary of findings Table A26, Annex 1](#)

We identified two RCTs including 680 patients in which fluvoxamine was compared with standard of care. Our results showed:

- Fluvoxamine may not reduce P-ACC, RR 0.99 (95% CI 0.81 to 1.21); RD -0.4% (95% CI -8.4% to 9.3%); Low certainty ⊕⊕○○

Full description of included studies

Tables 2 to 8 list all the identified studies that were included in this systematic review by intervention and P-ACC-related organ system affected. The treatments are arranged in alphabetical order. Study or author names, publication status, patient populations, interventions, sources of bias, outcomes, effect sizes, and certainty are listed for each study.

Table 2. Description of included studies and interventions effects for P-ACC-related asthenia or fatigue

1-MNA Uncertainty in potential benefits and harms. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Chudzik et al. (14) ; Peer reviewed; 2022	Patients with P-ACC (asthenia or fatigue after 30 days of acute COVID-19). 25 assigned to 1-MNA 58 mg a day and 25 assigned to standard of care.	Median age 49.5, male 32%, hypertension 14%, diabetes 2%	Not reported (NR)	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: No information Overall symptom improvement: No information Fatigue improvement: Very low certainty ⊕○○○ Functional capacity improvement: Very low certainty ⊕○○○ Strength improvement: No information Adverse events: No information Severe adverse events: No information

Actovegin

Actovegin may improve fatigue. However, certainty of the evidence was low. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Kutashov et al. (15) ; Peer reviewed; 2022	Patients with P-ACC (asthenia or fatigue after NR days of acute COVID-19). 222 assigned to Actovegin 1200 mg a day for 60 days and 222 assigned to standard of care.	Mean age 67.6, male 31.98%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: No information Overall symptom improvement: No information Fatigue improvement: RR 1.84 (95% CI 1.59 to 2.14); RD 39.7% (95% CI 27.7.6% to 53.6%); Low certainty ⊕⊕○○ Functional capacity improvement: No information Strength improvement: No information Adverse events: No information Severe adverse events: No information

ADAPT-232 (adaptogens)

ADAPT-232 may not improve fatigue. However, certainty of the evidence was low. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
------------------------------	---	---------------	-----------------------------	---------------------------------------	---

RCT

Karosanidze et al. (16) ; Peer reviewed; 2022	Patients with P-ACC (asthenia or fatigue after 30 days of acute COVID-19). 49 assigned to ADAPT-232 (adaptogens) 60 mL a day for 14 days and 50 assigned to standard of care.	Mean age 48.9, male 14%	NR	Low risk of bias	<p>HRQL improvement: No information</p> <p>Overall symptom improvement: No information</p> <p>Fatigue improvement: RR 1.02 (95% CI 0.84 to 1.24); RD 1.6% (95% CI -12.6% to 18.9%); Low certainty ⊕⊕○○</p> <p>Functional capacity improvement: No information</p> <p>Strength improvement: No information</p> <p>Adverse events: No information</p> <p>Severe adverse events: No information</p>
---	---	-------------------------	----	------------------	---

Arginine + Vitamin C

Uncertainty in potential benefits and harms. Further research is needed.

Study; publication status	Patients and interventions	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard
------------------------------	-------------------------------	---------------	-----------------------------	---------------------------------------	--------------------------------------

	analyzed				of care (SOC) and GRADE certainty of the evidence
RCT					
Tosato et al. (17) ; Peer reviewed; 2022	Patients with P-ACC (asthenia or fatigue after 28 days of acute COVID-19). 23 assigned to Arginine + Vitamin C 1.66 g/500 mg for 28 days and 23 assigned to standard of care.	Mean age 50.5 ± 14, male 34.8%, interval between COVID-19 and enrolment 254 days, hospitalization during COVID-19 56.5%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: No information Overall symptom improvement: No information Fatigue improvement: Very low certainty ⊕○○○ Functional capacity improvement: No information Strength improvement: Very low certainty ⊕○○○ Adverse events: No information Severe adverse events: No information
Aromatherapy Uncertainty in potential benefits and harms. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					

Hawkins et al (18) ; Peer reviewed; 2022	Patients with post COVID-19 condition (asthenia or fatigue after 150 days of acute COVID-19). 20 assigned to Aromatherapy Twice a day for 14 days and 20 assigned to standard of care.	Male 0%	NR	Low risk of bias	HRQL improvement: Very low certainty ⊕○○○ Overall symptom improvement: No information Fatigue improvement: Very low certainty ⊕○○○ Functional capacity improvement: No information Strength improvement: No information Adverse events: No information Severe adverse events: No information
--	--	---------	----	------------------	---

AXA1125 (amino acids + N-acetylcysteine)

AXA1125 may increase fatigue improvement but may not increase functional capacity improvement. However, certainty of the evidence was low. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
---------------------------	-------------------------------------	---------------	--------------------------	------------------------------------	---

RCT

Finnigan et al (19) ; Peer reviewed; 2022	Patients with post COVID-19 condition (asthenia or fatigue after 84 days of acute COVID-19). 21 assigned to AXA1125 33.9 gr twice a day for 4 weeks and 20 assigned to standard of care.	Mean age 43.6, male 31.7%,	NR	Low risk of bias	HRQL improvement: No information Overall symptom improvement: No information Fatigue improvement: RR 1.07 (95% CI 0.79 to 1.44);
---	--	----------------------------	----	------------------	--

					RD 5.1% (95% CI –16.6% to 34.5%); Low certainty ⊕⊕○○ Functional capacity improvement: RR 0.87 (95% CI 0.51 to 1.48); RD - 8.1% (95% CI – 30% to 29.3%); Low certainty ⊕⊕○○ Strength improvement: No information Adverse events: No information Severe adverse events: No information
--	--	--	--	--	---

Coenzyme Q10 (CQ10)

Uncertainty in potential benefits and harms. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
---------------------------	-------------------------------------	---------------	--------------------------	------------------------------------	---

RCT

Hansen et al. (20); Peer reviewed; 2022	Patients with P-ACC (asthenia or fatigue after 84 days of acute COVID-19). 59 assigned to coenzyme Q10 500 mg a day for 6 weeks and 60 assigned to standard of care.	Median age 49, male 25.2%, obesity 33.6%, interval between COVID-19 and enrolment 288.55 days, hospitalization during COVID-19 15.1%	NR	Low risk of bias	HRQL improvement: Very low certainty ⊕○○○ Overall symptom improvement: Very low certainty ⊕○○○ Fatigue improvement: No information Functional capacity
---	--	--	----	------------------	---

					improvement: No information Strength improvement: No information Adverse events: No information Severe adverse events: No information
--	--	--	--	--	--

Cognitive behavioral therapy

Cognitive behavioral therapy may increase fatigue improvement and functional capacity improvement. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
---------------------------	-------------------------------------	---------------	--------------------------	------------------------------------	---

RCT

Kuut et al. (21) Peer reviewed; 2022	Patients with post COVID-19 condition (asthenia or fatigue after 84 days of acute COVID-19). 57 assigned to Cognitive behavioral therapy for 17 weeks and 57 assigned to standard of care.	Mean age 46, male 27%, interval between COVID-19 and enrolment 188 days, hospitalization during COVID-19 11.5%	NR	High risk of bias Notes: Non-blinded study which might have introduced bias.	HRQL improvement: No information Overall symptom improvement: No information Fatigue improvement: RR 2.2 (95% CI 1.35 to 3.58); RD 31.6% (95% CI 9.2% to 68%); Low certainty ⊕⊕○○ Functional capacity improvement: RR 1.37 (95% CI 1.08 to 1.73);
---	--	--	----	---	---

					RD 22.4% (95% CI 4.8% to 44.7%); Low certainty ⊕⊕○○ Strength improvement: No information Adverse events: No information Severe adverse events: No information
--	--	--	--	--	---

Cytoflavin

Cytoflavin may not improve fatigue. However, certainty of the evidence was low. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
---------------------------	-------------------------------------	---------------	--------------------------	------------------------------------	---

RCT

CITADEL trial (22), Putilina et al.; Peer reviewed; 2022	Patients with P-ACC (asthenia or fatigue after 30 to 90 days of acute COVID-19). 50 assigned to cytoflavin 2 tablets a day for 25 days and 50 assigned to standard of care.	Mean age 40.4 ± 12, male 57%, hypertension 38%, diabetes 4%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: No information Overall symptom improvement: No information Fatigue improvement: RR 1.02 (95% CI 0.98 to 1.06); RD 2.1% (95% CI -1.9% to 6.2%); Low certainty ⊕⊕○○ Functional capacity improvement: No information Strength improvement: No
--	---	---	----	--	--

					<p>information</p> <p>Adverse events: No information</p> <p>Severe adverse events: No information</p>
<p>Enzymes + probiotics</p> <p>Enzymes + probiotics may improve fatigue. However, certainty of the evidence was low. Further research is needed.</p>					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Rathi et al. (23); Peer reviewed; 2022	<p>Patients with P-ACC (asthenia or fatigue after acute COVID-19). 100 assigned to enzymes + probiotics ImmunoSEB (500 mg/capsule) + ProbioSEB CSC3 (5 billion CFUs /capsule) and 100 assigned to standard of care.</p>	<p>Mean age 41.2 ± 13, male 63.5%, interval between COVID-19 and enrolment 19.5 days, one comorbidity 14.5%</p>	NR	<p>High risk of bias</p> <p>Notes: Concealment of allocation and blinding probably inappropriate.</p>	<p>HRQL improvement: No information</p> <p>Overall symptom improvement: No information</p> <p>Fatigue improvement: RR 6.07 (95% CI 3.79 to 9.71); RD 76% (95% CI 41.8% to 85%); Low certainty ⊕⊕○○</p> <p>Functional capacity improvement: No information</p> <p>Strength improvement: No information</p> <p>Adverse events: No information</p>

					Severe adverse events: No information
Fermented food supplements Uncertainty in potential benefits and harms. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Kharaeva et al. (24) ; Peer reviewed; 2022	Patients with P-ACC after moderate infection (asthenia or fatigue after acute COVID-19). 68 assigned to fermented food supplements 14 g twice a day for 20 days and 29 assigned to standard of care.	Age 38–69, male 51.5%, hypertension 36.1%, diabetes 15.5%, chronic lung disease 14.4%, obesity 19.6%, hospitalization during COVID-19 46.4%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: No information Overall symptom improvement: Very low certainty ⊕○○○ Fatigue improvement: No information
Kharaeva et al. (24) ; Peer reviewed; 2022	Patients with P-ACC after severe infection (asthenia or fatigue after 0 days of acute COVID-19). 64 assigned to fermented food supplements 14 g twice a day for 20 days and 27 assigned to standard of care.	Age 36–65, male 47.2%, diabetes 28.6%, chronic lung disease 20.9%, asthma 3.3%, chronic heart disease 37.5%, obesity 40.6%, hospitalization during COVID-19 41.8%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	Functional capacity improvement: No information Strength improvement: No information Adverse events: No information Severe adverse events: No information

Hydrogen (nasal) Uncertainty in potential benefits and harms. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Botek et al. (25); Peer reviewed; 2022	Patients with P-ACC (asthenia or fatigue after 21 to 35 days of acute COVID-19). 26 assigned to hydrogen (nasal) 300 mL/min for 14 days and 24 assigned to standard of care.	Mean age 40, male 52%, interval between COVID-19 and enrolment 25 days	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: No information Overall symptom improvement: No information Fatigue improvement: Very low certainty ⊕○○○ Functional capacity improvement: Very low certainty ⊕○○○ Strength improvement: No information Adverse events: No information Severe adverse events: No information

Immunodaat					
Uncertainty in potential benefits and harms. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Deshpande trial (26) ; Preprint; 2022	Patients with post COVID-19 condition. 26 assigned to Immunodaat 500 mg a day for 30 days and 28 assigned to standard of care.	Mean age 38.9, male 59.4%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: Very low certainty ⊕○○○ Overall symptom improvement: No information Fatigue improvement: Very low certainty ⊕○○○ Functional capacity improvement: No information Strength improvement: No information Adverse events: No information Severe adverse events: No information
Leronlimab					
Uncertainty in potential benefits and harms. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					

Gaylis et al. (27); Peer reviewed; 2022	Patients with P-ACC (asthenia or fatigue after 90 days of acute COVID-19). 27 assigned to Leronlimab 700 mg a week for 8 weeks and 26 assigned to standard of care.	NR	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: No information Overall symptom improvement: Very low certainty ⊕○○○ Fatigue improvement: No information Functional capacity improvement: No information Strength improvement: No information Adverse events: No information Severe adverse events: No information
---	---	----	----	--	---

Mindfulness training

Uncertainty in potential benefits and harms. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
------------------------------	---	---------------	-----------------------------	---------------------------------------	---

RCT

Hauswirth et al (28); Peer reviewed; 2022	Patients with post COVID-19 condition (asthenia or fatigue after 90 days of acute COVID-19). 17	Mean age 47.9, male 26.5%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: No information Overall symptom improvement: No
---	---	---------------------------	----	--	---

	assigned to a mindfulness based intervention (Rebalance®) 2 to 3 sessions (30 min) a week for 4 weeks and 17 assigned to standard of care.				<p>information</p> <p>Fatigue improvement: Very low certainty ⊕○○○</p> <p>Functional capacity improvement: No information</p> <p>Strength improvement: No information Adverse events: No information</p> <p>Severe adverse events: No information</p>
--	--	--	--	--	--

Physical training

Physical training may improve HRQL, fatigue and functional capacity. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Nambi et al. (29); Peer reviewed; 2022	Patients with P-ACC (sarcopenia after acute COVID-19). 36 assigned to aerobic training (high intensity) and 37 assigned to aerobic training (standard intensity).	Mean age 63.5, male 100%	NR	<p>High risk of bias</p> <p>Notes: Non-blinded study which might have introduced bias.</p>	<p>HRQL improvement: RR 1.41 (95% CI 0.81 to 2.58); RD 10% (95% CI -4.3% to 35.8%); Low certainty ⊕⊕○○</p>
Rodriguez-Blanco et al. (30) Peer reviewed; 2022	Patients with P-ACC (asthenia or fatigue after 40 days of acute COVID-19). 24 assigned to endurance training rehabilitation (ETR) (10 breathing and strength-based	Mean age 40.7, male 22.91%	NR	<p>High risk of bias</p> <p>Notes: Non-blinded study. Concealment of allocation probably inappropriate.</p>	<p>Overall symptom improvement: No information</p> <p>Fatigue improvement: RR 4.66 (95% CI 1.96 to 11.09); RD 44.4% (95% CI 11.7% to 87.9%);</p>

	exercises) for 14 days, and 24 assigned to standard of care.				Low certainty ⊕⊕○○
RECOVE trial (31) Jimeno-Almazán et al; Peer reviewed; 2022	Patients with post COVID-19 condition (asthenia or fatigue after 84 days of acute COVID-19). 43 assigned to physical training 3 days a week for 8 weeks and 20 assigned to standard of care.	Mean age 45.3 ± 8, male 31%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	Functional capacity improvement: RR 1.53 (95% CI 0.83 to 2.84); RD 10.9% (95% CI -3.5% to 37.8%); Low certainty ⊕⊕○○ Strength improvement: Very low certainty ⊕○○○
Ibrahim et al. (32) Peer reviewed; 2022	Patients with post COVID-19 condition (asthenia or fatigue after acute COVID-19). 24 assigned to physical training 4 times a week for 10 weeks and 24 assigned to standard of care.	Mean age 63, male 39.6%, diabetes 81.2%, chronic lung disease 66.6%, CHD 83.3%, cancer 12.5%,	NR	High risk of bias Notes: Concealment of allocation and blinding probably inappropriate.	Adverse events: No information Severe adverse events: No information

Phytochemicals

Phytochemicals may improve fatigue and HRQL. However, certainty of the evidence was low. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
UK Phyto-V trial (33) Thomas et al; Peer reviewed; 2022	Patients with P-ACC (asthenia or fatigue after acute COVID-19). 74 assigned to phytochemicals one capsule a day and 73 assigned to standard of care.	Mean age 53, male 56%, obesity 35%, interval between COVID-19 and enrolment 108 days, hospitalization during COVID-19 63%	NR	High risk of bias Notes: Concealment of allocation and blinding probably inappropriate.	HRQL improvement: RR 1.33 (95% CI 1.03 to 1.71); RD 18% (95% CI 1.8% to 39%); Low certainty ⊕⊕○○ Overall symptom improvement: No information

					<p>Fatigue improvement: RR 1.24 (95% CI 0.95 to 1.62); RD 13.1% (95% CI -2.5% to 33.5%); Low certainty ⊕⊕○○</p> <p>Functional capacity improvement: No information</p> <p>Strength improvement: No information</p> <p>Adverse events: No information</p> <p>Severe adverse events: No information</p>
--	--	--	--	--	--

Transcranial direct current stimulation (tDCS)

tDCS may improve fatigue and HRQL, and may not increase adverse events. However, certainty of the evidence was low. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Oliver-Mas et al. (34) ; Preprint; 2022	Patients with P-ACC (asthenia or fatigue after 180 days of acute COVID-19). 23 assigned to transcranial direct current stimulation (tDCS) 1 session a week for 8 weeks and 24 assigned to standard of care.	Mean age 45.6, male 21.3%, hypertension 12.8%, diabetes 4.3%, interval between COVID-19 and enrolment 620 days, hospitalization during COVID-19 14.9%	NR	Low risk of bias	<p>HRQL improvement: RR 1.37 (95% CI 1.09 to 1.71); RD – 26% (95% CI – 6.7% to 30%); Low certainty ⊕⊕○○</p> <p>Overall symptom improvement: No information</p>
Santana et al (35) ; Peer reviewed; 2022	Patients with post COVID-19 condition (asthenia or fatigue after 90 days of acute	Mean age 53, male 35.7%, hypertension 17.1%, diabetes 14.3%, chronic lung disease 5.7%, CHD	NR	Low risk of bias	<p>Fatigue improvement: RR 1.36 (95% CI 0.76 to 2.45); RD – 16.9% (95% CI – 11.2% to 53%);</p>

	COVID-19). 35 assigned to transcranial direct current stimulation (tDCS) 10 sessions and 35 assigned to standard of care.	7.1%, , hospitalization during COVID-19 25.7%			<p>Low certainty ⊕⊕○○</p> <p>Functional capacity improvement: No information</p> <p>Strength improvement: No information</p> <p>Adverse events: RR 0.83 (95% CI 0.26 to 2.73); RD – 3.4% (95% CI – 15.5% to 36%); Low certainty ⊕⊕○○)</p> <p>Severe adverse events: No information</p>
--	---	---	--	--	--

Telerehabilitation

Telerehabilitation may improve fatigue and functional capacity. However, certainty of the evidence was low. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
---------------------------	-------------------------------------	---------------	--------------------------	------------------------------------	---

RCT

King et al. (36); Preprint; 2022	Patients with P-ACC (asthenia or fatigue after 110 days of acute COVID-19). 11 assigned to telerehabilitation twice weekly for 10 weeks and 10 assigned to standard of care.	Mean age 48.5 ± 13, male 47.6%, interval between COVID-19 and enrolment 366 days, hospitalization during COVID-19 19%	NR	<p>High risk of bias</p> <p>Notes: Non-blinded study which might have introduced bias.</p>	<p>HRQL improvement: Very low certainty ⊕○○○</p> <p>Overall symptom improvement: No information</p> <p>Fatigue improvement: RR 1.63 (95% CI 1.04 to 2.54); RD 19.1% (95% CI 1.2% to 46.8%); Low certainty ⊕⊕○○)</p>
Simpson et al (37); Peer reviewed; 2022	Patients with post COVID-19 condition (asthenia or fatigue after 14 days of acute COVID-19). 15 assigned to	Mean age 58 ± 12, male 58%, interval between COVID-19 and enrolment 14 days, hospitalization during COVID-19 100%	NR	<p>High risk of bias</p> <p>Notes: Non-blinded study. Concealment of allocation probably inappropriate.</p>	

	telerehabilitation 45 to 60 min sessions, twice a week for 4 weeks and 12 assigned to standard of care.				Functional capacity improvement: RR 1.48 (95% CI 1.07 to 2.04); RD 13.8% (95% CI 2% to 29.8%); Low certainty ⊕⊕○○)
Longobardi et al. (38) ; Peer reviewed; 2022	Patients with post COVID-19 condition (asthenia or fatigue after 84 days of acute COVID-19). assigned to home physical training 3 times a week for 16 weeks and assigned to standard of care.	Mean age 61, male 50%, hypertension 56%, diabetes 36%, chronic lung disease 16%, CHD 20%, obesity 28%, interval between COVID-19 and enrolment 159 days, hospitalization during COVID-19 100%	NR	High risk of bias Notes: Non-blinded study which might have introduced bias.	Strength improvement: No information Adverse events: No information Severe adverse events: No information
Samper-Pardo et al. (39) ; Peer reviewed; 2022	Patients with post COVID-19 condition (asthenia or fatigue after 84 days of acute COVID-19). 52 assigned to Telerehabilitation through specifically designed app for 12 weeks and 48 assigned to standard of care.	Mean age 48.3 ± 9.26, male 20%, hypertension %, diabetes %, chronic lung disease %, asthma %, CHD %, CKD %, cancer %, obesity %, interval between COVID-19 and enrolment days, hospitalization during COVID-19 %	Corticosteroids %, remdesivir %, hydroxychloroquine %, lopinavir-ritonavir %, tocilizumab %, azithromycin %, convalescent plasma %; Vaccinated %	High risk of bias Notes: Non-blinded study which might have introduced bias.	

Table 3. Description of included studies and interventions effects for P-ACC-related dyspnea

ADAPT-232 (adaptogens) ADAPT-232 may not improve fatigue. However, certainty of the evidence was low. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Karosanidze et al. (16); Peer reviewed; 2022	Patients with P-ACC (asthenia or fatigue after 30 days of acute COVID-19). 49 assigned to ADAPT-232 (adaptogens) 60 mL a day for 14 days and 50 assigned to standard of care.	Mean age 48.9, male 14%	NR	Low risk of bias	HRQL improvement: No information Dyspnea improvement: RR 1. (95% CI 0.94 to 1.06); RD 0% (95% CI – 5.4% to 5.6%); Low certainty ⊕⊕○○ Functional capacity improvement: No information Pulmonary function improvement: No information Radiological response: No information Adverse events: No information Severe adverse events: No information
Antifibrotics Uncertainty in potential benefits and harms. Further research is needed.					

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Kerget et al (40) ; Peer reviewed; 2022	Patients with post COVID-19 condition (dyspnea and/or lung radiological abnormalities after 84 days of acute COVID-19). 15 assigned to pirfenidone 600 to 1800 mg a day for 3 months and 15 assigned to nintendanib 300 mg a day for 3 months	Mean age 65.6, male 40%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: No information Dyspnea improvement: No information Functional capacity improvement: Very low certainty ⊕○○○ Pulmonary function improvement: No information Radiological response: No information Adverse events: No information Severe adverse events: No
Diaphragm release Uncertainty in potential benefits and harms. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					

Nagy et al; (41); Peer reviewed; 2022	Patients with post COVID-19 condition (dyspnea and/or lung radiological abnormalities acute COVID-19). 26 assigned to diaphragm release three sessions a week for 6 weeks and 26 assigned to standard of care.	Mean age 40	NR	High risk of bias Notes: Concealment of allocation and blinding probably inappropriate.	HRQL improvement: No information Dyspnea improvement: Very low certainty ⊕○○○ Functional capacity improvement: No information Pulmonary function improvement: No information Radiological response: No information Adverse events: No information Severe adverse events: No information
---	--	-------------	----	--	--

Nebivolol

Uncertainty in potential benefits and harms. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
------------------------------	---	---------------	-----------------------------	---------------------------------------	---

RCT

Dal Negro et al (42); Peer reviewed; 2022	Patients with post COVID-19 condition (dyspnea and/or lung radiological abnormalities after 84 days of acute COVID-19). 8 assigned to Nebivolol 2.5 mg a day and 8 assigned to standard of care.	Mean age 50.5 ± 17.2, male 63%	NR	High risk of bias Notes: Concealment of allocation and blinding probably inappropriate.	HRQL improvement: No information Dyspnea improvement: Very low certainty ⊕○○○ Functional capacity improvement: No information
---	--	--------------------------------	----	--	--

					Pulmonary function improvement: No information Radiological response: No information Adverse events: No information Severe adverse events: No
Respiratory training/rehabilitation Respiratory training probably improves HRQL and dyspnea, and may improve functional capacity					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
ENO Breathe trial (43), Philip et al.; Peer reviewed; 2022	Patients with P-ACC (dyspnea and/or lung radiological abnormalities after 30 days of acute COVID-19). 58 assigned to ENO Breathe 6-week program and 71 assigned to standard of care.	Mean age 49.5 ± 12, male 17.3%, interval between COVID-19 and enrolment 320 days, hospitalization during COVID-19 17.3%	NR	High risk of bias Notes: Non-blinded study which might have introduced bias.	HRQL improvement: RR 1.47 (95% CI 1.14 to 1.89); RD 20.7% (95% CI 6.2% to 39.3%); Moderate certainty ⊕⊕⊕○ Dyspnea improvement: RR 1.81 (95% CI 1.47 to 2.22); RD 23.3% (95% CI 13.5% to 35.1%); Moderate certainty ⊕⊕⊕○
McNarry et al. (44); Peer reviewed; 2022	Patients with P-ACC (dyspnea and/or lung radiological abnormalities after acute COVID-19). 37 assigned to inspiratory muscle training 3 sessions a week for 8 weeks and 37 assigned to standard of care.	Mean age 46.6 ± 12, male 12.8%, interval between COVID-19 and enrolment 270 days	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate. Intention-to-treat (ITT) analysis for primary outcome not available.	Functional capacity improvement: RR 1.65 (95% CI 1.11 to 2.46); RD 22.7% (95% CI

Srinivasan et al. (45) ; Peer reviewed; 2022	Patients with P-ACC (dyspnea and/or lung radiological abnormalities after acute COVID-19). 24 assigned to respiratory training 3 times a day for 6 weeks and 24 assigned to standard of care.	NR	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	3.8% to 51%); Low certainty ⊕⊕○○ Pulmonary function improvement: Very low certainty ⊕○○○ Radiological response: No information
Rodriguez-Blanco et al. ; (30) Peer reviewed; 2022	Patients with P-ACC (asthenia or fatigue after 40 days of acute COVID-19). 24 assigned to respiratory training (10 breathing and strength-based exercises) for 14 days, and 24 assigned to standard of care.	Mean age 40.7, male 22.91%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	Adverse events: No information Severe adverse events: No
InsCOVID trial (46); Palau et al; Peer reviewed; 2022	Patients with post COVID-19 condition (dyspnea and/or lung radiological abnormalities after 90 days of acute COVID-19). 13 assigned to inspiratory muscle training twice a day for 12 weeks and 13 assigned to standard of care.	Mean age 50.4 ± 12.2, male 58%, hypertension 12%, interval between COVID-19 and enrolment 362 days, hospitalization during COVID-19 100%	NR	High risk of bias Notes: Non-blinded study which might have introduced bias.	
RECOVER trial (47), Romanet et al.; Preprint; 2022	Patients with P-ACC (dyspnea and/or lung radiological abnormalities after 90 days of acute COVID-19). 27 assigned to endurance training rehabilitation (ETR) two (1 h) sessions per week for 10 weeks and 33	Mean age 58.2, male 61.6%, diabetes 36.7%, chronic lung disease 8.3%, chronic heart disease 5%, cancer 5%, interval between COVID-19 and enrolment 173 days, hospitalization during COVID-19 100%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	

	assigned to standard of care.				
RECOVER trial ; (31); Jimeno-Almazán et al; Peer reviewed; 2022	Patients with post COVID-19 condition (dyspnea and/or lung radiological abnormalities after 84 days of acute COVID-19). 40 assigned to respiratory training twice a day for 8 weeks and 20 assigned to standard of care.	Mean age 45.3 ± 8, male 31%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	
Del Corral et al ; (48) Peer reviewed; 2022	Patients with post COVID-19 condition (dyspnea and/or lung radiological abnormalities after 84 days of acute COVID-19). 44 assigned to Respiratory training home-based, 40 minutes a day, 6 days a week for 8 weeks and 44 assigned to standard of care.	Mean age 46.4, male 28.5%, interval between COVID-19 and enrolment 350.7 days, hospitalization during COVID-19 31.8%	Corticosteroids 5.7%	Low risk of bias	
Vallier et al ; (49) Peer reviewed; 2022	Patients with P-ACC (dyspnea and/or lung radiological abnormalities after acute COVID-19). 8 assigned to home pulmonary rehabilitation four times a week for 4 weeks and 9 assigned to inpatient	Mean age 54.8 ± 16, male 70.6%, interval between COVID-19 and enrolment 141 days, hospitalization during COVID-19 76.5%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: Very low certainty ⊕○○○ Dyspnea improvement: Very low certainty ⊕○○○ Functional capacity

	rehabilitation four times a week for 4 weeks				improvement: Very low certainty ⊕○○○ Pulmonary function improvement: No information Radiological response: No information Adverse events: No information Severe adverse events: No information
Simpson et al (37) ; Peer reviewed; 2022	Patients with post COVID-19 condition (asthenia or fatigue after 14 days of acute COVID-19). 15 assigned to telerehabilitation 45 to 60 min sessions, twice a week for 4 weeks and 12 assigned to standard of care.	Mean age 58 ± 12, male 58%, interval between COVID-19 and enrolment 14 days, hospitalization during COVID-19 100%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	Radiological response: No information Adverse events: No information Severe adverse events: No information
Rutkowski et al (50) ; Peer reviewed; 2022	Patients with post COVID-19 condition (dyspnea and/or lung radiological abnormalities after of acute COVID-19). 18 assigned to VR respiratory training five sessions a week for 3 weeks and 14 assigned to conventional respiratory training.	Mean age 57.8 ± 4.9, male 37.5%	NR	High risk of bias Notes: Non-blinded study which might have introduced bias.	HRQL improvement: No information Dyspnea improvement: Very low certainty ⊕○○○ Functional capacity improvement: Very low certainty ⊕○○○
Stavrou et al (51) ; Peer reviewed; 2022	Patients with post COVID-19 condition (dyspnea and/or lung radiological abnormalities after 60 days of acute COVID-19). 10 assigned to VR respiratory training and 10 assigned to standard of care.	Mean age 53.9, male 80%, interval between COVID-19 and enrolment 60 days, hospitalization during COVID-19 100%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	Pulmonary function improvement: No information Radiological response: No information Adverse events: No information Severe adverse events: No information

Kusumawardani et al (52); Peer reviewed; 2022	Patients with post COVID-19 condition (dyspnea and/or lung radiological abnormalities after acute COVID-19). 10 assigned to incentive spirometry 5 times a day for four weeks and 10 assigned to conventional respiratory training.	Mean age 46, male 65%, hypertension 5%, diabetes 5%, obesity 55%, interval between COVID-19 and enrolment 22.5 days, hospitalization during COVID-19 100%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: No information Dyspnea improvement: No information Functional capacity improvement: No information Pulmonary function improvement: Very low certainty ⊕○○○ Radiological response: No information Adverse events: No information Severe adverse events: No information
---	---	---	----	--	--

Steroids (high dose)

High dose steroids may not improve dyspnea and may not increase adverse events. However, certainty of the evidence was low. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
---------------------------	-------------------------------------	---------------	--------------------------	------------------------------------	---

RCT

COLDSTER trial (53); Dhooria et al; Peer reviewed; 2022	Patients with post COVID-19 condition (dyspnea and/or lung radiological abnormalities after 21 to 49 days of acute COVID-19). 65 assigned to High dose steroids Prednisone 40 mg a day descending progressively to 10	Mean age 57, male 68%, one commorbiditie 73%	NR	High risk of bias Notes: Non-blinded study which might have introduced bias.	HRQL improvement: No information Dyspnea improvement: RR 1 (95% CI 0.87 to 1.15); RD 0% (95% CI –11.1% to 12.7%); Low certainty ⊕⊕○○ Functional
---	---	--	----	---	--

	mg a day for 6 weeks and 65 assigned to standard of care.				<p>capacity improvement: No information</p> <p>Pulmonary function improvement: No information</p> <p>Radiological response: Very low certainty ⊕○○○</p> <p>Adverse events: RR 0.92 (95% CI 0.75 to 1.13); RD – 6.2% (95% CI – 19.3% to 10%); Low certainty ⊕⊕○○</p> <p>Severe adverse events: Very low certainty ⊕○○○</p>
--	---	--	--	--	--

Treamid

Treamid may improve dyspnea and pulmonary function, but may not improve functional capacity. Treamid may increase adverse events. However, certainty of the evidence was low. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
---------------------------	-------------------------------------	---------------	--------------------------	------------------------------------	---

RCT

Bazdyrev et al. (54) ; Peer reviewed; 2022	Patients with P-ACC (dyspnea and/or lung radiological abnormalities after acute COVID-19). 29 assigned to treamid 50 mg a day for 28 days and 30 assigned to standard of care.	Mean age 55 ± 11, male 44.1%	NR	Low risk of bias	<p>HRQL improvement: No information</p> <p>Dyspnea improvement: RR 1.96 (95% CI 0.9 to 4.25); RD 21.7% (95% CI –2.3% to 73.7%); Low certainty ⊕⊕○○</p> <p>Functional capacity improvement:</p>
--	--	------------------------------	----	------------------	---

					RR 1.10 (95% CI 0.64 to 1.90); RD 4.3% (95% CI -16.2% to 39.8%); Low certainty ⊕⊕○○ Pulmonary function improvement: RR 2.48 (95% CI 1 to 6.17); RD 24.7% (95% CI 0% to 86.1%); Low certainty ⊕⊕○○ Radiological response: Very low certainty ⊕○○○ Adverse events: RR 1.19 (95% CI 0.56 to 2.50); RD -5.5% (95% CI -12.7% to 43.6%); Low certainty ⊕⊕○○ Severe adverse events: No information
--	--	--	--	--	--

Table 4. Description of included studies and interventions effects for PCC neurocognitive symptoms or sleep disturbances

Actovegin Actovegin may improve cognition. However, certainty of the evidence was low. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					

Kutashov et al (15); Peer reviewed; 2022	Patients with P-ACC (asthenia or fatigue after NR days of acute COVID-19). 222 assigned to Actovegin 1200 mg a day for 60 days and 222 assigned to standard of care.	Mean age 67.6, male 31.98%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: No information Overall symptom improvement: No information Cognitive improvement: RR 1.19 (95% CI 1.06 to 1.33); RD 12.7% (95% CI 4.2% to 22.3%); Low certainty ⊕⊕○○ Depression improvement: No information Adverse events: No information Severe adverse events: No information
--	--	----------------------------	----	--	---

Electric stimulation

Uncertainty in potential benefits and harms. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
---------------------------	-------------------------------------	---------------	--------------------------	------------------------------------	---

RCT

Zulbaran-Rojas et al (55); Peer reviewed; 2022	Patients with post COVID-19 condition (neurocognitive after acute COVID-19). 10 assigned to Electrical stimulation and 8 assigned to standard of care.	Mean age 51.7, male 27.8%, hypertension 44.4%, diabetes 33.3%, interval between COVID-19 and enrolment 299 days, hospitalization during COVID-19 100%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: No information Overall symptom improvement: No information Cognitive improvement: No information Depression improvement: No information
--	--	---	----	--	--

					<p>information</p> <p>Adverse events: No information</p> <p>Severe adverse events: No information</p>
<p>Hyperbaric oxygen (HBO)</p> <p>HBO may improve HRQL. However, certainty of the evidence was low. Further research is needed.</p>					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Zilberman-Itskovich et al. (56); Peer reviewed; 2022	<p>Patients with P-ACC (neurocognitive symptoms after 90 days of acute COVID-19). 37 assigned to HBO 1 session a day for 40 days and 36 assigned to standard of care.</p>	<p>Mean age 48, male 39.7%, hypertension 8.2%, diabetes 2.7%, chronic lung disease 0%, asthma 4.1%, cancer 0%, obesity 27.4%, interval between COVID-19 and enrolment 165 days, hospitalization during COVID-19 16.4%</p>	NR	Low risk of bias	<p>HRQL improvement: RR 1.30 (95% CI 0.84 to 2); RD 13.9% (95% CI -7.4% to 46.9%); Low certainty ⊕⊕○○</p> <p>Overall symptom improvement: No information</p> <p>Cognitive improvement: Very low certainty ⊕○○○</p> <p>Depression improvement: Very low certainty ⊕○○○</p> <p>Adverse events: No information</p> <p>Severe adverse events: No information</p>

Mindfulness training Uncertainty in potential benefits and harms. Further research is needed.					
Study; publication status	Study; publication status	Study; publication status	Study; publication status	Study; publication status	Study; publication status
RCT					
Hausswirth et al (57) ; Peer reviewed; 2022	Patients with post COVID-19 condition (asthenia or fatigue after 90 days of acute COVID-19). 17 assigned to a mindfulness based intervention (Rebalance®) 2 to 3 sessions (30 min) a week for 4 weeks and 17 assigned to standard of care.	Mean age 47.9, male 26.5%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: No information Overall symptom improvement: No information Cognitive improvement: No information Depression improvement: Very low certainty ⊕○○○ Adverse events: No information Severe adverse events: No information
Palmitoylethanolamide + luteolin Uncertainty in potential benefits and harms. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Versace et al (58) ; Peer reviewed; 2022	Patients with post COVID-19 condition (neurocognitive after acute COVID-19). 17 assigned to Palmitoylethanolam	Mean age 50.8, male 35.3%	NR	High risk of bias Notes: pseudo-randomized	HRQL improvement: No information Overall symptom improvement: No information

	ide + Luteolin 1400/400mg a day for 8 weeks and 17 assigned to standard of care.				Cognitive improvement: Very low certainty ⊕○○○ Depression improvement: No information Adverse events: No information Severe adverse events: No information
Physical training Uncertainty in potential benefits and harms. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Kalayeh et al (59) ; Preprint; 2022	Patients with post COVID-19 condition (sleep disturbances after 84 days of acute COVID-19). 17 assigned to endurance training rehabilitation (ETR) Three times a week for eight weeks and 15 assigned to standard of care.	Mean age 25, male 100%, interval between COVID-19 and enrolment 165 days,	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: Very low certainty ⊕○○○ Overall symptom improvement: No information Cognitive improvement: No information Depression improvement: No information Sleep quality improvement: Very low certainty ⊕○○○

					Adverse events: No information Severe adverse events: No information
--	--	--	--	--	--

Transcutaneous auricular vagus nerve stimulation (taVNS)

Uncertainty in potential benefits and harms. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
------------------------------	---	---------------	-----------------------------	---------------------------------------	---

RCT

Badran et al. (60) ; Preprint; 2022	Patients with P-ACC (neurocognitive symptoms after acute COVID-19). 6 assigned to transcutaneous auricular vagus nerve stimulation (taVNS) 2 (1 h) sessions a day for 4 weeks and 6 assigned to standard of care.	Mean age 48.5 ± 11.3, male 33.3%	NR	Low risk of bias	HRQL improvement: No information Overall symptom improvement: No information Cognitive improvement: No information Depression improvement: No information Adverse events: No information Severe adverse events: No information
---	---	----------------------------------	----	------------------	---

Transcranial direct current stimulation (tDCS)

tDCS may not improve fatigue and may not increase adverse events. However, certainty of the evidence was low. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
------------------------------	---	---------------	-----------------------------	---------------------------------------	---

RCT

Oliver-Mas et al. (34) ; Preprint; 2022	Patients with P-ACC (asthenia or fatigue after 180 days of acute COVID-19). 23 assigned to transcranial direct current stimulation (tDCS) 1 session a week for 8 weeks and 24 assigned to standard of care.	Mean age 45.6, male 21.3%, hypertension 12.8%, diabetes 4.3%, interval between COVID-19 and enrolment 620 days, hospitalization during COVID-19 14.9%	NR	Low risk of bias	<p>HRQL improvement: No information</p> <p>Overall symptom improvement: No information</p> <p>Cognitive improvement: RR 0.59 (95% CI 0.33 to 1.05); RD – 27.5% (95% CI – 44.8% to 3.4%); Low certainty ⊕⊕○○</p> <p>Depression improvement: No information</p> <p>Adverse events: No information</p> <p>Severe adverse events: No information</p>
---	---	---	----	------------------	--

Table 5. Description of included studies and interventions effects for PCC olfactory and/or gustatory dysfunction

ADAPT-232 (adaptogens)					
ADAPT-232 may not improve olfactory symptoms. However, certainty of the evidence was low. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Karosanidze et al. (16) ; Peer reviewed; 2022	Patients with P-ACC (asthenia or fatigue after 30 days of acute COVID-19). 49 assigned to ADAPT-232 (adaptogens) 60 mL a day for 14 days and 50 assigned to standard of care.	Mean age 48.9, male 14%	NR	Low risk of bias	<p>HRQL improvement: No information</p> <p>Overall symptom improvement: No information</p> <p>Olfactory symptoms improvement: RR 0.89 (95% CI 0.79 to 1.01); RD – 10.3% (95% CI – 20.5% to 1.4%); Low certainty ⊕⊕○○</p> <p>Gustatory symptoms improvement: No information</p> <p>Adverse events: No information</p> <p>Severe adverse events: No information</p>
Diode laser					
Uncertainty in potential benefits and harms. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					

Shabaan et al ; (61); Peer reviewed; 2022	Patients with post COVID-19 condition (Olfactory and/or gustatory dysfunction after 28 days of acute COVID-19). 18 assigned to Diode laser 6-minute session and 18 assigned to standard of care.	Mean age 41.5, male 22.2%, hospitalization during COVID-19 22.2%	NR	Low risk of bias	HRQL improvement: No information Overall symptom improvement: No information Olfactory symptoms improvement: Very low certainty ⊕○○○ Gustatory symptoms improvement: No information Adverse events: No information Severe adverse events: No information
---	--	--	----	------------------	---

Olfactory training

Uncertainty in potential benefits and harms. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Di Stadio et al. (62); Peer reviewed; 2022	Patients with P-ACC (olfactory and/or gustatory dysfunction after 180 days of acute COVID-19). 76 assigned to olfactory training and 88 assigned to standard of care.	Mean age 40.7, male 27.6%, hypertension 1.7%, diabetes 0%, chronic heart disease 5.2%	Steroids 44%, vitamins 20.7%, alpha lipoic/nicetile 26.7%	High risk of bias Notes: Non-blinded study which might have introduced bias.	HRQL improvement: No information Overall symptom improvement: No information Olfactory symptoms improvement: Very low certainty ⊕○○○ Gustatory symptoms improvement: No
Pires et al. (63); Preprint; 2022	Patients with P-ACC (olfactory and/or gustatory dysfunction after 30 days of acute COVID-19). 26 assigned to	Mean age 37.6, male 35%	Steroids (nasal) 23.8%	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: No information Overall symptom improvement: No information Olfactory symptoms improvement: Very low certainty ⊕○○○ Gustatory symptoms improvement: No

	advanced olfactory training with 8 essential oils: rose, eucalyptus, clove and lemon, citronella, mint, vanilla and cedarwood and 54 assigned to standard of care.				information Adverse events: No information Severe adverse events: No information
COVANOS trial (64) , Lechner et al; Peer reviewed; 2022	Patients with P-ACC (olfactory and/or gustatory dysfunction after 30 days of acute COVID-19). 25 assigned to olfactory training for 12 weeks and 26 assigned to standard of care.	Mean age 44, male 13.8%, hypertension 8.9%, diabetes 1.1%, chronic lung disease 0%, asthma 12.6%, chronic heart disease 0%, cancer 2.1%	NR	High risk of bias Notes: Non-blinded study which might have introduced bias.	

Palmitoylethanolamide + Luteolin

Palmitoylethanolamide + Luteolin may improve olfactory symptoms. However, certainty of the evidence was low. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
---------------------------	-------------------------------------	---------------	--------------------------	------------------------------------	---

RCT

Di Stadio et al; (62) Peer reviewed; 2023	Patients with post COVID-19 condition (Olfactory and/or gustatory dysfunction after 180 days of acute COVID-19). 94 assigned to Palmitoylethanolamide + Luteolin 700/70 mg a day and 36 assigned to standard of care.	Mean age 40.4 ± , male 46.2%, hypertension 2.3%, diabetes 0%	NR	High risk of bias Notes: Concealment of allocation and blinding probably inappropriate.	HRQL improvement: No information Overall symptom improvement: No information Olfactory symptoms improvement: RR 3.11 (95% CI 1.47 to 6.66); RD 35.5% (95% CI 7.8% to 83.3%); Low certainty ⊕⊕○○ Gustatory symptoms
---	---	--	----	--	---

					improvement: No information Adverse events: No information Severe adverse events: No information
Pentasodium diethylenetriamine pentaacetate (DTPA) The effects of DTPA are uncertain. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Imam et al; (65); Peer reviewed; 2022	Patients with post COVID-19 condition (Olfactory and/or gustatory dysfunction after 0 days of acute COVID-19). 33 assigned to DTPA 2% nasal spray, three times a day for one month and 33 assigned to standard of care.	Mean age 39.4, male 42.4%, hypertension 22.7%, diabetes 22.7%, asthma 7.6%, interval between COVID-19 and enrolment 97 days,	Corticosteroids 27.3%	High risk of bias Notes: Concealment of allocation and blinding probably inappropriate.	HRQL improvement: No information Overall symptom improvement: No information Olfactory symptoms improvement: Very low certainty ⊕○○○ Gustatory symptoms improvement: No information Adverse events: No information Severe adverse events: No information
Steroids (nasal) Uncertainty in potential benefits and harms. Further research is needed.					

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
RC 4-7-2020 trial (66), Abdelalim et al.; Peer reviewed; 2022	Patients with P- ACC (olfactory and/or gustatory dysfunction after acute COVID-19). 50 assigned to Mometasone 2 puffs (100 µg) once daily in each nostril for 3 weeks and 50 assigned to standard of care.	Mean age 29, male 46%, hypertension 14%, diabetes 16%, hospitalization during COVID-19 31%	Steroids 13%	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: No information Overall symptom improvement: No information Olfactory symptoms improvement: Very low certainty ⊕○○○ Gustatory symptoms improvement: No information Adverse events: No information Severe adverse events: No information
Steroids Steroids may not improve olfactory nor gustatory symptoms. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Vaira et al. (67); Peer reviewed; 2022	Patients with P- ACC (olfactory and/or gustatory dysfunction after acute COVID-19). 9 assigned to prednisone 1 mg/kg a day and 9	Mean age 42.1, male 38.8%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: No information Overall symptom improvement: No information

	assigned to standard of care.				Olfactory symptoms improvement: RR 1.09 (95% CI 0.83 to 1.44); RD 3.3% (95% CI –6.2% to 16.1%); Low certainty ⊕⊕○○
Schepens et al (68) ; Peer reviewed; 2022	Patients with post COVID-19 condition (Olfactory and/or gustatory dysfunction after 28 days of acute COVID-19). 57 assigned to Prednisone 40 mg a day for 10 days and 56 assigned to standard of care.	Median age 49, male 36.5%, interval between COVID-19 and enrolment 56 days	Vaccinated 79.1%	Low risk of bias	Gustatory symptoms improvement: RR 1.01 (95% CI 0.67 to 1.53); RD 0.5% (95% CI –14.6% to 23.3%); Low certainty ⊕⊕○○ Adverse events: No information Severe adverse events: No information

Theophylline (nasal)

Uncertainty in potential benefits and harms. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
---------------------------	-------------------------------------	---------------	--------------------------	------------------------------------	---

RCT

SCENT2 trial (69) ; Gupta et al; Peer reviewed; 2022	Patients with post COVID-19 condition (Olfactory and/or gustatory dysfunction after 90 days of acute COVID-19). 26 assigned to Theophylline (nasal) 400 mg twice a day for 6 weeks and 25 assigned to standard of care.	Mean age 44.7, male 29.4%, interval between COVID-19 and enrolment 387 days	NR	Low risk of bias	HRQL improvement: Very low certainty ⊕○○○ Overall symptom improvement: No information Olfactory symptoms improvement: Very low certainty ⊕○○○ Gustatory
--	---	---	----	------------------	--

					symptoms improvement: No information Adverse events: No information Severe adverse events: Very low certainty ⊕○○○
--	--	--	--	--	---

Table 6. Description of included studies and interventions effects for PCC cardiovascular system symptoms

Ivabradine Uncertainty in potential benefits and harms. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Jadhav et al. (70) ; Peer reviewed; 2022	Patients with P-ACC (cardiovascular symptoms after 0 to 14 days of acute COVID-19). 25 assigned to Ivabradine 5 to 10 mg and 25 assigned to standard of care.	Mean age 48.8 ± 7.66	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: No information Overall symptom improvement: No information Tachycardia improvement: Very low certainty ⊕○○○ Adverse events: No information Severe adverse events: No information

Table 7. Description of included studies and interventions effects for PCC psychological distress

Virtual reality informational video Virtual reality informational video may improve depression, post-traumatic stress, and psychological distress. However, certainty of the evidence was low. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
ICU-VR trial (71), Vlase et al.; Peer reviewed; 2022	Patients with P-ACC (psychological distress after 90 days of acute COVID-19). 45 assigned to virtual reality 14-minute informational video session once and 44 assigned to standard of care.	Mean age 60, male 36%	NR	High risk of bias Notes: Non-blinded study. Concealment of allocation probably inappropriate.	HRQL improvement: No information Depression improvement: RR 1.21 (95% CI 0.95 to 1.54); RD 14% (95% CI – 3.7% to 36.7%); Low certainty ⊕⊕○○ Post-traumatic stress improvement: RR 1.18 (95% CI 0.98 to 1.42); RD 13.8% (95% CI –1.5% to 32.3%); Low certainty ⊕⊕○○ Psychological distress improvement: RR 1.49 (95% CI 1.08 to 2.05); RD 25.5% (95% CI 4.1% to 55.1%); Low certainty ⊕⊕○○ Adverse events: No information Severe adverse events: No information

Table 8. Description of included studies and interventions effects for P-ACC-related thromboembolic risk

Anticoagulants (prophylactic dose) Anticoagulants may not have an important effect on mortality, VTE, major bleeding and HRQL. However, certainty of the evidence was low because of imprecision. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
MICHELLE trial (72), Ramacciotti et al.; Peer reviewed; 2022	Patients with P-ACC (at increased risk of VTE after acute COVID-19). 159 assigned to rivaroxaban 10 mg a day for 35 days and 159 assigned to standard of care.	Mean age 57.1, male 60%, interval between COVID-19 and enrolment 8 days, hospitalization during COVID-19 100%	NR	High risk of bias Notes: Non-blinded study which might have introduced bias to symptoms, VTE and adverse events outcomes.	Mortality: RR 0.85 (95% CI 0.38 to 0.88); RD -0.3% (95% CI -1.2% to 1.8%); Low certainty ⊕⊕○○ HRQL improvement: RR 0.99 (95% CI 0.78 to 1.24); Low certainty ⊕⊕○○ VTE (symptomatic): RR 1 (95% CI 0.29 to 3.45); RD 0% (95% CI -2.3% to 7.9%); Low certainty ⊕⊕○○ Major bleeding: RR 2.01 (95% CI 0.18 to 22.1); RD 0.1% (95% CI -0.1% to 1.2%); Low certainty ⊕⊕○○ Severe adverse events: No information
ACTIV-4C trial (73); Wang et al; Peer reviewed; 2022	Patients with post COVID-19 condition (thromboembolic events after 0 days of acute COVID-19). 607 assigned to Apixaban 5 mg a day for 30 days and 610 assigned to standard of care.	Median age 54, male 50.5%, hypertension 46.7%, diabetes 28.3%, chronic lung disease 6.1%, asthma 13.7%, CHD 5.6%, interval between COVID-19 and enrolment 0 days, hospitalization during COVID-19 100%	NR	Low risk of bias	

Table 9. Description of included studies and interventions effects for PIMS-TS

Steroids Steroids may reduce time to discharge and respiratory support requirements. However, certainty of the evidence was low. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
Swissped RECOVERY trial (74); Welzel et al; Peer reviewed; 2022	Patients with PIMS-TS. 37 assigned to methylprednisolone 10 mg/kg a day for 3 days and 38 assigned to IVIG 2 gr/kg once	Mean age 9.1, male 75%, underlying chronic disease 11%	NR	High risk of bias Notes: Non-blinded study which might have introduced bias.	Mortality: No information Time to discharge reduction: RR 1.09 (95% CI 0.88 to 1.39); RD 4.5% (95% CI -6% to 19.5%); Low certainty ⊕⊕○○ Respiratory support: RR 0.49 (95% CI 0.27 to 0.89); RD -28.2% (95% CI -40.5% to -5.9%); Low certainty ⊕⊕○○ Inotropic requirements: Very low certainty ⊕○○○ LVEF <55%: Very low certainty ⊕○○○ Arrhythmia: Very low certainty ⊕○○○ VTE: Very low certainty ⊕○○○ Major bleeding: No information

Table 10. Description of included studies and interventions effects for P-ACC prophylaxis

Convalescent plasma					
Convalescent may not reduce P-ACC. However, certainty of the evidence was low. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
CSSC-004 trial (75); Kelly et al; Preprint; 2022	Patients with mild to moderate COVID-19. 445 assigned to convalescent plasma 250 ml once and 437 assigned to standard of care.	Median age 43, male 42.6%, hypertension 23.5%, diabetes 8.2%, obesity 16%,	Vaccinated 22%	High risk of bias Notes: Significant loss to follow-up	Mortality: No information HRQL improvement: No information P-ACC: RR 0.93 (95% CI 0.77 to 1.12); RD -2.4% (95% CI -7.9% to 4.2%); Low certainty ⊕⊕○○ Adverse events: No information Severe adverse events: No information
Fluvoxamine					
Fluvoxamine may not reduce P-ACC. However, the certainty of the evidence was low. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
COVID-OUT trial (76); Bramante et al; Preprint; 2022	Patients with mild to moderate COVID-19. 298 assigned to Fluvoxamine 50 mg once followed by 100 mg a day for 14 days and 297 assigned to	Median age 44.5, male 45.8%, hypertension 26.9%, diabetes 1.1%, obesity 47.2%,	Corticosteroids 1.5%, monoclonal antibodies 4.2%; Vaccinated 56.4%	High risk of bias Notes: Significant loss to follow-up	Mortality: No information HRQL improvement: No information P-ACC: RR 0.99 (95% CI 0.81 to

	standard of care.				1.21); RD -0.4% (95% CI -8.4% to 9.3%); Low certainty ⊕⊕○○
Farahani et al (77) ; Peer reviewed; 2022	Patients with post COVID-19 condition (P-ACC prophylaxis after 0 days of acute COVID-19). 42 assigned to Fluvoxamine 100 mg a day for 10 days and 43 assigned to standard of care.	Mean age 38.5, male 51.2%, hypertension 15.3%, diabetes 6.5%, CHD 11.6%,	Vaccinated 100%	Low risk of bias	Adverse events: No information Severe adverse events: No information

Ivermectin

Ivermectin may not reduce P-ACC. However, certainty of the evidence was low. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
---------------------------	-------------------------------------	---------------	--------------------------	------------------------------------	---

RCT

COVID-OUT - Ivermectin trial (76) ; Bramante et al; Preprint; 2022	Patients with mild to moderate COVID-19. 377 assigned to Ivermectin 390-470 mcg/kg per day for 3 days and 361 assigned to standard of care.	Median age 45.5, male 44%, hypertension 26.7%, diabetes 2%, obesity 48.8%	Corticosteroids 1.5%, monoclonal antibodies 4.2%; Vaccinated 52.2%	High risk of bias Notes: Significant loss to follow-up	Mortality: No information HRQL improvement: No information P-ACC: RR 0.99 (95% CI 0.61 to 1.62); RD 0% (95% CI -1.7% to -2.6%); Low certainty ⊕⊕○○ Adverse events: No information Severe adverse events: No information
--	---	---	--	---	--

Leflunomide

Leflunomide may not reduce P-ACC. However, certainty of the evidence was low. Further research is needed.

Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
---------------------------	-------------------------------------	---------------	--------------------------	------------------------------------	---

RCT					
DEFEAT-COVID trial (78); Kralj-Hans et al; Peer reviewed; 2022	Patients with moderate to severe COVID-19). 81 assigned to leflunomide 100 mg/day for 3 days followed by 20 mg/day for 7 days and 91 assigned to standard of care.	Mean age 55.8, male 67%, diabetes 22%, chronic lung disease 12%, CHD 39%, cancer 3%, obesity 4%, interval between COVID-19 and enrolment 0 days, hospitalization during COVID-19 100%	Corticosteroids 95%, remdesivir %, hydroxychloroquine 47%, tocilizumab 2.3%,	High risk of bias Notes: Non-blinded study which might have introduced bias.	Mortality: No information HRQL improvement: No information P-ACC: RR 1.28 (95% CI 0.92 to 1.77); RD 11.2% (95% CI -3.2% to 31.1%); Low certainty ⊕⊕○○ Adverse events: No information Severe adverse events: Very low certainty ⊕○○○
Metformin					
Metformin may reduce P-ACC. However, certainty of the evidence was low. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
COVID-OUT - Metformin trial (76); Bramante et al; Preprint; 2022	Patients with mild to moderate COVID-19. 564 assigned to metformin 1500 mg a day for 14 days and 561 assigned to standard of care.	Median age 45.5, male 45.3%, hypertension 22.8%, diabetes 1.6%, obesity 47.4%	Steroids 1.5%, remdesivir %, monoclonal antibodies 4.2%; Vaccinated 55.6%	High risk of bias Notes: Significant loss to follow-up	Mortality: No information HRQL improvement: No information P-ACC: RR 0.59 (95% CI 0.39 to 0.88); RD -4.3% (95% CI -6.4% to -1.2%); Low certainty ⊕⊕○○ Adverse events: No information Severe adverse events: No information

Remdesivir Remdesivir may not reduce P-ACC. However, certainty of the evidence was low. Further research is needed.					
Study; publication status	Patients and interventions analyzed	Comorbidities	Additional interventions	Risk of bias and study limitations	Interventions effects vs standard of care (SOC) and GRADE certainty of the evidence
RCT					
SOLIDARITY - Finland trial (79); Nevalainen et al; Peer reviewed; 2022	Patients with post COVID-19 condition (P-ACC prophylaxis after 0 days of acute COVID-19). 98 assigned to Remdesivir 200 mg once followed by 100 mg a day for 10 days and 83 assigned to standard of care.	Mean age 58.4, male 60.2%, diabetes 22.1%, hospitalization during COVID-19 100%	Corticosteroids 71.8%	Low risk of bias	Mortality: Very low certainty ⊕○○○ HRQL improvement: No information P-ACC: RR 1.06 (95% CI 0.53 to 2.13); RD 0.8% (95% CI -6.9% to 16.4%); Low certainty ⊕⊕○○ Adverse events: No information Severe adverse events: No information

References

1. A clinical case definition of post COVID-19 condition by a Delphi consensus. 6 October, 2021. Available from: <https://apps.who.int/iris/rest/bitstreams/1376291/retrieve>
2. A clinical case definition for post COVID-19 condition in children and adolescents by expert consensus (16 February 2023). Available from: <https://www.who.int/publications/i/item/WHO-2019-nCoV-Post-COVID-19-condition-CA-Clinical-case-definition-2023-1>
3. Zhang H, Zang C, Xu Z, Zhang Y, Xu J, Bian J, et al. Data-driven identification of post-acute SARS-CoV-2 infection subphenotypes. *Nat Med* [Internet]. 2022 Dec 1 [cited 2023 Jan 6]; Available from: <https://www.nature.com/articles/s41591-022-02116-3>
4. Nalbandian A, Sehgal K, Gupta A, Madhavan MV, McGroder C, Stevens JS, et al. Post-acute COVID-19 syndrome. *Nat Med*. 2021 Apr;27(4):601–15.
5. Malik P, Patel K, Pinto C, Jaiswal R, Tirupathi R, Pillai S, et al. Post-acute COVID-19 syndrome (PCS) and health-related quality of life (HRQoL)—A systematic review and meta-analysis. *Journal of Medical Virology*. 2022 Jan;94(1):253–62.
6. Han Q, Zheng B, Daines L, Sheikh A. Long-Term Sequelae of COVID-19: A Systematic Review and Meta-Analysis of One-Year Follow-Up Studies on Post-COVID Symptoms. *Pathogens*. 2022 Feb 19;11(2):269.
7. World Health Organization. Clinical management of COVID-19 - Living guideline. Geneva: WHO; 15 September 2022. Available from: <https://reliefweb.int/report/world/clinical-management-covid-19-living-guideline-15-september-2022>
8. The L·OVE Platform. Methods for the special L·OVE of coronavirus infection [Internet] Santiago: Epistemonikos Foundation; 2020 [cited 7 December 2020]. Available from: <https://app.iloveevidence.com/covid-19>
9. Guyatt GH, Thorlund K, Oxman AD, Walter SD, Patrick D, Furukawa TA, et al. GRADE guidelines: 13. Preparing Summary of Findings tables and evidence profiles—continuous outcomes. *Journal of Clinical Epidemiology*. 2013 Feb;66(2):173–83.
10. Hultcrantz M, Rind D, Akl EA, et al. The GRADE Working Group clarifies the construct of certainty of evidence. *J Clin Epidemiol* 2017; 87: 4–13.
11. Zeng L, Brignardello-Petersen R, Hultcrantz M, et al. GRADE guidelines 32: GRADE offers guidance on choosing targets of GRADE certainty of evidence ratings. *Journal of Clinical Epidemiology* 2021; 137: 163–75.

12. Sterne JAC, Savović J, Page MJ, Elbers RG, Blencowe NS, Boutron I, et al. RoB 2: A revised tool for assessing risk of bias in randomised trials. *BMJ* 2019;366:l4898. Available from: <https://doi.org/10.1136/bmj.l4898>.
13. Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008; 336: 924–26.
14. Chudzik M, Burzyńska M, Kapusta J. Use of 1-MNA to Improve Exercise Tolerance and Fatigue in Patients after COVID-19. *Nutrients*. 2022 Jul 22;14(15):3004.
15. Kutashov VA. Actovegin use in patients with cognitive impairment after coronavirus infection (COVID-19). *RJTAO*. 2021 Apr 25;13(2):65–72.
16. Karosanidze I, Kiladze U, Kirtadze N, Giorgadze M, Amashukeli N, Parulava N, et al. Efficacy of Adaptogens in Patients with Long COVID-19: A Randomized, Quadruple-Blind, Placebo-Controlled Trial. *Pharmaceuticals*. 2022 Mar 11;15(3):345.
17. Tosato M, Calvani R, Picca A, Ciciarello F, Galluzzo V, Coelho-Júnior HJ, et al. Effects of L-Arginine Plus Vitamin C Supplementation on Physical Performance, Endothelial Function, and Persistent Fatigue in Adults with Long COVID: A Single-Blind Randomized Controlled Trial. *Nutrients*. 2022 Nov 23;14(23):4984.
18. Hawkins J, Hires C, Keenan L, Dunne E. Aromatherapy blend of thyme, orange, clove bud, and frankincense boosts energy levels in post-COVID-19 female patients: A randomized, double-blinded, placebo controlled clinical trial. *Complementary Therapies in Medicine*. 2022 Aug;67:102823.
19. Finnigan LEM, Cassar MP, Koziel MJ, Pradines J, Lamlum H, Azer K, et al. Efficacy and tolerability of an endogenous metabolic modulator (AXA1125) in fatigue-predominant long COVID: a single-centre, double-blind, randomised controlled phase 2a pilot study. *eClinicalMedicine*. 2023 Apr;101946.
20. Hansen KS, Mogensen TH, Agergaard J, Schiøttz-Christensen B, Østergaard L, Vibholm LK, et al. High-dose coenzyme Q10 therapy versus placebo in patients with post COVID-19 condition: A randomized, phase 2, crossover trial. *The Lancet Regional Health - Europe*. 2022 Nov;100539.
21. Kuut TA, Müller F, Csorba I, Braamse A, Aldenkamp A, Appelman B, et al. Efficacy of cognitive behavioral therapy targeting severe fatigue following COVID-19: results of a randomized controlled trial. *Clinical Infectious Diseases*. 2023 May 8;ciad257.
22. Putilina MV, Teplova NV, Bairova KI, Petrikeeva AE, Shabalina NI. The result of prospective randomized study CITADEL — the efficacy and safety of drug cytoflavin in postcovid rehabilitation. *Z nevrologii i psikiatrii im SS Korsakova*. 2021;121(10):45.

23. Rathi A, Jadhav SB, Shah N. A Randomized Controlled Trial of the Efficacy of Systemic Enzymes and Probiotics in the Resolution of Post-COVID Fatigue. *Medicines*. 2021 Aug 30;8(9):47.
24. Kharaeva Z, Shokarova A, Shomakhova Z, Ibragimova G, Trakhtman P, Trakhtman I, et al. Fermented *Carica papaya* and *Morinda citrifolia* as Perspective Food Supplements for the Treatment of Post-COVID Symptoms: Randomized Placebo-Controlled Clinical Laboratory Study. *Nutrients*. 2022 May 25;14(11):2203.
25. Botek M, Krejčí J, Valenta M, McKune A, Sládečková B, Konečný P, et al. Molecular Hydrogen Positively Affects Physical and Respiratory Function in Acute Post-COVID-19 Patients: A New Perspective in Rehabilitation. *IJERPH*. 2022 Feb 10;19(4):1992.
26. Deshpande S, Mundhe N, Deshpande V, Tamoli S, Mahadik S, Pawar V. Potential use of Immunodaat® (Botanical extract of Elderberry -*Sambucus Nigra* L.) in the management of Post Covid-19 symptoms- a comparative, multi-centric, randomized, clinical study [Internet]. *Allergy and Immunology*; 2022 Oct [cited 2023 Mar 23]. Available from: <http://medrxiv.org/lookup/doi/10.1101/2022.10.04.22280680>
27. Gaylis NB, Ritter A, Kelly SA, Pourhassan NZ, Tiwary M, Sacha JB, et al. Reduced Cell Surface Levels of C-C Chemokine Receptor 5 and Immunosuppression in Long Coronavirus Disease 2019 Syndrome. *Clinical Infectious Diseases*. 2022 Sep 30;75(7):1232–4.
28. Hausswirth C, Schmit C, Rougier Y, Coste A. Positive Impacts of a Four-Week Neuro-Meditation Program on Cognitive Function in Post-Acute Sequelae of COVID-19 Patients: A Randomized Controlled Trial. *IJERPH*. 2023 Jan 11;20(2):1361.
29. Nambi G, Abdelbasset WK, Alrawaili SM, Elsayed SH, Verma A, Vellaiyan A, et al. Comparative effectiveness study of low versus high-intensity aerobic training with resistance training in community-dwelling older men with post-COVID 19 sarcopenia: A randomized controlled trial. *Clin Rehabil*. 2022 Jan;36(1):59–68.
30. Rodriguez-Blanco C, Bernal-Utrera C, Anarte-Lazo E, Gonzalez-Gerez JJ, Saavedra-Hernandez M. A 14-Day Therapeutic Exercise Telerehabilitation Protocol of Physiotherapy Is Effective in Non-Hospitalized Post-COVID-19 Conditions: A Randomized Controlled Trial. *JCM*. 2023 Jan 18;12(3):776.
31. Jimeno-Almazán A, Buendía-Romero Á, Martínez-Cava A, Franco-López F, Sánchez-Alcaraz BJ, Courel-Ibáñez J, et al. Effects of a concurrent training, respiratory muscle exercise, and self-management recommendations on recovery from post-COVID-19 conditions: the RECOVE trial. *Journal of Applied Physiology*. 2023 Jan 1;134(1):95–104

32. Ibrahim AA, Hussein HM, Ali MS, Kanwal R, Acar T, Shaik DH, et al. A randomized controlled trial examining the impact of low vs. moderate-intensity aerobic training in post-discharge COVID-19 older subjects. *European Review for Medical and Pharmacological Sciences*. 2023 May;27(9):4280–91.
33. Thomas R, Williams M, Aldous J, Yanagisawa Y, Kumar R, Forsyth R, et al. A Randomised, Double-Blind, Placebo-Controlled Trial Evaluating Concentrated Phytochemical-Rich Nutritional Capsule in Addition to a Probiotic Capsule on Clinical Outcomes among Individuals with COVID-19—The UK Phyto-V Study. *COVID*. 2022 Mar 22;2(4):433–49.
34. Oliver-Mas S, Delgado-Alonso C, Delgado-Álvarez A, Díez-Cirarda M, Cuevas C, Fernández-Romero L, et al. Transcranial Direct Current Stimulation (tDCS) for Post-COVID Fatigue: A Randomized, Double-Blind, Controlled Pilot Study. *SSRN Journal* [Internet]. 2022 [cited 2022 Dec 21]; Available from: <https://www.ssrn.com/abstract=4216601>
35. Santana K, França E, Sato J, Silva A, Queiroz M, de Farias J, et al. Non-invasive brain stimulation for fatigue in post-acute sequelae of SARS-CoV-2 (PASC). *Brain Stimulation*. 2023 Jan;16(1):100–7.
36. King M, Byrne A, Denehy L, Graham P, Douglas B, de Toni P, et al. Feasibility of a Group-Based Telerehabilitation Intervention for Long COVID Management. [Internet]. In Review; 2022 Mar [cited 2022 Dec 27]. Available from: <https://www.researchsquare.com/article/rs-1452186/v1>
37. Simpson AJ, Green A, Nettleton M, Hyde L, Shepherdson J, Killingback C, et al. Group-based pulmonary telerehabilitation is feasible, safe, beneficial and well-received in patients who have been hospitalised with COVID-19. *ERJ Open Res*. 2023 Mar;9(2):00373–2022.
38. Longobardi I, Goessler K, de Oliveira Júnior GN, Prado DML do, Santos JVP, Meletti MM, et al. Effects of a 16-week home-based exercise training programme on health-related quality of life, functional capacity, and persistent symptoms in survivors of severe/critical COVID-19: a randomised controlled trial. *Br J Sports Med*. 2023 May 10;bjsports-2022-106681.
39. Samper-Pardo M, León-Herrera S, Oliván-Blázquez B, Méndez-López F, Domínguez-García M, Sánchez-Recio R. Effectiveness of a telerehabilitation intervention using ReCOVery APP of long COVID patients: a randomized, 3-month follow-up clinical trial. *Sci Rep*. 2023 May 16;13(1):7943.

40. Kerget B, Çil G, Araz Ö, Alper F, Akgün M. Comparison of two antifibrotic treatments for lung fibrosis in post-COVID-19 syndrome: A randomized, prospective study. *Medicina Clínica*. 2023 Mar;S0025775323000817.
41. Nagy EN, Elimy DA, Ali AY, Ezzelregal HG, Elsayed MM. Influence of Manual Diaphragm Release Technique Combined with Inspiratory Muscle Training on Selected Persistent Symptoms in Men with Post-Covid-19 Syndrome: A Randomized Controlled Trial. *JRM*. 2022 Oct 20;54:jrm00330.
42. Dal Negro RW, Turco P, Povero M. Nebivolol: an effective option against long-lasting dyspnoea following COVID-19 pneumonia - a pivotal double-blind, cross-over controlled study. *Multidis Res Med [Internet]*. 2022 Dec 23 [cited 2023 Mar 23];17. Available from: <https://mrmjournal.org/mrm/article/view/886>
43. Philip KEJ, Owles H, McVey S, Pagnuco T, Bruce K, Brunjes H, et al. An online breathing and wellbeing programme (ENO Breathe) for people with persistent symptoms following COVID-19: a parallel-group, single-blind, randomised controlled trial. *The Lancet Respiratory Medicine*. 2022 Sep;10(9):851–62.
44. McNarry MA, Berg RMG, Shelley J, Hudson J, Saynor ZL, Duckers J, et al. Inspiratory muscle training enhances recovery post-COVID-19: a randomised controlled trial. *Eur Respir J*. 2022 Oct;60(4):2103101.
45. Srinivasan V., Kandakurti P.K., Alagesan J., Suganthirababu P., Jenifer Augustina S., Anitha A., et al. Efficacy of pursed lip breathing with bhastrika pranayama vs incentive spirometry in rehabilitating post Covid 19 follow up-a randomized control study. *Turkish Journal of Physiotherapy and Rehabilitation*. 2021;32(3):402–7.
46. Palau P, Domínguez E, Gonzalez C, Bondía E, Albiach C, Sastre C, et al. Effect of a home-based inspiratory muscle training programme on functional capacity in postdischarged patients with long COVID: the InsCOVID trial. *BMJ Open Resp Res*. 2022 Dec;9(1):e001439.
47. Romanet C, Wormser J, Fels A, Lucas P, Prudat C, Sacco E, et al. Effectiveness of endurance training rehabilitation after hospitalisation in intensive care for COVID-19-related acute respiratory distress syndrome on dyspnoea (RECOVER): a randomised controlled, open-label multicentre trial [Internet]. *Respiratory Medicine*; 2022 Aug [cited 2022 Dec 30]. Available from: <http://medrxiv.org/lookup/doi/10.1101/2022.08.29.22279327>
48. Del Corral T, Fabero-Garrido R, Plaza-Manzano G, Fernández-de-las-Peñas C, Navarro-Santana M, López-de-Uralde-Villanueva I. Home-based respiratory muscle training on

- quality of life and exercise tolerance in long-term post-COVID-19: Randomized controlled trial. *Annals of Physical and Rehabilitation Medicine*. 2023 Feb;66(1):101709.
49. Vallier JM, Simon C, Bronstein A, Dumont M, Jobic A, Paleiron N, et al. Randomized controlled trial of home-based vs. hospital-based pulmonary rehabilitation in post COVID-19 patients. *Eur J Phys Rehabil Med* [Internet]. 2023 Jan [cited 2023 Feb 9]; Available from: <https://www.minervamedica.it/index2.php?show=R33Y9999N00A23012602>
 50. Rutkowski S, Bogacz K, Rutkowska A, Szczegielniak J, Casaburi R. Inpatient post-COVID-19 rehabilitation program featuring virtual reality—Preliminary results of randomized controlled trial. *Front Public Health*. 2023 Feb 6;11:1121554.
 51. Stavrou VT, Vavougios GD, Kalogiannis P, Tachoulas K, Touloudi E, Astara K, et al. Breathlessness and exercise with virtual reality system in long-post-coronavirus disease 2019 patients. *Front Public Health*. 2023 Feb 23;11:1115393.
 52. Kusumawardani RI, Tinduh D, Poerwandari D, Marhana IA, Melaniani S. The effectiveness of incentive spirometry exercise on pulmonary function in COVID-19 survivors: a randomized controlled trial study. *Bali Medical Journal*. 2023;12(1):539–44.
 53. Dhooria S, Chaudhary S, Sehgal IS, Agarwal R, Arora S, Garg M, et al. High-dose versus low-dose prednisolone in symptomatic patients with post-COVID-19 diffuse parenchymal lung abnormalities: an open-label, randomised trial (the COLDSTER trial). *Eur Respir J*. 2022 Feb;59(2):2102930.
 54. Bazdyrev E, Panova M, Brachs M, Smolyarchuk E, Tsygankova D, Gofman L, et al. Efficacy and safety of Treamid in the rehabilitation of patients after COVID-19 pneumonia: a phase 2, randomized, double-blind, placebo-controlled trial [Internet]. In Review; 2022 Jul [cited 2022 Dec 27]. Available from: <https://www.researchsquare.com/article/rs-1845321/v1>
 55. Zulbaran-Rojas A, Lee M, Bara RO, Flores-Camargo A, Spitz G, Finco MG, et al. Electrical stimulation to regain LOWER EXTREMITY muscle perfusion and endurance in patients with POST-ACUTE sequelae of SARS CoV -2: A randomized controlled trial. *Physiological Reports* [Internet]. 2023 Mar [cited 2023 May 5];11(5). Available from: <https://onlinelibrary.wiley.com/doi/10.14814/phy2.15636>
 56. Zilberman-Itskovich S, Catalogna M, Sasson E, Elman-Shina K, Hadanny A, Lang E, et al. Hyperbaric oxygen therapy improves neurocognitive functions and symptoms of post-COVID condition: randomized controlled trial. *Sci Rep*. 2022 Jul 12;12(1):11252.

57. Hausswirth C, Schmit C, Rougier Y, Coste A. Positive Impacts of a Four-Week Neuro-Meditation Program on Cognitive Function in Post-Acute Sequelae of COVID-19 Patients: A Randomized Controlled Trial. *IJERPH*. 2023 Jan 11;20(2):1361.
58. Versace V, Ortelli P, Dezi S, Ferrazzoli D, Alibardi A, Bonini I, et al. Co-ultramicronized palmitoylethanolamide/luteolin normalizes GABAB-ergic activity and cortical plasticity in long COVID-19 syndrome. *Clinical Neurophysiology*. 2023 Jan;145:81–8.
59. Kalayeh MB, Gaeini AA, Kordi MR, Kalayeh MB. Effects of Eight-Week Resistance Training on the Quality of Life and Sleep Quality of Untrained Men with a History of COVID-19 [Internet]. In Review; 2023 Apr [cited 2023 May 5]. Available from: <https://www.researchsquare.com/article/rs-2818393/v1>
60. Badran BW, Huffman SM, Dancy M, Austelle CW, Bikson M, Kautz SA, et al. A pilot randomized controlled trial of supervised, at-home, self-administered transcutaneous auricular vagus nerve stimulation (taVNS) to manage long COVID symptoms [Internet]. In Review; 2022 Jun [cited 2022 Dec 30]. Available from: <https://www.researchsquare.com/article/rs-1716096/v1>
61. Shabaan AA, Kassem I, Mahrous AI, Aboulmagd I, Badrah M, Attalla M, et al. Diode laser in management of loss of taste sensation in patients with post-COVID syndrome: a randomized clinical trial. *BMC Oral Health*. 2023 May 6;23(1):263.
62. Di Stadio A, Cantone E, De Luca P, Di Nola C, Massimilla EA, Motta G, et al. Parosmia COVID-19 Related Treated by a Combination of Olfactory Training and Ultramicronized PEA-LUT: A Prospective Randomized Controlled Trial. *Biomedicine*. 2023 Apr 6;11(4):1109.
63. Pires ÍAT, Steffens ST, Mocelin AG, Shibukawa DE, Leahy L, Saito FL, et al. Intensive Olfactory Training in Post-COVID Patients: A Randomized Multicenter Clinical Trial [Internet]. 2022 Jan [cited 2022 Dec 29]. Available from: <https://preprints.scielo.org/index.php/scielo/preprint/view/3301/version/3492>
64. Lechner M, Liu J, Counsell N, Gillespie D, Chandrasekharan D, Ta NH, et al. The COVANOS trial – insight into post-COVID olfactory dysfunction and the role of smell training. *Rhin*. 2022 Jun 1;60(3):188–99.
65. Imam MS, Abdelazim MH, Abdelazim AH, Ismaiel WF, Gamal M, Abourehab MAS, et al. Efficacy of pentasodium diethylenetriamine pentaacetate in ameliorating anosmia post COVID-19. *American Journal of Otolaryngology*. 2023 Jul;44(4):103871.
66. Abdelalim AA, Mohamady AA, Elsayed RA, Elawady MA, Ghallab AF. Corticosteroid nasal spray for recovery of smell sensation in COVID-19 patients: A randomized controlled trial. *American Journal of Otolaryngology*. 2021 Mar;42(2):102884.

67. Vaira LA, Hopkins C, Petrocelli M, Lechien JR, Cutrupi S, Salzano G, et al. Efficacy of corticosteroid therapy in the treatment of long- lasting olfactory disorders in COVID-19 patients. *Rhin.* 2020 Dec 1;0(0):0–0.
68. Schepens EJA, Blijleven EE, Boek WM, Boesveldt S, Stokroos RJ, Stegeman I, et al. Prednisolone does not improve olfactory function after COVID-19: a randomized, double-blind, placebo-controlled trial. *BMC Med.* 2022 Nov 16;20(1):445.
69. Gupta S, Lee JJ, Perrin A, Khan A, Smith HJ, Farrell N, et al. Efficacy and Safety of Saline Nasal Irrigation Plus Theophylline for Treatment of COVID-19–Related Olfactory Dysfunction: The SCENT2 Phase 2 Randomized Clinical Trial. *JAMA Otolaryngol Head Neck Surg.* 2022 Sep 1;148(9):830.
70. Kartik Pandurang J, Pankaj V J. Ivabradine versus carvedilol in the management of palpitation with sinus tachycardia among recovered COVID-19 patients. *J Cardiol Cardiovasc Me.* 2020 Dec 23;5(3):176–80.
71. Vlakte JH, van Bommel J, Wils EJ, Bienvenu J, Hellemons ME, Korevaar TI, et al. Intensive Care Unit–Specific Virtual Reality for Critically Ill Patients With COVID-19: Multicenter Randomized Controlled Trial. *J Med Internet Res.* 2022 Jan 31;24(1):e32368.
72. Ramacciotti E, Barile Agati L, Calderaro D, Aguiar VCR, Spyropoulos AC, de Oliveira CCC, et al. Rivaroxaban versus no anticoagulation for post-discharge thromboprophylaxis after hospitalisation for COVID-19 (MICHELLE): an open-label, multicentre, randomised, controlled trial. *The Lancet.* 2022 Jan;399(10319):50–9.
73. Wang TY, Wahed AS, Morris A, Kreuziger LB, Quigley JG, Lamas GA, et al. Effect of Thromboprophylaxis on Clinical Outcomes After COVID-19 Hospitalization. *Ann Intern Med.* 2023 Apr;176(4):515–23.
74. Welzel T, Atkinson A, Schöbi N, Andre MC, Bailey DGN, Blanchard-Rohner G, et al. Methylprednisolone versus intravenous immunoglobulins in children with paediatric inflammatory multisystem syndrome temporally associated with SARS-CoV-2 (PIMS-TS): an open-label, multicentre, randomised trial. *The Lancet Child & Adolescent Health.* 2023 Feb;S2352464223000202.
75. Gebo KA, Heath SL, Fukuta Y, Zhu X, Baksh S, Abraham AG, et al. Early Treatment, Inflammation and Post-COVID Conditions [Internet]. *Infectious Diseases (except HIV/AIDS)*; 2023 Feb [cited 2023 Mar 17]. Available from: <http://medrxiv.org/lookup/doi/10.1101/2023.02.13.23285855>
76. Bramante C, Buse JB, Liebovitz D, Nicklas J, Puskarich M, Cohen KR, et al. Outpatient Treatment of COVID-19 and the Development of Long COVID Over 10 Months: A

Multi-Center, Quadruple-Blind, Parallel Group Randomized Phase 3 Trial [Internet]. SSRN; 2023 [cited 2023 Mar 14]. Available from:

<https://www.ssrn.com/abstract=4375620>

77. Farahani RH, Ajam A, Naeini AR. Effect of fluvoxamine on preventing neuropsychiatric symptoms of post COVID syndrome in mild to moderate patients, a randomized placebo-controlled double-blind clinical trial. *BMC Infect Dis.* 2023 Mar 31;23(1):197.
78. Kralj-Hans I, Li K, Wesek A, Lamorgese A, Omar F, Ranasinghe K, et al. Leflunomide treatment for patients hospitalised with COVID-19: DEFEAT-COVID randomised controlled trial. *BMJ Open.* 2023 Apr;13(4):e068179.
79. Nevalainen OPO, Horstia S, Laakkonen S, Rutanen J, Mustonen MJ, Kalliala IEJ, et al. Effect of remdesivir post hospitalization for COVID-19 infection from the randomized SOLIDARITY Finland trial. *Nat Commun.* 2022 Oct 18;13(1):6152.

Annex 1. Summary of findings tables

Summary of findings Table A1.

Population: Patients with P-ACC-related asthenia or fatigue

Intervention: Actovegin

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		SOC	Actovegin		
Fatigue improvement	Relative risk: 1.54 (CI 95% 1.59 - 2.14) Based on data from 444 participants in 1 study Follow up 90 days	471 per 1000	725 per 1000 Difference: 254 more per 1000 (CI 95% 278 more - 537 more)	Low Due to very serious risk of bias ¹	Actovegin may improve fatigue

1. **Risk of Bias: very serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias, Inadequate sequence generation/ generation of comparable groups, resulting in potential for selection bias, Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias; **Imprecision: no serious.** 95% CI include important benefits and harms.

Summary of findings Table A2.

Population: Patients with P-ACC-related asthenia or fatigue

Intervention: ADAPT-232

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the evidence (Quality of evidence)	Plain language summary
		SOC	ADAPT-232		
Fatigue improvement	Relative risk 1.02 (95% CI 0.84 to 1.24) Based on data from 99 participants in 1 study Follow-up 21 days	800 per 1000	816 per 1000 Difference: 16 more per 1000 (95% CI 128 fewer to 192 more)	Low Due to very serious imprecision ¹	Adapt-232 may have little or no difference on fatigue improvement

1. **Imprecision: very serious.** 95% CI includes important benefits and harms.

Summary of findings Table A3.

Population: Patients with P-ACC-related asthenia or fatigue

Intervention: Cytoflavin

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the evidence (Quality of evidence)	Plain language summary
		SOC	Cytoflavin		
Fatigue improvement ^a	Relative risk 1.02 (95% CI 0.98 to 1.06) Based on data from 200 participants in 1 study Follow-up 25 days	979 per 1000	999 per 1000 Difference: 20 more per 1000 (95% CI 20 fewer to 21 more)	Low Due to serious risk of bias, Due to serious imprecision ^b	Cytoflavin may have little or no difference on fatigue improvement

a. Decrease in 12 units of the MFI score.

b. **Risk of bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias; Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias. **Imprecision: serious.** Low number of patients.

Summary of findings Table A4.

Population: Patients with P-ACC-related asthenia or fatigue

Intervention: Enzymes + probiotics

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the evidence (Quality of evidence)	Plain language summary
		SOC	Enzymes + probiotics		
Fatigue improvement	Relative risk 6.07 (95% CI 3.71 to 9.71) Based on data from 200 participants in 1 study Follow-up 25 days	150 per 1000	911 per 1000 Difference: 761 more per 1000 (95% CI 407 more to 850 more)	Low Due to serious risk of bias, Due to serious imprecision ^a	Enzymes + probiotics may increase fatigue improvement

- a. **Risk of bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias; Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias. **Imprecision: serious.** Low number of patients.

Summary of findings Table A5.

Population: Patients with P-ACC-related asthenia or fatigue

Intervention: Phytochemicals

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		SOC	Phytochemicals		
HRQL improvement	Relative risk: 1.33 (CI 95% 1.03 - 1.71) Based on data from 147 participants in 1 study Follow up 30 days	543 per 1000	722 per 1000 Difference: 179 more per 1000 (CI 95% 16 more - 386 more)	Low Due to serious risk of bias, Due to serious imprecision ¹	Phytochemicals may increase HRQL improvement
Fatigue improvement	Relative risk: 1.24 (CI 95% 0.95 - 1.62) Based on data from 147 participants in 1 study Follow up 30 days	539 per 1000	668 per 1000 Difference: 129 more per 1000 (CI 95% 27 fewer - 334 more)	Low Due to serious risk of bias, Due to serious imprecision ²	Phytochemicals may increase fatigue improvement

1. **Risk of Bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias, Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias; **Imprecision: serious.** Low number of patients.
2. **Risk of Bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias, Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias; **Imprecision: serious.** Low number of patients.

Summary of findings Table A6.

Population: Patients with P-ACC-related asthenia or fatigue

Intervention: Transcranial direct current stimulation (tDCS)

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates SOC Transcranial direct current stimulation (tDCS)	Certainty of the Evidence (Quality of evidence)	Plain language summary
Fatigue improvement	Relative risk: 1.36 (CI 95% 0.76 - 2.45) Based on data from 117 participants in 2 studies Follow up 32.5 days	468 636 per 1000 per 1000 Difference: 168 more per 1000 (CI 95% 112 fewer - 672 more)	Low Due to very serious imprecision ¹	Transcranial direct current stimulation (tdcs) may have little or no difference on fatigue improvement
HRQL improvement	Relative risk: 1.37 (CI 95% 1.09 - 1.71) Based on data from 70 participants in 1 studies Follow up 35 days	705 966 per 1000 per 1000 Difference: 261 more per 1000 (CI 95% 63 more - 295 more)	Low Due to very serious imprecision ²	Transcranial direct current stimulation (tdcs) may improve HRQL
Adverse events	Relative risk: 0.83 (CI 95% 0.26 - 2.73) Based on data from 47 participants in 1 studies Follow up 30 days	208 173 per 1000 per 1000 Difference: 35 fewer per 1000 (CI 95% 154 fewer - 360 more)	Low Due to very serious imprecision ³	Transcranial direct current stimulation (tdcs) may have little or no difference on adverse events

1. **Imprecision: very serious.** 95% CI include important benefits and harms;

2. **Imprecision: very serious.** 95% CI include important benefits and harms;

3. **Imprecision: very serious.** 95% CI include important benefits and harms;

Summary of findings Table A7.

Population: Patients with P-ACC-related dyspnea

Intervention: ADAPT-232

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the evidence (Quality of evidence)	Plain language summary
		SOC	ADAPT-232		
Dyspnea improvement	Relative risk 1.0 (95% CI 0.94 to 1.06) Based on data from 99 participants in 1 study Follow-up 21 days	980 per 1000	980 per 1000 Difference: 0 fewer per 1000 (95% CI 59 fewer to 20 more)	Low Due to very serious imprecision ^a	ADAPT-232 may have little or no difference on dyspnea improvement

a. **Imprecision: very serious.** 95% CI includes important benefits and harms.

Summary of findings Table A8.

Population: Patients with P-ACC-related dyspnea

Intervention: Endurance training

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the evidence (Quality of evidence)	Plain language summary
		SOC	Endurance training		
HRQL improvement ^a	Relative risk 1.48 (95% CI 0.92 to 2.37) Based on data from 60 participants in 1 study Follow-up 21 days	441 per 1000	980 per 1000 Difference: 0 fewer per 1000 (95% CI 59 fewer to 20 more)	Low Due to serious risk of bias, Due to serious imprecision ^b	Endurance training may increase HRQL improvement
Dyspnea improvement ^c	Relative risk 2.03 (95% CI 0.98 to 4.21) Based on data from 60 participants in 1 study Follow-up 21 days	236 per 1000	980 per 1000 Difference: 0 fewer per 1000 (95% CI 59 fewer to 20 more)	Low Due to serious risk of bias, Due to serious imprecision ^d	Endurance training may increase dyspnea improvement

a. Increment of 7 units in the SF-12 scale.

b. **Risk of bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias; Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias. **Imprecision: serious.** Low number of patients.

c. Increment of 7 units in the SF-12 scale.

d. **Risk of bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias; Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias. **Imprecision: serious.** Low number of patients.

Summary of findings Table A9.

Population: Patients with P-ACC-related dyspnea

Intervention: High dose steroids (i.e., prednisone 40 mg a day)

Comparator: Standard dose steroids (i.e., prednisone 10 mg a day)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the evidence (Quality of evidence)	Plain language summary
		Standard dose steroids	High dose steroids		
Dyspnea improvement	Relative risk 1.0 (95% CI 0.87 to 1.15) Based on data from 130 participants in 1 study Follow-up 42 days	862 per 1000	862 per 1000	Low Due to serious risk of bias, Due to serious imprecision ^a	High dose steroids may have little or no difference on dyspnea improvement
Radiological response	Relative risk 1.33 (95% CI 0.69 to 2.59) Based on data from 60 participants in 1 study Follow-up 21 days	185 per 1000	246 per 1000	Very low Due to serious risk of bias, Due to very serious imprecision ^b	We are uncertain whether high dose steroids increases or decreases radiological response
Adverse events	Relative risk 0.92 (95% CI 0.75 to 1.13) Based on data from 60 participants in 1 study Follow-up 21 days	769 per 1000	707 per 1000	Low Due to serious risk of bias, Due to serious imprecision ^c	High dose steroids may have little or no difference on adverse events
Severe adverse events	Relative risk 3.0 (95% CI 0.32 to 28.09) Based on data from 60 participants in 1 study Follow-up 21 days	15 per 1000	45 per 1000	Very low Due to serious risk of bias, Due to very serious imprecision ^d	We are uncertain whether high dose steroids increases or decreases severe adverse events

- Risk of bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias; Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias. **Imprecision: serious.** Low number of patients.
- Risk of bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias; Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias. **Imprecision: very serious.** 95% CI includes important benefits and harms.
- Risk of bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias; Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias. **Imprecision: serious.** Low number of patients.
- Risk of bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias; Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias. **Imprecision: very serious.** 95% CI includes important benefits and harms.

Summary of findings Table A10.

Population: Patients with P-ACC-related dyspnea

Intervention: Respiratory training/rehabilitation

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		SOC	Respiratory training		
HRQL improvement	Relative risk: 1.47 (CI 95% 1.14 - 1.89) Based on data from 351 participants in 5 studies Follow up 88 days	441 per 1000	648 per 1000 Difference: 207 more per 1000 (CI 95% 62 more - 392 more)	Moderate Due to serious risk of bias ¹	Respiratory training probably increases HRQL improvement
Functional capacity improvement	Relative risk: 1.65 (CI 95% 1.11 - 2.46) Based on data from 88 participants in 2 studies Follow up 56 days	349 per 1000	576 per 1000 Difference: 227 more per 1000 (CI 95% 38 more - 510 more)	Low Due to very serious imprecision ²	Respiratory training may increase functional capacity improvement
Dyspnea improvement	Relative risk: 1.81 (CI 95% 1.47 - 2.22) Based on data from 474 participants in 8 studies Follow up 75 days	287 per 1000	519 per 1000 Difference: 232 more per 1000 (CI 95% 135 more - 350 more)	Moderate Due to serious risk of bias ³	Respiratory training probably increases dyspnea improvement
Pulmonary function improvement	Relative risk: 1.39 (CI 95% 0.8 - 2.41) Based on data from 74 participants in 2 studies Follow up 66 days	459 per 1000	638 per 1000 Difference: 179 more per 1000 (CI 95% 92 fewer - 647 more)	Very low Due to serious risk of bias, Due to very serious imprecision ⁴	We are uncertain whether respiratory training increases or decreases pulmonary function improvement

1. **Risk of Bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias, Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias;
2. **Imprecision: very serious.** 95% CI include important benefits and harms;
3. **Risk of Bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias, Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias;
4. **Risk of Bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias, Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias;
Imprecision: very serious. 95% CI include important benefits and harms;

Summary of findings Table A11.

Population: Patients with P-ACC-related dyspnea

Intervention: Treamid

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the evidence (Quality of evidence)	Plain language summary
		SOC	Treamid		
Functional capacity improvement	Relative risk 1.1 (95% CI 0.64 to 1.9) Based on data from 59 participants in 1 study Follow-up 28 days	445 per 1000	490 per 1000	Low Due to very serious imprecision ^a	Treamid may have little or no difference on functional capacity improvement
Dyspnea improvement	Relative risk 1.96 (95% CI 0.9 to 4.25) Based on data from 59 participants in 1 study Follow-up 28 days	227 per 1000	445 per 1000	Low Due to very serious imprecision ^b	Treamid may increase dyspnea improvement
Pulmonary function improvement	Relative risk 2.48 (95% CI 1.0 to 6.17) Based on data from 59 participants in 1 study Follow-up 28 days	167 per 1000	414 per 1000	Low Due to very serious imprecision ^c	Treamid may increase pulmonary function improvement
Adverse events	Relative risk 1.19 (95% CI 0.56 to 2.5) Based on data from 59 participants in 1 study Follow-up 28 days	290 per 1000	345 per 1000	Low Due to very serious imprecision ^d	Treamid may increase adverse events

a. **Imprecision: very serious.** 95% CI includes important benefits and harms.

b. **Imprecision: very serious.** 95% CI includes important benefits and harms.

c. **Imprecision: very serious.** 95% CI includes important benefits and harms.

d. **Imprecision: very serious.** 95% CI includes important benefits and harms.

Summary of findings Table A12.

Population: Patients with P-ACC-related neurocognitive symptoms or sleep disturbances

Intervention: Actovegin

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		SOC	Actovegin		
Cognitive improvement	Odds ratio: 1.19 (CI 95% 1.06 - 1.33) Based on data from 444 participants in 1 study	673 per 1000	710 per 1000 Difference: 37 more per 1000 (CI 95% 13 more - 384 fewer)	Low Due to very serious risk of bias ¹	Actovegin may improve cognition

1. **Risk of Bias: very serious.** Inadequate sequence generation/ generation of comparable groups, resulting in potential for selection bias, Inadequate concealment of allocation during randomization process, resulting in potential for selection bias, Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias, Inadequate/lack of blinding of outcome assessors, resulting in potential for detection bias; **Indirectness: serious.** Non appropriately established MID; **Imprecision: no serious.** 95% CI include important benefits and harms.

Summary of findings Table A13.

Population: Patients with P-ACC-related neurocognitive symptoms or sleep disturbances

Intervention: Hyperbaric oxygen (HBO)

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the evidence (Quality of evidence)	Plain language summary
		SOC	HBO		
HRQL improvement	Relative risk 1.3 (95% CI 0.84 to 2.0) Based on data from 73 participants in 1 study	469 per 1000	610 per 1000	Low Due to very serious imprecision ^a	HBO may increase HRQL improvement
		Difference: 141 more per 1000 (95% CI 75 fewer to 469 more)			
Cognitive improvement	Odds ratio 2.84 (95% CI 1.09 to 7.37) Based on data from 73 participants in 1 study	667 per 1000	850 per 1000	Very low Due to extremely serious imprecision, Due to serious indirectness ^b	We are uncertain whether HBO increases or decreases cognitive improvement
		Difference: 183 more per 1000 (95% CI 19 more to 22 more)			
Depression improvement	Odds ratio 35.9 (95% CI 2.72 to 474.6) Based on data from 73 participants in 1 study Follow-up 28 days	681 per 1000	987 per 1000	Very low Due to extremely serious imprecision, Due to serious indirectness ^c	We are uncertain whether HBO increases or decreases depression improvement
		Difference: 306 more per 1000 (95% CI 172 more to 312 more)			

a. **Imprecision: very serious.** 95% CI includes important benefits and harms.

b. **Indirectness: serious.** Non appropriately established minimal important difference (MID). **Imprecision: extremely serious.** 95% CI includes important benefits and harms.

c. **Indirectness: serious.** Non appropriately established MID. **Imprecision: extremely serious.** 95% CI includes important benefits and harms.

Summary of findings Table A14.

Population: Patients with P-ACC-related neurocognitive symptoms or sleep disturbances

Intervention: Transcranial direct current stimulation (tDCS)

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the evidence (Quality of evidence)	Plain language summary
		SOC	Transcranial direct current stimulation (tDCS)		
Cognitive improvement	Relative risk 0.59 (95% CI 0.33 to 1.05) Based on data from 47 participants in 1 study Follow-up 30 days	667 per 1000	394 per 1000 Difference: 273 fewer per 1000 (95% CI 447 fewer to 33 more)	Low Due to very serious imprecision ^a	tDCS may have little or no difference on cognitive improvement

a. **Imprecision: very serious.** 95% CI includes important benefits and harms.

Summary of findings Table A15.

Population: Patients with P-ACC-related olfactory and/or gustatory dysfunction

Intervention: ADAPT-232

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the evidence (Quality of evidence)	Plain language summary
		SOC	ADAPT-232		
Olfactory symptoms improvement	Relative risk 0.89 (95% CI 0.79 to 1.01) Based on data from 99 participants in 1 study Follow-up 21 days	960 per 1000	854 per 1000 Difference: 106 fewer per 1000 (95% CI 202 fewer to 10 more)	Low Due to very serious imprecision ^a	ADAPT-232 may have little or no difference on olfactory symptoms

a. **Imprecision: very serious.** 95% CI includes important benefits and harms.

Summary of findings Table A16.

Population: Patients with P-ACC-related olfactory and/or gustatory dysfunction

Intervention: Palmitoylethanolamide + Luteolin

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		SOC	Palmitoylethanolamide + Luteolin		
Olfactory symptoms improvement	Relative risk: 3.13 (CI 95% 1.47 - 666.0) Based on data from 130 participants in 1 studies Follow up 90 days	167 per 1000	523 per 1000 Difference: 356 more per 1000 (CI 95% 78 more - 833 more)	Low Due to very serious imprecision ¹	Palmitoylethanolamide + luteolin may increase olfactory symptoms improvement

1. **Imprecision: very serious.** 95% CI include important benefits and harms;

Summary of findings Table A17.

Population: Patients with P-ACC-related olfactory and/or gustatory dysfunction

Intervention: Steroids

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		SOC	Steroids		
Olfactory symptoms improvement	Relative risk: 1.09 (CI 95% 0.83 - 1.44) Based on data from 131 participants in 2 studies Follow up 52 days	365 per 1000	398 per 1000 Difference: 33 more per 1000 (CI 95% 62 fewer - 161 more)	Low Due to very serious imprecision ¹	Steroids may have little or no difference on olfactory symptoms
Gustatory symptoms improvement	Relative risk: 1.01 (CI 95% 0.67 - 1.53) Based on data from 113 participants in 1 study Follow up 84 days	443 per 1000	447 per 1000 Difference: 4 more per 1000 (CI 95% 146 fewer - 235 more)	Low Due to serious risk of bias, Due to serious imprecision ²	Steroids may have little or no difference on gustatory symptoms

1. **Imprecision: very serious.** Low number of patients, Wide confidence intervals.

2. **Risk of Bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias, Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias;
Imprecision: serious. Low number of patients.

Summary of findings Table A18.

Population: Patients with P-ACC-related psychological distress

Intervention: Virtual reality informational video

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		SOC	Virtual informational video		
Depression improvement	Relative risk 1.21 (95% CI 0.95 to 1.54) Based on data from 89 participants in 1 study Follow-up 90 days	682 per 1000	825 per 1000	Low Due to serious risk of bias, Due to serious imprecision ^a	Virtual reality informational video may increase depression improvement
Post-traumatic stress disorder improvement	Relative risk 1.18 (95% CI 0.98 to 1.42) Based on data from 89 participants in 1 study Follow-up 90 days	773 per 1000	912 per 1000	Low Due to serious risk of bias, Due to serious imprecision ^b	Virtual reality informational video may increase post-traumatic stress disorder improvement
Psychologic distress improvement	Relative risk 1.49 (95% CI 1.08 to 2.05) Based on data from 89 participants in 1 study Follow-up 90 days	523 per 1000	779 per 1000	Low Due to serious risk of bias, Due to serious imprecision ^c	Virtual reality informational video may increase psychological distress improvement

a. **Risk of bias: serious. Imprecision: serious.** Low number of patients.

b. **Risk of bias: serious. Imprecision: serious.** Low number of patients.

c. **Risk of bias: serious. Imprecision: serious.** Low number of patients.

Summary of findings Table A19.

Population: Patients with P-ACC-related thromboembolic risk

Intervention: Anticoagulants

Comparator: Standard of care

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		SOC	Anticoagulants		
HRQL improvement	Relative risk: 0.99 (CI 95% 0.78 - 1.24) Based on data from 1217 participants in 1 study Follow up 90 days			Low Due to very serious imprecision ¹	Anticoagulants may have little or no difference on HRQL
VTE	Relative risk: 1.0 (CI 95% 0.29 - 3.45) Based on data from 1535 participants in 2 studies Follow up 32.5 days	32 per 1000	32 per 1000 Difference: 0 fewer per 1000 (CI 95% 23 fewer - 78 more)	Low Due to very serious imprecision ²	Anticoagulants may have little or no difference on VTE
Mortality	Relative risk: 0.85 (CI 95% 0.38 - 1.88) Based on data from 1535 participants in 2 studies Follow up 32.5 days	20 per 1000	17 per 1000 Difference: 3 fewer per 1000 (CI 95% 12 fewer - 18 more)	Low Due to very serious imprecision ³	Anticoagulants may have little or no difference on mortality
Major bleeding	Relative risk: 2.01 (CI 95% 0.18 - 22.1) Based on data from 1535 participants in 2 studies Follow up 32.5 days	1 per 1000	2 per 1000 Difference: 1 more per 1000 (CI 95% 1 fewer - 21 more)	Low Due to very serious imprecision ⁴	Anticoagulants may have little or no difference on major bleeding

1. **Imprecision: very serious.** Low number of patients, Wide confidence intervals.

2. **Imprecision: very serious.** Low number of patients, Wide confidence intervals.

3. **Imprecision: very serious.** Low number of patients, Wide confidence intervals.

4. **Imprecision: very serious.** Low number of patients, Wide confidence intervals.

Summary of findings Table A20.

Population: Patients with PIMS-TS

Intervention: Steroids

Comparator: IVIG

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		IVIG	Steroids		
Time to discharge time reduction ¹	Relative risk: 1.09 (CI 95% 0.88 - 1.39) Based on data from 75 participants in 1 study Follow up 28	500 per 1000	545 per 1000 Difference: 45 more per 1000 (CI 95% 60 fewer - 195 more)	Low Due to serious risk of bias, Due to serious imprecision ²	Steroids may decrease time to discharge
Respiratory support	Relative risk: 0.49 (CI 95% 0.27 - 0.89) Based on data from 75 participants in 1 study Follow up 28	553 per 1000	271 per 1000 Difference: 282 fewer per 1000 (CI 95% 404 fewer - 61 fewer)	Low Due to serious risk of bias, Due to serious imprecision ³	Steroids may decrease respiratory support requirements
Inotropic requirements	Relative risk: 0.68 (CI 95% 0.35 - 1.32) Based on data from 75 participants in 1 study Follow up 28	395 per 1000	269 per 1000 Difference: 126 fewer per 1000 (CI 95% 257 fewer - 126 more)	Very low Due to serious risk of bias, Due to serious imprecision, Due to very serious imprecision ⁴	We are uncertain whether steroids increases or decreases inotropic requirements
Left ventricular fraction deterioration	Relative risk: 0.57 (CI 95% 0.21 - 1.54) Based on data from 75 participants in 1 study Follow up 28	237 per 1000	135 per 1000 Difference: 102 fewer per 1000 (CI 95% 187 fewer - 128 more)	Very low Due to serious risk of bias, Due to serious imprecision, Due to very serious imprecision ⁵	We are uncertain whether steroids increases or decreases LVEF deterioration
Arrhythmia	Relative risk: 2.05 (CI 95% 0.19 - 21.7) Based on data from 75 participants in 1 study Follow up 28	26 per 1000	53 per 1000 Difference: 27 more per 1000 (CI 95% 21 fewer - 538 more)	Very low Due to serious risk of bias, Due to serious imprecision, Due to very serious imprecision ⁶	We are uncertain whether steroids increases or decreases Arrhythmias
Venous thromboembolic events	Relative risk: 0.34 (CI 95% 0.01 - 8.14) Based on data from 75 participants in 1 study Follow up 28	39 per 1000	13 per 1000 Difference: 26 fewer per 1000 (CI 95% 39 fewer - 278 more)	Very low Due to serious risk of bias, Due to serious imprecision, Due to very serious imprecision ⁷	We are uncertain whether steroids increases or decreases VTE

1. Proportion of patients discharged on day 6.
2. **Risk of Bias: serious.** Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias, Inadequate/lack of blinding of outcome assessors, resulting in potential for detection bias; **Imprecision: serious.** Wide confidence intervals.
3. **Risk of Bias: serious.** Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias, Inadequate/lack of blinding of outcome assessors, resulting in potential for detection bias; **Imprecision: serious.** Wide confidence intervals.
4. **Risk of Bias: serious.** Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias, Inadequate/lack of blinding of outcome assessors, resulting in potential for detection bias; **Imprecision: very serious.** Wide confidence intervals.
5. **Risk of Bias: serious.** Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias, Inadequate/lack of blinding of outcome assessors, resulting in potential for detection bias; **Imprecision: very serious.** Wide confidence intervals.

6. **Risk of Bias: serious.** Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias, Inadequate/lack of blinding of outcome assessors, resulting in potential for detection bias; **Imprecision: very serious.** Wide confidence intervals.
7. **Risk of Bias: serious.** Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias, Inadequate/lack of blinding of outcome assessors, resulting in potential for detection bias; **Imprecision: very serious.** Wide confidence intervals.

Summary of findings Table A21.

Population: Patients with COVID-19

Intervention: Metformin to prevent P-ACC

Comparator: SOC

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		SOC	Metformin to prevent P-ACC		
P-ACC	Relative risk: 0.59 (CI 95% 0.39 - 0.88) Based on data from 1125 participants in 1 study Follow up 300 days	105 per 1000	62 per 1000 Difference: 43 fewer per 1000 (CI 95% 64 fewer - 13 fewer)	Low Due to serious risk of bias, Due to serious imprecision ¹	Metformin may reduce P-ACC

1. **Risk of Bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias, Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias;
Imprecision: serious. Low number of patients.

Summary of findings Table A22.

Population: Patients with COVID-19
Intervention: Ivermectin to prevent P-ACC
Comparator: SOC

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		SOC	Ivermectin to prevent P-ACC		
P-ACC	Relative risk: 0.99 (CI 95% 0.61 - 1.62) Based on data from 738 participants in 1 study Follow up 300 days	105 per 1000	104 per 1000 Difference: 1 fewer per 1000 (CI 95% 41 fewer - 65 more)	Low Due to serious risk of bias, Due to serious imprecision ¹	Ivermectin may not reduce P-ACC

1. **Risk of Bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias, Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias;
Imprecision: serious. Low number of patients.

Summary of findings Table A23.

Population: Patients with COVID-19

Intervention: Convalescent plasma to prevent P-ACC

Comparator: SOC

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		SOC	CP to prevent P-ACC		
P-ACC	Relative risk: 0.93 (CI 95% 0.77 - 1.12) Based on data from 882 participants in 1 study Follow up 90 days	343 per 1000	319 per 1000 Difference: 24 fewer per 1000 (CI 95% 79 fewer - 41 more)	Low Due to serious risk of bias, Due to serious imprecision ¹	Convalescent plasma may not reduce P-ACC

1. **Risk of Bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias, Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias;
Imprecision: serious. Low number of patients.

Summary of findings Table A24.

Population: Patients with COVID-19

Intervention: Remdesivir to prevent P-ACC

Comparator: SOC

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		SOC	Remdesivir to prevent P-ACC		
Mortality	Relative risk: 0.85 (CI 95% 0.25 - 2.83) Based on data from 181 participants in 1 study Follow up 365 days	60 per 1000	51 per 1000 Difference: 9 fewer per 1000 (CI 95% 45 fewer - 110 more)	Very low Due to serious risk of bias, Due to very serious imprecision	We are uncertain whether remdesivir to prevent p-acc increases or decreases mortality
P-ACC	Relative risk: 1.06 (CI 95% 0.53 - 2.13) Based on data from 181 participants in 1 study Follow up 365 days	145 per 1000	154 per 1000 Difference: 9 more per 1000 (CI 95% 68 fewer - 164 more)	Low Due to serious risk of bias, Due to serious imprecision	Remdesivir may not reduce P-ACC

1. **Risk of Bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias, Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias; **Imprecision: very serious.** Low number of patients, Wide confidence intervals.
2. **Risk of Bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias, Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias; **Imprecision: serious.** Low number of patients.

Summary of findings Table A25.

Population: Patients with COVID-19

Intervention: Leflunomide to prevent P-ACC

Comparator: SOC

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		SOC	Leflunomide to prevent P-ACC		
P-ACC	Relative risk: 1.28 (CI 95% 0.92 - 1.77) Based on data from 172 participants in 1 study Follow up 90 days	407 per 1000	521 per 1000 Difference: 114 more per 1000 (CI 95% 33 fewer - 313 more)	Low Due to serious risk of bias, Due to serious imprecision ¹	Remdesivir may not reduce P-ACC
Severe adverse events	Relative risk: 1.76 (CI 95% 0.81 - 3.85) Based on data from 214 participants in 1 study Follow up 90 days	82 per 1000	144 per 1000 Difference: 62 more per 1000 (CI 95% 16 fewer - 234 more)	Very low Due to serious risk of bias, Due to very serious imprecision ²	We are uncertain whether remdesivir to prevent p- acc increases or decreases mortality

1. **Risk of Bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias, Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias; **Imprecision: serious.** Low number of patients.
2. **Risk of Bias: serious.** Inadequate concealment of allocation during randomization process, resulting in potential for selection bias, Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias; **Imprecision: very serious.** Low number of patients, Wide confidence intervals.

Summary of findings Table A26.

Population: Patients with COVID-19

Intervention: Fluvoxamine to prevent P-ACC

Comparator: SOC

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		SOC	Fluvoxamine to prevent P-ACC		
P-ACC	Relative risk: 0.99 (CI 95% 0.81 - 1.21) Based on data from 680 participants in 2 studies Follow up 192 days	444 per 1000	440 per 1000 Difference: 4 fewer per 1000 (CI 95% 84 fewer - 93 more)	Low Due to very serious imprecision ¹	Fluvoxamine may not reduce P-ACC

5. **Imprecision: very serious.** Wide confidence intervals.

Summary of findings Table A27.

Population: Patients with P-ACC-related asthenia or fatigue

Intervention: AXA1125

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		SOC	AXA1125		
Fatigue improvement	Relative risk: 1.07 (CI 95% 0.79 - 1.44) Based on data from 41 participants in 1 study Follow up 28 days	780 per 1000	835 per 1000 Difference: 55 more per 1000 (CI 95% 164 fewer - 343 more)	Low Due to very serious imprecision ¹	Axa1125 may increase fatigue improvement
Functional capacity improvement	Relative risk: 0.87 (CI 95% 0.51 - 1.48) Based on data from 41 participants in 1 study Follow up 28 days	607 per 1000	528 per 1000 Difference: 79 fewer per 1000 (CI 95% 297 fewer - 291 more)	Low Due to very serious imprecision ²	Axa1125 may not increase functional capacity improvement
Adverse events	Relative risk: 2.62 (CI 95% 1.0 - 6.89) Based on data from 41 participants in 1 study Follow up 28 days	200 per 1000	524 per 1000 Difference: 324 more per 1000 (CI 95% 0 fewer - 800 more)	Very low Due to extremely serious imprecision ³	We are uncertain whether axa1125 improves or worsen adverse events

1. **Imprecision: very serious.** 95% CI include important benefits and harms.

2. **Imprecision: very serious.** 95% CI include important benefits and harms.

3. **Imprecision: ~extreme_serious.** 95% CI include important benefits and harms.

Summary of findings Table A28.

Population: Patients with P-ACC-related asthenia or fatigue

Intervention: Telerehabilitation

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		SOC	Telerehabilitation		
HRQL improvement	Relative risk: 1.49 (CI 95% 0.6 - 3.66) Based on data from 48 participants in 2 studies Follow up 63 days	256 per 1000	966 per 1000	Very low Due to very serious imprecision, Due to serious risk of bias ¹	We are uncertain whether telerehabilitation increases or decreases hrql improvement
Fatigue improvement	Relative risk: 1.63 (CI 95% 1.04 - 2.54) Based on data from 89 participants in 3 studies Follow up 79 days	304 per 1000	636 per 1000	Low Due to serious risk of bias, Due to serious imprecision ²	Telerehabilitation may increase fatigue improvement
Functional capacity improvement	Relative risk: 2.11 (CI 95% 1.11 - 3.99) Based on data from 62 participants in 2 studies Follow up 91 days	281 per 1000	593 per 1000	Low Due to serious risk of bias, Due to serious imprecision ³	Telerehabilitation may increase functional capacity improvement

1. **Risk of Bias: serious.** Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias, Inadequate/lack of blinding of outcome assessors, resulting in potential for detection bias; **Imprecision: very serious.** 95% CI include important benefits and harms, Wide confidence intervals;
2. **Risk of Bias: serious.** Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias, Inadequate/lack of blinding of outcome assessors, resulting in potential for detection bias; **Imprecision: serious.** Low number of patients;
3. **Risk of Bias: serious.** Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias, Inadequate/lack of blinding of outcome assessors, resulting in potential for detection bias; **Imprecision: serious.** 95% CI include important benefits and harms, Low number of patients;

Summary of findings Table A29.

Population: Patients with P-ACC-related asthenia or fatigue

Intervention: Cognitive behavioral therapy

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		SOC	Cognitive behavioral therapy		
Fatigue improvement	Relative risk: 2.2 (CI 95% 1.35 - 3.58) Based on data from 114 participants in 1 studies Follow up 119 days	263 per 1000	636 per 1000 Difference: 168 more per 1000 (CI 95% 112 fewer - 672 more)	Low Due to serious risk of bias, Due to serious imprecision ¹	Cognitive behavioral therapy may increase fatigue improvement
Functional capacity improvement	Relative risk: 1.37 (CI 95% 1.08 - 1.73) Based on data from 114 participants in 1 studies Follow up 119 days	614 per 1000	841 per 1000 Difference: 227 more per 1000 (CI 95% 49 more - 448 more)	Low Due to serious risk of bias, Due to serious imprecision ²	Cognitive behavioral therapy may increase functional capacity improvement

1. **Risk of Bias: serious.** Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias, Inadequate/lack of blinding of outcome assessors, resulting in potential for detection bias; **Imprecision: serious.** Low number of patients;
2. **Risk of Bias: serious.** Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias, Inadequate/lack of blinding of outcome assessors, resulting in potential for detection bias; **Imprecision: serious.** 95% CI include important benefits and harms, Low number of patients;

Summary of findings Table A30.

Population: Patients with P-ACC-related asthenia or fatigue

Intervention: Physical training

Comparator: Standard of care (SOC)

Outcome Timeframe	Study results and measurements	Absolute effect estimates		Certainty of the Evidence (Quality of evidence)	Plain language summary
		SOC	Physical training		
HRQL improvement	Relative risk: 1.44 (CI 95% 0.81 - 2.58) Based on data from 121 participants in 2 studies Follow up 63 days	227 per 1000	636 per 1000 Difference: 168 more per 1000 (CI 95% 112 fewer - 672 more)	Low Due to serious risk of bias, Due to serious imprecision ¹	Physical training may increase HRQL improvement
Functional capacity improvement	Relative risk: 1.53 (CI 95% 0.83 - 2.84) Based on data from 96 participants in 2 studies Follow up 42 days	205 per 1000	314 per 1000 Difference: 109 more per 1000 (CI 95% 35 fewer - 377 more)	Low Due to serious risk of bias, Due to serious imprecision ²	Physical training may increase functional capacity improvement
Fatigue improvement	Relative risk: 4.66 (CI 95% 1.96 - 11.09) Based on data from 96 participants in 2 studies Follow up 35 days	121 per 1000	564 per 1000 Difference: 443 more per 1000 (CI 95% 116 more - 879 more)	Low Due to serious risk of bias, Due to serious imprecision ³	Physical training may increase fatigue improvement
Strength improvement	Relative risk: 3.03 (CI 95% 0.85 - 10.79) Based on data from 73 participants in 1 studies Follow up 56 days	76 per 1000	230 per 1000 Difference: 154 more per 1000 (CI 95% 11 fewer - 744 more)	Very low Due to serious risk of bias, Due to very serious imprecision ⁴	We are uncertain whether cognitive behavioral therapy increases or decreases strength improvement

1. **Risk of Bias: serious.** Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias, Inadequate/lack of blinding of outcome assessors, resulting in potential for detection bias; **Imprecision: serious.** Low number of patients;
2. **Risk of Bias: serious.** Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias, Inadequate/lack of blinding of outcome assessors, resulting in potential for detection bias; **Imprecision: serious.** 95% CI include important benefits and harms, Low number of patients;
3. **Risk of Bias: serious.** Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias, Inadequate/lack of blinding of outcome assessors, resulting in potential for detection bias; **Imprecision: serious.** Low number of patients;
4. **Risk of Bias: serious.** Inadequate/lack of blinding of participants and personnel, resulting in potential for performance bias, Inadequate/lack of blinding of outcome assessors, resulting in potential for detection bias; **Imprecision: very serious.** Low number of patients;

This review compiles the evidence on potential therapeutic options for post COVID-19 condition (PCC). Included are all the identified clinical forms, symptoms and manifestations of PCC for which an intervention was assessed in at least one randomized controlled trial.