

Salt-Smart Americas:

A Guide for
Country-Level Action

Less
than

5g
/day

Salt-Smart Americas:

A Guide for
Country-Level Action



Also published in Spanish (2013) with the title:
La reducción de la sal en las Américas: una guía para la acción en los países
ISBN 978-75-31769-3

PAHO HQ Library Cataloguing-in-Publication Data

Pan American Health Organization.

Salt-Smart Americas: A Guide for Country-Level Action. Washington, DC : PAHO, 2013.

1. Sodium Chloride, Dietary. 2. Cardiovascular Disease. 3. Behavior. 4. Food Industry. 5. Americas. I. Title.

ISBN 978-92-75-11769-9

(NLM Classification: WB 424)

The Pan American Health Organization welcomes requests for permission to reproduce or translate its publications, in part or in full. Applications and inquiries should be addressed to the Department of Knowledge Management and Communications (KMC), Pan American Health Organization, Washington, D.C., U.S.A. (pubrights@paho.org). The Noncommunicable Diseases and Disabilities Unit of the Pan American Health Organization ncdbulletin@paho.org will be glad to provide the latest information on any changes made to the text, plans for new editions, and reprints and translations already available.

© Pan American Health Organization, 2013. All rights reserved.

Publications of the Pan American Health Organization enjoy copyright protection in accordance with the provisions of Protocol 2 of the Universal Copyright Convention. All rights are reserved.

The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of the Secretariat of the Pan American Health Organization concerning the status of any country, territory, city or area or of its authorities, or concerning the delimitation of its frontiers or boundaries.

The mention of specific companies or of certain manufacturers' products does not imply that they are endorsed or recommended by the Pan American Health Organization in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by the Pan American Health Organization to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall the Pan American Health Organization be liable for damages arising from its use.



Table of Contents

Preface	ix
Acknowledgements	x
Recommendations for the Americas	1
<hr/>	
Policy Statement: Preventing Cardiovascular Disease in the Americas by Reducing Dietary Salt Intake Population-Wide	3
<hr/>	
Policy Goal	4
Audience	4
Rationale	4
Recommendations for Policy and Action	4
To National Governments	4
To Nongovernmental Organizations, Health Care Organizations, Associations of Health Professionals	5
To the Food Industry	5
To the Pan American Health Organization	5
Background	6
Prevalence of Hypertension	6
Non-Optimal Blood Pressure, Health and Salt	6
Population-Wide Salt Reduction is Cost-Effective and Equitable	6
Fortification Alternatives	7
References	8
<hr/>	
PAHO/WHO Regional Expert Group for Cardiovascular Disease Prevention Through Population-Wide Dietary Salt Reduction. Final Report November 2011	9
<hr/>	
Executive Summary	10
Key Messages	10
Recommendations	11
Introduction	15
The Consumption and Overconsumption of Salt	15
High Salt Intake Is Unsafe	15
Lowering Salt Intake Is Cost-effective and Cost-saving	16
WHO Platforms for Dietary Salt Reduction	16
WHO Population Salt Reduction Strategy	16
The PAHO Initiative – Cardiovascular Disease Prevention through Population-wide Dietary Salt Reduction	17
Background	17
The Expert Group and its Sub-groups	18
Purpose of this Report	19
Tools, Resources and Achievements	19
Advocacy and Communication	19
Surveillance	20
Food Industry Engagement	21
Synchronizing Salt Iodization and Salt Reduction Programs	23
Economic Studies on the Cost-effectiveness and Cost-savings of Population Level Dietary Salt Reduction	24
Status of Action in the Americas	24
Country-specific Summaries	24
Food Categories for which Salt Content Is Being Reduced	28
Lessons Learned So Far	29
Challenges	30
Momentum and Potential in the Region	31
References	32
Appendix 1: Organizations that Have Endorsed the Policy Statement as of September 2011	34

Appendix 2: Dissemination Activities	36
Appendix 3: Country-specific Targets and Timelines	37

Population Awareness-Building 39

Applying a Social Marketing Framework to Salt Reduction 41

Situation Analysis	42
Literature Review and Research Findings	43
Target Audience	43
Audience Analysis	44
Overview of Supply Side	45
Demand Side/Communication Goals and Objectives	45
Marketing Mix Strategies	46
Monitoring and Evaluation Plan	51
Work Plan and Timeline	52
Budget/Funding Sources	52
Postscript: How Can Social Marketing Strengthen Current Efforts?	53
Appendix 1: Reviews, Outlines and Forms	54
Global Literature Review Synthesis	54
Social Marketing Plan Outline	55
Marketing Mix Planning Tool	57
Creative Brief Form	57
Bravo Criteria Checklist	58
Implementation Plan Template	58

Monitoring and Evaluation of Salt Consumption, Sources of Salt and Knowledge and Behaviors 61

Protocol For Population Level Sodium Determination In 24-Hour Urine Samples 63

Section 1: Introduction	64
Overview of the PAHO/WHO Protocol for Population Level Sodium Determination in 24-Hour Urine Samples	64
Rationale for Population Level Sodium Determination in 24-Hour Urine Samples	64
Section 2: Field Protocol	66
Overview of the Field Protocol	66
Planning and Conducting a 24-Hour Urine Collection Study	66
Selecting the Sample	67
Implementation Plan	69
Applying for Ethical Approval	70
Timeframes and Data Collection Considerations	70
Adapting the PAHO/WHO Protocol for Sodium Determination in 24-Hour Urine Samples	70
Pilot Testing	71
Section 3: Data Collection Guide	71
Instructions for Field Staff, Equipment and Analytic Methods	71
Guide to Physical Measurements	72
Measuring Blood Pressure and Heart Rate	73
Measuring Height	74
Measuring Weight	74
Measuring Waist Circumference	75
Measuring Hip Circumference	75
Section 4: Questionnaire on Knowledge, Attitudes, and Behavior toward Dietary Salt	76

Section 5: Detailed Instructions for Participants in 24-Hour Urine Collection	77
Section 6: Household Salt Collection and Iodine Determination	78
Section 7: Use of Spot Urine to Estimate 24-Hour Excretion of Sodium, Potassium and Iodine	78
Section 8: Dataset for Health Economic Analysis	79
References	79
PAHO/WHO Sodium Detection in 24-Hour Urine Samples Questionnaire	82
A Review of Methods to Determine the Main Sources of Salt in the Diet	91
Section 1: Introduction	92
Section 2: Food Consumption Surveys	92
Overview	92
Methods that Provide Direct or Primary Data on Food Consumption	93
Using Indirect or Secondary Data Sources	95
Attaching Questions or Modules to Existing Chronic Noncommunicable Disease Risk Factor Surveys	96
Section 3: Sodium Content of Foods	97
Overview	97
Methods that Provide Direct or Primary Data	97
Food Composition Tables and Databases from Selected Countries and Regions	97
Using Indirect or Secondary Data Sources	98
Section 4: Salt Added at the Table and during Cooking	99
Qualitative Methods	99
Direct Quantitative Methods	99
Indirect “Subtraction” Method	100
Secondary Sources	100
Section 5: Culturally or Regionally-specific High-sodium Foods	100
Section 6: Recommendations	100
Determining Baseline Data on the Major Contributors to Salt Intake	101
Establishing and Maintaining Food Composition Databases and Tables	101
Deciding on Targets and Interventions	102
Monitoring and Evaluation	102
References	103
Appendix 1: Examples from the Region	104
Protocol for the Survey on Knowledge, Attitudes and Behavior toward Dietary Salt and Health	109
Introduction	110
Justification	110
Main Objective	110
Hypothesis	110
Methodology and Data Analysis	110
References	111
Appendix 1: Questionnaire on Knowledge, Attitudes, and Behavior toward Dietary Salt and Health	112
Optimizing Salt and Iodine Intake	115
White Paper on Improving Public Health by Optimizing Salt and Iodine Intakes	117
Background	118
The Common Goal	119
A Framework for Collaborative Action	119
References	120

Improving Public Health in the Americas by Optimizing Sodium and Iodine Intakes	121
Key Messages	122
Background	122
Presentation Summaries	123
Meeting Outcomes	129
Conclusions and Next Steps	130
References	131
Appendix 1: Meeting Participants and Contributors to the Position Document	133

Targets Setting and Voluntary Industry Engagement 137

A Guide for Setting Targets and Timelines to Reduce the Salt Content of Food	139
Key Messages	140
Introduction	140
Purpose of the Guide	140
Step 1 – Secure the National Strategy to Reduce Dietary Salt	141
Step 2 – Prepare Data	144
Step 3 – Identify the Key Stakeholders	146
Step 4 – Plan the Meetings	148
Step 5 – Monitor Performance	148
Dealing with Technical Issues, Controversies, Barriers	149
Appendix 1: Table 1 - Targets and Timelines for Food Categories in Common in Argentina, Brazil, Canada, Chile and the National Salt Reduction Initiative in the United States (As of January 2013)	151
Appendix 2: Links to National Targets and Timelines Including Industry Commitments Made Public	156
Appendix 3: Pros and Cons for Types of Targets that Can Be Set	157

Preface

The Region of the Americas has taken on the challenge of addressing noncommunicable diseases. These diseases are the leading causes of death in every country of the Region. Hypertension, or high blood pressure, is the primary risk factor for cardiovascular disease (CVD) and greatly contributes to CVD attributable deaths in the Americas.

Salt reduction is recognized as the most cost-effective intervention in the population-based prevention of hypertension. The Pan American Health Organization (PAHO) initiative “Cardiovascular Disease Prevention through Population-wide Dietary Salt Reduction” supports the countries of the Americas by providing effective tools, strategies, and interventions to reduce sodium intake levels among all people in the Region.

High levels of dietary salt consumption remain prevalent in the Americas and greatly exceed the upper international limit of 5g/person/day. Recent estimates of sodium consumption from Argentina (12.0 g of salt/day), Brazil (11.0 g of salt/day), Chile (9.0 g of salt/day), and the United States (8.7 g of salt/day) confirm the unfavorably high salt intakes, which contribute to hypertension and CVD. Sources of dietary sodium vary by country. For example, in the United States and Canada, 75% comes from processed food, while in Brazil, 70% comes from discretionary salt added during cooking or consumption at the table.

In 2009, only three countries reported having national strategies to reduce salt intake at the population level. Recognizing the need to translate knowledge into action, PAHO spearheaded its regional initiative to encourage governments, civil society, and the private sector across the Americas to develop a policy framework that would address the prevention and control of noncommunicable diseases and hypertension through proven and effective salt reduction programs. Four years later, 10 countries stand at the forefront of a growing momentum involving a diverse cross-section of society’s leaders and stakeholders who have partnered together to achieve this goal.

In 2013, a year dedicated to the fight against hypertension and marked by the celebration of World Hypertension Day, PAHO is pleased to launch *Salt-Smart Americas: A Guide for Country-Level Action*. While highlighting the recommendations, protocols, and guidelines developed under the PAHO regional initiative, the publication also underscores the important teamwork of distinguished public health experts, country program leaders, and civil society representatives and their unwavering commitment to galvanize, lead, strengthen, and support action at both the local and national levels.

Despite the Region’s successes, PAHO recognizes that complex issues remain that will require our careful and sustained attention. We will continue to develop and tailor technical expertise that addresses country needs for the creation or expansion of national action, as well as to catalyze and stimulate nongovernmental organizations, civil society, the private sector, and international agencies to meaningfully participate in and contribute to reducing salt overconsumption at the regionwide level.

Carissa F. Etienne
Director

Acknowledgements

The Pan American Health Organization wishes to thank the following partners for their contributions to this publication.

PAHO/WHO Expert Group Members 2009-2011

Norm Campbell | Chair of the Expert Group, University of Calgary, Canada

Omar Dary | Lead for the Iodine Sub-group, A2Z Project, United States

Rainford Wilks | Co-chair for the Caribbean, Tropical Medicine Research Institute, Jamaica

Hasan Hutchinson | Health Canada

Ana Beatriz Pinto de Almeida Vasconcellos | Co-chair for Latin America, September 2009 – March 2011, Ministry of Health, Brazil

Mary L'Abbé | University of Toronto, Canada

Patricia Jaime | Co-chair for Latin America, April 2011 –September 2011, Ministry of Health, Brazil

Darwin Labarthe | Northwestern University Feinberg School of Medicine, United States

Sonia Angell | Centers for Disease Control and Prevention, United States

Hubert Linders | Fundraising Officer and Project Coordinator, Regional Office for Latin America and the Caribbean, Chile

Simón Barquera | National Institute of Public Health, Mexico

Carlos Monteiro | University of São Paulo, Brazil

Adriana Blanco-Metzler | Institute for Research and Education in Nutrition and Health (INCIENSA), Costa Rica

Tito Pizarro | Ministry of Health, Chile

Francesco Cappuccio | Lead for the Surveillance Sub-group, WHO Collaborating Centre for Nutrition, Warwick University, England

Marcello Tavella | Program for the Prevention of Infarct in Argentina (PROPIA), Argentina

Beatriz Champagne | Lead for the Advocacy Sub-group, InterAmerican Heart Foundation, United States

Ricardo Uauy | Lead for the Industry Sub-group, Institute for Nutrition and Food Technology (INTA), Chile

Dan Chisholm | World Health Organization, Switzerland

Lianne Vardy | Canadian Public Health Agency

Ricardo Correa-Rotter | Salvador Zubiran National Institute of Medical Sciences and Nutrition (INNSZ), Mexico

Godfrey Xuereb | World Health Organization, Switzerland

Secretariat

Branka Legetic | Regional Advisor HSD/NCD, PAHO, Washington DC

Barbara Legowski | Consultant, Ottawa, Canada

Norm Campbell | University of Calgary, Canada

PAHO/WHO Technical Advisory Group Members 2012-2015

Norm Campbell | Co-chair of TAG, University of Calgary, Canada

Ricardo Correa-Rotter | Co-chair of TAG, National Institute of Medical Sciences and Nutrition (INNSZ), Mexico

ADVOCACY AND COMMUNICATION

Beatriz Champagne | Co-lead, InterAmerican Heart Foundation, United States

Trevor Hassell | Co-lead, Healthy Caribbean Coalition, Barbados

Hubert Linders | Fundraising Officer and Project Coordinator, Regional Office for Latin America and the Caribbean, Chile

SURVEILLANCE

Sonia Angell | Co-lead, CDC, United States

Mary L'Abbe | Co-lead, University of Toronto, Canada

Adriana Blanco-Metzler | INCIENSA, Costa Rica

INDUSTRY

Ricardo Uauy | Co-lead, University of Chile, INTA, Chile

Daniel Ferrante | Co-lead, Ministry of Health, Argentina

IODINE

Omar Dary | Co-lead, Nutrition and micronutrient expert, United States

Eduardo Augusto Fernandes Nilson | Co-lead, Ministry of Health, Brazil

ECONOMICS

Kirsten Bibbins-Domingo | Lead, University of San Francisco, United States

SCIENTIFIC REVIEW

Lawrence J. Appel | Lead, Johns Hopkins University, United States

Norm Campbell | University of Calgary, Canada

Secretariat

Branka Legetic | Regional Advisor HSD/NCD, PAHO, Washington DC

Barbara Legowski | Consultant, Ottawa, Canada

Norm Campbell | University of Calgary, Canada



**Recommendations
for the Americas**

**Policy Statement:
Preventing Cardiovascular
Disease in the Americas by
Reducing Dietary Salt Intake
Population-Wide**

The PAHO/WHO Regional Expert Group on Cardiovascular Disease Prevention through Dietary Salt Reduction has produced this policy statement. It has the rationale and recommendations for a population-based approach to reduce dietary salt intake among all people in the Americas, be they adults or children.

Policy Goal

A gradual and sustained drop in dietary salt intake to reach national targets or in their absence, the internationally recommended target of less than 5g/day/person by 2020.

Audience

Policy and decision makers in government, leaders in nongovernmental organizations (representing consumers, health, scientific and health care professionals), civil society, the food industry (including food processors and distributors), among food importers and exporters, and in PAHO.

Rationale

- Increased blood pressure world-wide is the leading risk factor for death and the second leading risk for disability by causing heart disease, stroke and kidney failure.
- In the Americas, between 1/5 and 1/3 of all adults has hypertension and once age 80 is reached, over 90% can be expected to be hypertensive.
- In 2001, the management of non-optimal blood pressure i.e. systolic pressure over 115 mmHg consumed about 10% of the world's overall healthcare expenditures.
- As dietary salt consumption increases, so does blood pressure. Typical modern diets provide excessive amounts of salt, from early childhood through adulthood.
- The recommended intake of salt is less than 5g/day/person. In the Americas, intake can be over double the recommended level. All age groups including children are affected.
- Adding salt at the table is not the only problem. In most populations by far the the largest amount of dietary salt comes from ready-made meals and pre-prepared foods, including bread, processed meats, and even breakfast cereals.
- Reducing salt consumption population-wide is one of the most cost-effective measures available to public health. It can lower the rates of a number of related chronic diseases and conditions at an estimated cost of between 4 (US) cents and 32 (US) cents per person per year. Population-wide interventions can also distribute the benefits of healthy blood pressure equitably.

- Governments are justified in intervening directly to reduce population-wide salt consumption because salt additives in food are so common. People are unaware of how much salt they are eating in different foods and of the adverse effects on their health. Children are especially vulnerable.
- Salt intake can be reduced without compromising micronutrient fortification efforts.

Recommendations for Policy and Action

The recommendations below are consistent with the World Health Organization's three pillars for successful dietary salt reduction: product reformulation; consumer awareness and education campaigns; and environmental changes to make healthy choices the easiest and most affordable options for all people.

To National Governments

- Seek endorsement of this policy statement by ministries of health, agriculture and trade, by food regulatory agencies, national public health leaders, nongovernmental organizations (NGOs), academia, and relevant food industries.
- Develop sustainable, funded, scientifically based salt reduction programs that are integrated into existing food, nutrition, health and education programs. The programs should be socially inclusive and include major socioeconomic, racial, cultural, gender and age subgroups and specifically children. Components should include:
 - Standardized food labeling such that consumers can easily identify high and low salt foods.
 - Educating people including children about the health risks of high dietary salt and how to reduce salt intake as part of a healthy diet.
- Initiate collaboration with relevant domestic food industries to set gradually decreasing targets, with timelines, for salt levels according to food categories, by regulation or through economic incentives or disincentives with government oversight.
- Regulate or otherwise encourage domestic and multi-

national food enterprises to adopt the lowest of a) best in class (salt content to match the lowest in the specific food category) and b) best in world for the national market (match the lowest salt content of the specific food produced by the company elsewhere in the world).

- Develop a national surveillance system with regular reporting to identify dietary salt intake levels and the major sources of dietary salt. Monitor progress towards the national target(s) for dietary salt intake or the internationally recommended target.
- Review national salt fortification policies and recommendations to be in concordance with the recommended salt intake.
- Extend official support to the Codex Alimentarius committee on food labeling for salt/sodium to be included as a mandatory component of nutrition labels.
- Develop legislative or regulatory frameworks to implement the World Health Organization (WHO) recommendations on advertising of food products and beverages to children.

To Nongovernmental Organizations, Health Care Organizations, Associations of Health Professionals

- Endorse this policy statement.
- Educate memberships on the health risks of high dietary salt and how to reduce salt intake. Encourage involvement in advocacy. Monitor and promote presentations on dietary salt at national meetings and the publication of articles on dietary salt.
- Promote and advocate media releases on dietary salt reduction to reach the public, including children and particularly women given their integral roles in family health and food preparation.
- Broadly disseminate relevant literature.
- Educate policy and decision makers on the health benefits of lowering blood pressure among normotensive and hypertensive people, regardless of age.
- Advocate policies and regulations that will contribute to population-wide reductions in dietary salt.
- Promote coalition building, increase organizational capacity for advocacy and develop advocacy tools to promote civil society actions.

To the Food Industry

- Endorse this policy statement.
- Make current best in class and best in world low salt products and practices universal across global markets as soon as possible. Make salt substitutes readily available at affordable prices.

- Institute reformulation schedules for a gradual and sustained reduction in the salt content of all existing salt-containing food products, restaurant and ready-made meals to contribute to achieving the internationally recommended target or national targets where applicable. Make all new food product formulations inherently low in salt.
- Use standardized, clear and easy-to-understand food labels that include information on salt content.
- Promote the health benefits of low salt diets to all peoples of the Americas.

To the Pan American Health Organization

- Ensure good communications and information sharing between regional and international initiatives to foster best practices.
- Develop a template for national report cards and report to Member States on comparative national baselines and progress at pre specified time points (e.g. in 2010 the baseline, progress in 2015 and 2020).
- Work with Member States to monitor dietary salt consumption in the Americas.
- Develop and foster a network of endorsing governments, NGOs, and expert champions on dietary salt in the Pan American region.
- Develop a web based 'toolbox' with educational materials and programs on dietary salt for the public, patients, health care professionals that are culturally appropriate to sub-regions of the Americas.
- Develop and advocate conflict of interest guidelines to assist health organizations and scientists in the Pan American region in their interactions with the food industry.
- Foster research on the economic and health impacts of high dietary salt in the countries and sub-regions of the Pan American region.
- Assist Member States to revise national and sub-regional fortification programs to be consistent with efforts to reduce dietary salt.
- Collaborate with the Food and Agriculture Organization (FAO), UNICEF, the Codex Alimentarius Commission and other relevant UN bodies to achieve a consistent and coordinated approach to reducing dietary salt.
- Educate policy and decision makers on the health benefits of lowering blood pressure among normotensive and hypertensive people, regardless of age.
- Advocate policies and regulations that will contribute to population-wide reductions in dietary salt.

Background

Prevalence of Hypertension

About one in four adults worldwide had hypertension in 2000 (1). As populations age, rates of hypertension will increase. The Framingham study found that 90% of normotensive people aged 55 to 65 will develop high blood pressure if they reach average life expectancy (2). By 2025, without intervention, 29% of adults around the world are expected to have hypertension (1).

In Canada, one in five adults has hypertension (3) and in the United States, 29% of adults were estimated to be hypertensive in 2003-04 (4). In the different countries of Latin America, the prevalence of hypertension ranges from 26 to 42% of the general adult population (5).

Non-Optimal Blood Pressure, Health and Salt

WHO states that increasing blood pressure worldwide is the leading risk factor for death (6) and the second leading risk for disability by causing heart disease, stroke and kidney failure (7). Whereas most health care professionals consider systolic blood pressure at 140 mmHg and over to be “hypertension”, the relative risk for cardiovascular diseases (CVD) begins to rise when blood pressure goes above 115 mmHg. Thus a much wider range of non-optimal blood pressure is adversely affecting health, and has been attributed to most CVD deaths from ischemic heart disease and stroke (8).

There is strong evidence that salt added to food is a major factor increasing the blood pressure in normotensive and hypertensive people, whether adults or children. A high salt diet also increases the risk of left ventricular hypertrophy and kidney damage, is a probable cause of gastric cancer, and has possible associations with osteoporosis, calcium containing renal stones and increased severity of asthma. Because salty foods cause thirst they are likely an important contributor to obesity, especially among children and adolescents, through association with increased consumption of high-calorie soft drinks (9,10,11).

A technical report for the WHO and FAO recommends salt intake of less than 5g/day/person, the target for a healthy diet, equivalent to 2000 mg of sodium (12). Among the countries in the Americas where standardized and comparable sodium excretion was studied, salt intake was found to be as high as 11.5g/day/person (13). Data for the United States for 2005-06 show average daily intake of sodium among people aged 2 years and over to be 1.5 times the recommended

upper limit (UL) (14). In Canada, over 85% of men and 60% of women between 19 and 70 years of age have salt intake exceeding the UL. Over 90% of Canadian children aged 4 to 8, and 83% of girls and 97% of boys aged 9 to 13 ingest more than the recommended maximum. The situation is the same in almost 80% of Canadian children between ages 1 and 3 (15).

Population-Wide Salt Reduction Is Cost-Effective and Equitable

In 2001, the management of nonoptimal blood pressure and its resulting diseases consumed about 10% of global healthcare expenditures, considered a conservative estimate. If the welfare losses due to premature death are added, the costs could be 20 times higher (8). Effectively lowering blood pressure on a universal scale requires actions with population-wide reach. Individual advice and instruction, part of any comprehensive approach to healthy blood pressure, have a limited impact. On the other hand, reducing salt in the diet of whole populations, not only what is used at the table but more importantly what is added to processed and ready-made foods like bread, processed meats and breakfast cereals, can distribute the benefits of lowered blood pressure broadly and equitably (16,17).

Governments are justified in taking a population based approach to reduce salt intake because salt additives in food are so common. People are unaware of how much salt they are eating in different foods and of the adverse effects on their health. Children are especially vulnerable.

Lowering blood pressure through population-wide salt intake reduction is cost effective (17,18). A strategy that combines mass-media awareness campaigns with regulation of the salt content of food products has been estimated to cost between \$0.04 and \$0.32 US per person per year. Over 10 years, the strategy is predicted to avert 8.5 million deaths world-wide, mostly from CVD (17).

The savings to healthcare budgets can be dramatic. Researchers in the UK estimate that achieving dietary salt intake of less than 6g/day could potentially reduce the need for anti-hypertensive drugs by as much as 30% (19). Already, a 10% reduction in salt intake in the UK since 2000-01, attributed to the combined gradual and sustained efforts of industry lowering the salt in certain food products and to the Food Standards Agency's information campaign, has yielded an annual cost saving benefit of £1.5 billion (20).

In the US, if average population intake fell to 5g/day, there could be 11 million fewer cases of hypertension, saving approximately \$18 billion in healthcare and

gaining about \$32 billion in quality adjusted life years (21). In Canada, reducing salt food additives is estimated to decrease hypertension prevalence by 30% and almost double the rate of successful treatment and control. Direct savings to the health system just from reduced hypertension management costs were estimated at \$430 million/year (22).

Fortification Alternatives

Salt is used in some areas of the Americas as a vehicle for iodine and similarly in some cases to fortify fluoride intake. Alternative vehicles for fortification exist, such as vegetable oils and milk. Changes in practice need to be coordinated with policies to reduce dietary salt.

Acknowledgements

Norm Campbell, Chair

(University of Calgary – Canada)

Rainford Wilks, Co-chair for the Caribbean

(Tropical Medicine Research Institute – Jamaica)

Ana Beatriz Pinto de Almeida Vasconcellos Co-chair for Latin America

(Ministry of Health – Brazil)

Simón Barquera

(National Institute of Public Health – Mexico)

Adriana Blanco-Metzler

(Institute for Research and Education in Nutrition and Health (INCIENSA) – Costa Rica)

Ezzedine Boutrif

(FAO – Italy)

Francesco Cappuccio

(Warwick University – England)

Beatriz Champagne

(InterAmerican Heart Foundation – United States)

Ricardo Correa-Rotter

(Salvador Zubiran National Institute of Medical Sciences and Nutrition (INNSZ) – Mexico)

Omar Dary

(A2Z Project – United States)

Darwin Labarthe

(CDC – United States)

Mary L'Abbe

(University of Toronto – Canada)

Hubert Linders

(Fundraising Officer and Project Coordinator, Regional Office for Latin America and the Caribbean - Chile)

Carlos Monteiro

(University of São Paulo – Brazil)

Tito Pizarro

(Ministry of Health – Chile)

Jorge Polônia

(University of Fernando Pessoa – Portugal)

Marcelo Tavella

(Program for the Prevention of Infarct in Argentina (PROPIA) – Argentina)

Ricardo Uauy

(Institute for Nutrition and Food Technology (INTA) – Chile)

Lianne Vardy

(Canadian Public Health Agency)

Sonia Angell

(NYC Health, New York, United States)

References

- 1 Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, He J.** Global burden of hypertension: analysis of worldwide data. *Lancet*. 2005;365:217-23.
- 2 Vasan RS, Beiser A, Seshadri S, Larson MG, Kannel WB, D'Agostino RB, Levy D.** Residual lifetime risk for developing hypertension in middle-aged women and men: The Framingham Heart Study. *JAMA*. 2002;287:1003-10.
- 3 Joffres MR, Ghadirian P, Fodor JG, Petrasovits A, Chockalingam A, Hamet P.** Awareness, treatment and control of hypertension in Canada. *Am J Hypertens*. 1997;10:1097-1102.
- 4 Ong KL, Cheung BMY, Man YB, Lau CP, Lam KSL.** Prevalence, awareness, treatment and control of hypertension among United States adults 1999–2004. *Hypertension*. 2007;49:69-75.
- 5 Sanchez RA, Ayala M, Baglivo H, Velazquez C, Burlando G, Kohlmann O, Jimenez J, Jaramillo PL, Brando A, Valdes G, Alcocer L, Bendersky M, Ramirez AJ, Zanchetti A;** Latin America Expert Group. Latin American guidelines on hypertension. *J Hypertens*. 2009;27:905-22.
- 6 World Health Organization.** The World Health Report 2002: Reducing risks, promoting healthy life. Available at: <http://www.who.int/whr/2002/en/> Accessed September 13, 2009.
- 7 Hsu C, McCulloch CE, Darbinian J, Go AS, Iribarren C.** Elevated blood pressure and risk of end-stage renal disease in subjects without baseline kidney disease. *Arch Intern Med*. 2005;165:923-28.
- 8 Gaziano TA, Bitton A, Anand S, Weinstein MC** for the International Society of Hypertension. The global cost of non-optimal blood pressure. *J Hypertens*. 2009; 27:1472-77.
- 9 He FJ, MacGregor GA.** A comprehensive review on salt and health and current experience of worldwide salt reduction programmes. *J Hum Hypertens*. 2009;23: 363-84.
- 10 Feng J, MacGregor GA.** Importance of salt in determining blood pressure in children: Meta-analysis of controlled trials. *Hypertension*. 2006;48:861-69.
- 11 Mohan S, Campbell NRC, Willis K.** Effective population-wide public health interventions to promote sodium reduction. *CMAJ*. 2009;DOI:10.1503/cmaj.090361. Available at: <http://www.cmaj.ca/cgi/rapidpdf/cmaj.090361v1.pdf>. Accessed October 20, 2009.
- 12 World Health Organization.** Reducing salt intake in populations: Report of a WHO forum and technical meeting, 5-7 October 2006, Paris, France. Available at: http://www.who.int/dietphysicalactivity/Salt_Report_VC_april07.pdf. Accessed September 13, 2009.
- 13 Intersalt Comparative Research Group.** Intersalt: an international study of electrolyte excretion and blood pressure. Results for 24 hour urinary sodium and potassium excretion. *BMJ*. 1988;297:319-28.
- 14 United States Centers for Disease Control and Prevention.** Application of lower sodium intake recommendations to adults - United States, 1999-2006. *MMWR Morb Mortal Wkly Rep*. 2009;58:281-3.
- 15 Garriguet D.** Sodium consumption at all ages. *Statistics Canada Health Reports*. 2007;18:47-58. Available at: <http://www.statcan.gc.ca/pub/82-003-x/2006004/article/sodium/9608-eng.pdf>. Accessed September 7, 2009.
- 16 Feng JH, MacGregor GA.** Salt in food. *Lancet*. 2005; 365: 844-45.
- 17 Asaria P, Chisholm D, Mathers C, Ezzati M, Beaglehole R.** Chronic disease prevention: health effects and financial costs of strategies to reduce salt intake and control tobacco use. *Lancet*. 2007;370:2044–53.
- 18 Murray CJ, Lauer JA, Hutubessy RC, Niessen L, Tomijima N, Rodgers A, Lawes CM, Evans DB.** Effectiveness and costs of interventions to lower systolic blood pressure and cholesterol: a global and regional analysis on reduction of cardiovascular disease risk. *Lancet*. 2003; 361: 717-25.
- 19 Walker J, MacKenzie AD, Dunning J.** Does reducing your salt intake make you live longer? *Interact Cardiovasc Thorac Surg*. 2007;6:793-98.
- 20 Food Standards Agency (UK).** Agency publishes 2012 salt reduction targets. Monday 18 May 2009. Available at: <http://www.food.gov.uk/news/pressreleases/2009/may/saltpreleases>. Accessed September 22, 2009.
- 21 Palar K, Sturm R.** Potential societal savings from reduced sodium consumption in the US adult population. *Am J Health Promotion*. 2009;24:49-57.
- 22 Joffres MR, Campbell NR, Manns B, Tu K.** Estimate of the benefits of a population-based reduction in dietary sodium additives on hypertension and its related health care costs in Canada. *Can J Cardiol*. 2007;23:437-43.

**PAHO/WHO Regional Expert
Group For Cardiovascular
Disease Prevention
Through Population-Wide
Dietary Salt Reduction**

Final Report
November 2011

Executive Summary

High salt* diets are a major cause of raised blood pressure, increasing the risk for cardiovascular and kidney disease and death. Already in 2000, about one in four adults world-wide had hypertension. Prevalence is expected to rise from a combination of people generally living longer and prevention of raised blood pressure being largely ineffective up to now.

Reducing dietary salt at the population level is the most cost-effective public health measure available to lower blood pressure and mortality. It can save lives and healthcare dollars across low, middle and high-income countries. Knowing this, the Pan American Health Organization (PAHO) in September 2009 convened a group of independent international experts on salt and health to guide the first two-years of a regional initiative – Cardiovascular Disease Prevention through Dietary Salt Reduction.

The Expert Group and its five sub-groups, supported by additional technical expertise, addressed issues and prepared resources to assist Member States in engaging the food industry on product reformulation; advocacy and communication; surveillance (sources of salt in the diet, salt intake and public knowledge and opinions on salt and health); salt fortification with iodine; analysis of health and economic impacts of dietary salt reduction; and review of scientific evidence. The work is consistent with and supportive of the concurrent WHO global initiative for dietary salt reduction.

This report marks the completion of the first two years of the regional initiative and the 24-month mandate of the Expert Group. The report summarizes the progress made and the lessons learned to date by the stakeholders, and it features the achievements of and resources prepared by the Expert Group and its sub-groups. It also presents the Expert Group's key messages and recommendations directed to the major stakeholders in dietary salt reduction: Member States; the food and salt industries; non-government organizations (NGOs) and civil society; PAHO/WHO and other international organizations.

There is momentum and potential for a second phase

*The word salt is used to refer to sodium and the term “reducing dietary salt intake” implies the reduction of total sodium intake from all dietary sources including, for example, salt added during food manufacturing and processing, sodium additives such as monosodium glutamate and other sodium-based preservatives or taste enhancers, as well as salt added at the table and in cooking. Where salt substitutes are advocated (partially replacing sodium with potassium and other components) sodium reduction may be achieved without salt reduction thereby requiring the distinction between the two terms.

of activity in the region. Several of the challenges identified by countries, whether active or not yet, can be addressed through the dissemination and facilitated implementation of the tools and resources collected and developed by the Expert Group. The target and timeline commitments made by the multinational and large national food companies in Brazil and to the US National Salt Reduction Initiative (NSRI) (which in addition received restaurant chain commitments) are evidence of what can be anticipated with regards to food reformulation and can be examples to other markets and countries in the region. The occasions for dissemination are also opportunities to continue and expand networking and sharing of plans, resources and approaches. For low- and middle-income countries that have not yet launched dietary salt reduction initiatives and are considering where to start, the experiences in the southern cone countries, especially regarding bread – a major source of salt – provide important lessons and direction. The national experiences with and tools for public information and awareness raising campaigns are potentially of great value to these countries – they can inform public health policy makers who need to build the broad public support that can secure the issue on political agendas.

Key Messages

Salt intake exceeding biologically adequate levels has a causal and direct relationship with greater-than-optimum levels of blood pressure.

Countries should launch national initiatives to reduce the overconsumption of salt as part of non-communicable disease prevention or healthy nutrition policies, taking advantage of the lessons learned and momentum in other countries and the region, using entry points and action sequences most appropriate to country contexts, adopting at least the internationally recommended target of less than 2000 mg sodium or 5 g salt from all sources per person per day by 2020.

Countries should use the most reliable and valid methods feasible to determine a baseline of the main sources of salt in the national diet (salt in commercially processed foods and restaurant/catered foods that is added before products are sold, and the salt or high salt products added at the table and in cooking [personal discretionary use]) and population level salt intake. While secondary data can supply estimates of total salt intake, there should be provision made for 24-hour urine sampling to confirm findings.

Countries should take advantage of health economic analysis models and methods currently available to

demonstrate the disease burden of high salt diets and conversely the benefits to population health and economic development of reducing the overconsumption of salt.

Where packaged foods and food service establishments (restaurants and caterers) are a major source of dietary salt or are emerging as such, countries working with the food industry should set targets and timelines for reducing salt content. When doing so, countries should review the reformulation commitments and low salt products made by food companies in and for other markets in the region and elsewhere as evidence of what can be done. Countries can begin with voluntary targets for high volume food categories followed by targets for all foods that contribute salt to the diet. Should voluntary efforts lag or fail, the salt content of foods should be regulated within the national food safety and health protection legislative frameworks to ensure market-wide impact and coverage of all relevant food categories.

Countries should include broad public awareness and education campaigns in their national initiatives targeting high salt intake. The campaigns should mobilize relevant non-government organizations, consumer organizations, civil society, public policy makers, public health and healthcare professionals and the food and salt industries and their trade associations, and be guided by national knowledge on population level salt intake and the main sources of salt in the diet.

Countries should establish systems to objectively and transparently monitor food industry progress against the targets and timelines and evaluate and publicly disclose whether or not industry commitments for salt reduction, whether voluntary or mandatory, are being met. At the same time countries should continue to monitor population salt intake levels and sources of dietary salt through risk factor studies to determine the impacts of food product reformulations and reductions in the personal use of salt.

Countries should routinely monitor population level iodine intake as dietary salt is reduced and, where iodine deficiency is a concern, develop policies for the voluntary or mandatory use of iodized salt or iodine-containing premixes in commercially produced food at levels appropriate to population iodine needs.

Recommendations

To Member States

Recognizing that people in the Americas are over consuming salt, resulting in raised blood pressure and increased risk of cardiovascular and kidney disease, and understanding that unhealthy diet is a major threat to

the health of populations and to sustained economic growth and productive workforces, the governments of Member States should take a leadership role and:

Reduce exposure

Launch national multi-stakeholder initiatives to reduce people's exposure to high salt diets as part of national non-communicable disease prevention or nutrition policies. Taking guidance from the *WHO Global Strategy on Diet, Physical Activity and Health*, use entry points and action sequences most appropriate to country contexts, using processes where collaboration with stakeholders is transparent and conflicts of interest are minimized.

Strengthen surveillance

Develop or adjust and sustain surveillance mechanisms, using the most reliable and valid methods feasible in country contexts, to first determine a baseline and then monitor the main sources of salt in the diet (salt in commercially processed foods and in foods prepared by food service establishments (restaurants/caterers) where the salt is added before products are sold, and the salt added at the table and in cooking [personal discretionary use]) and population level salt intake. While secondary sources e.g. salt industry sales per capita, can supply estimates of salt intake, there should be provision made for 24-hour urine sampling to confirm findings of total salt intake. Refer to the tools and guidance prepared by PAHO/WHO specific for surveillance of dietary salt.

Adopt timelines

Establish a schedule of national targets and timelines for a salt reduction program based on the knowledge of salt intake levels and the main sources of salt in the national diet, setting the ultimate target as at least the internationally recommended intake of less than 2000 mg sodium or 5 g salt from all sources per person per day by 2020. Establish interim targets e.g. at 3 and 5 year points as a means to monitor progress and allow adjustment of targets as needed.

Address economics

Estimate the health and economic development benefits of whole populations achieving the optimal level of salt intake to make the case for concerted all-of-society action. Take advantage of currently available models and methods.

Set targets

Engage multinational and national food industries – food manufacturers and food service establishments

– in meaningful dialogues to establish sustainable and transparent processes where the industries commit to follow schedules to meet gradually lowered and measurable targets for the salt content of food categories, taking into account, as evidence of what can be done, the low salt products, reformulation schedules, targets and technologies applied by food companies in other countries where national salt reduction programs are underway. Closely monitor and disclose publicly the progress against commitments.

Consider regulation

Consider regulating the salt content of foods within the national food safety and health protection legislative frameworks to ensure market-wide impact and coverage of all relevant food categories.

Institute advocacy

Build or improve the capacities of the public institutions responsible for developing and maintaining a national initiative, specific to national needs, and demonstrate to policy makers at all levels and in various sectors the benefits and potential contributions of population level dietary salt reduction.

Clarify labeling

Mandate clear and simple labeling of packaged foods and foods prepared by food service establishments for the purchaser to easily identify the amount of salt contained in each serving (or per reference amount used in the national labeling system) and easily understand if the product is high versus low in salt content. At the same time, discourage the marketing of processed foods as “healthy” solely on the basis of being sodium-reduced given that unhealthy properties may remain e.g. high density, high fat, high sugar.

Require reporting

Require food manufacturers and food service chains to annually disclose and report the nutritional content of their products including their sales volumes and use this information to track trends in the salt content of the products and their consumption. Establish the mechanisms e.g. databases necessary to monitor trends. Databases should be publically accessible for use by consumers and researchers and civil society.

Communicate effectively

Coordinate and collaborate with, and where necessary train non-government organizations, consumer organizations and civil society to assist in broad and multifaceted educational campaigns to ensure that con-

sumers, public health and healthcare professionals, public policy makers and the food and salt industries are informed and educated about the harmful effects of excess salt on health, about the main sources of salt in their diets and the amounts in food products.

Protect children

Refer to the 2010 WHO *Set of Recommendations on the Marketing of Foods and Non-alcoholic Beverages to Children* to guide the design of or strengthen existing national policies to reduce children’s exposure to the marketing of foods high in saturated fats, trans fatty acids, added sugars or salt.

Standardize procurement

Develop standardized food procurement policies for all public institutions and for the use of public funds consistent with healthy nutrition guidelines. Policies should include criteria limiting the amount of salt in products purchased.

Enlist workplaces

Encourage workplace wellness programs in public and private sectors that include access to healthy food choices including those that are low in salt.

Monitor iodine

Monitor iodine intake at the population level in order to adjust iodization levels of salt, both discretionary salt (added at the table and in cooking) as well as that used by the food industry where applicable. Where iodine deficiency disorders remain a problem, pursue or strengthen policies for the voluntary or mandatory use of iodized salt or iodine-containing premixes in commercially produced food at levels appropriate to population iodine needs. Engage with PAHO and other international agencies to develop an overarching comprehensive action framework for countries in the region that complements existing regional and national efforts for dietary salt reduction.

Disseminate successes

Use the sub-regional economic platforms to facilitate dissemination of best practices that reduce the

* CARMEN is an acronym for Collaborative Action for Risk Factor Prevention and Effective Management of Non-communicable Diseases and Conjunto de Acciones para la Reducción Multifactorial de Enfermedades No Transmisibles – the network of countries in the Caribbean and Latin America that are applying integrated community-based approaches to preventing and managing non-communicable diseases and their risk factors.

salt content of the food supply for food safety and food quality purposes and to reduce salt in foods imported and exported across common markets.

Utilize CARMEN

Through the CARMEN network* promote the advances of national programs and disseminate lessons learned.

To the Food and Salt Industries

Recognizing that progress has been made in certain markets with new and reformulated low/no salt products, all multinational and national food manufacturers, private label retailers, food service establishments (restaurants and caterers) and the food services sector should:

Be transparent

Make public the plans to reduce salt in all products, prioritizing mainstream and biggest sellers, and monitor progress on the plans and make public the results.

Set targets

Cooperate with national governments in setting and making public reformulation targets and timelines that will meaningfully contribute to reaching the ultimate national intake target and implement corresponding product development and reformulation schedules.

Supply globally

Supply to all countries in the Americas now, and continue to supply as they become available, the products with the best nutrient composition including the lowest salt content in a company's product line available anywhere in the world.

Transfer technologies

Transfer the technologies that have delivered low/no salt content products in such countries as the United Kingdom and the United States to all branches and/or subsidiaries in countries in the Americas including the products with internationally recognized brands.

Communicate responsibly

Ensure that public messages related to salt and health used in advertising, marketing and promotional activity is consistent with and reinforce those of WHO and the PAHO Policy Statement. Inform consumers responsibly that taste preferences for salt can be changed.

Clarify labelling

Ensure clear and simple labelling of packaged foods

and foods prepared by food service establishment for the purchaser to easily identify the amount of salt contained in each serving (or per reference amount used in the national labelling system) and easily understand if the product is high versus low in salt content.

Supply information

Regularly provide up-to-date nutrition information to government in the format specified by government for the purposes of monitoring the salt content of foods.

Protect children

Act according to the WHO Set of Recommendations on the Marketing of Foods and Non-alcoholic Beverages to Children.

Honor commitments

To members of the International Food and Beverage Alliance who pledged to WHO in 2008 to support its Global Strategy on Diet, Physical Activity and Health, act meaningfully on the commitments made with regards to reformulation and new food product development to offer consumers low/no salt alternatives world-wide.

Recognizing that the salt in packaged foods and foods prepared by food service establishments are increasingly if not already replacing discretionary salt as the main source of salt in the diet, and that these processed foods should therefore contain appropriate amounts of iodized salt or include iodine in fortification premixes, food manufacturers should:

Conduct research

Invest in research to determine how to efficiently manufacture foods to deliver iodine at levels appropriate to population needs.

To Nongovernment Organizations and Civil Society

Recognizing that NGOs and civil society, including also researchers and the academic community, have shown themselves to be powerful advocates and activists, their work can be parallel to or stimulate that of governments to move public health agendas. They can also motivate private sector responses to public issues or demands. Where not already mobilized on the issue of overconsumption of salt, NGOs, consumer associations and civil society should:

Build capacities

Build internal capacity to advance salt reduction

efforts. Societies of healthcare professionals and organizations representing consumers or patients should mobilize their memberships and build coalitions to impress upon individuals the need to reduce the personal use of salt and to pressure food manufacturers and food service establishments to reduce the salt content of their products.

Standardize messages

Use standardized educational messages to promote a consistent understanding of the health issues related to high salt diets and solutions. Emphasize public education where discretionary salt – added at the table and in cooking – is the major source of salt in the diet. Where commercially prepared and restaurant/catered foods are or are becoming the main sources of salt, raise public awareness and mobilize consumers to demand greater choice and control over the salt content in the foods they buy. Use the fact sheets and other materials made available by PAHO/WHO, designed to facilitate common messaging to make the case.

Advocate transparency

Advocate processes where public health collaboration with stakeholders is transparent and conflicts of interest are minimized.

Monitor industry

Be the “watch dog” to monitor food industry adherence to voluntary or mandatory reformulation schedules or the government oversight of targets, and keep consumers informed of progress or lack there-of.

Advocate healthy procurement

Advocate standardized healthy food procurement policies for public and private sectors that include placing limits on the salt content of foods purchased and served.

Advocate labelling

Advocate clear and simple labelling of packaged foods and foods in food service establishments for the purchaser to easily identify the amount of salt contained in each serving (or per reference amount used in the national labelling system) and easily understand if the product is high versus low in salt content.

Engage governments

Engage with and strengthen relationships with governments using existing communication channels and broadening them when the opportunities arise.

Engage media

Engage with the media as broadly as possible to disseminate sound scientific evidence and advances in salt reduction in support of campaigns and advocacy for public and private sector actions. Put forward champions or prominent public figures as advocates to strengthen the actions to reduce population level salt consumption and initiate/support consumer demand for salt reduction across the food supply.

Protect children

Facilitate and support efforts to implement the WHO Set of Recommendations on the Marketing of Foods and Non-alcoholic Beverages to Children.

Engage internationally

Engage with PAHO and other international agencies to develop an overarching comprehensive action framework for the region to leverage existing regional and national NGO and civil society efforts for dietary salt reduction.

To Pan American Health Organization

Recognizing its primary role in facilitating dialogue, providing technical support to Member States and intervening with national and international agencies on behalf of Member States, PAHO should:

Advocate all-of-society approaches

Advocate among governments and all stakeholders the need for a population-based all-of-society approach to salt reduction as a proven cost-effective and cost-saving intervention.

Build databases

Develop, maintain and make publicly available a database of national policies with a dietary salt reduction component including descriptions of mechanisms and instruments being applied and the national targets and timelines set for food category reductions.

Facilitate dialogues

Facilitate meaningful dialogue among national governments, the food and salt industries, NGOs and civil society to develop an overarching comprehensive and multisector action framework for the region that complements existing regional and national efforts for dietary salt reduction and salt iodization programs.

Utilize CARMEN

Ensure that learning and best practices in countries that are lowering dietary salt are shared through the

CARMEN network and continue to provide technical and expert support to countries as needed.

Encourage networks

Encourage development of other networks and more generally the linking of similar initiatives to optimize the exchange of specific experiences and knowledge.

Facilitate analyses

Facilitate the national assessments of the population health and economic development advantages of lowering dietary salt through regional or sub-regional workshops on health economics modelling.

Stimulate research

Stimulate country-based and regional collaboration with research communities to evaluate and publish relevant findings, especially the evidence of cost-effectiveness and cost-savings from lower dietary salt, along with the initiatives and innovations implemented in low- and middle-income countries as models appropriate to the economies and cultures of the Americas.

Protect children

Facilitate countries to implement the WHO Set of Recommendations on the Marketing of Foods and Non-alcoholic Beverages to Children.

Facilitate synchronization

Promote and support the synchronization of national salt iodization with salt reduction programs by facilitating collaboration with international stakeholders and with them jointly support country-level pilot studies.

Advocate reporting

Engage the relevant international and national agencies to advocate transparent and publicly available food composition data and database development.

Introduction

The Consumption and Overconsumption of Salt

The human body requires sodium to regulate body fluids and maintain critical body functions. Over the millennia of human evolution, the small amount of sodium found naturally in foods was physiologically sufficient by virtue of the body having developed mechanisms to retain and conserve it (1).

It has been approximately 5000 years since salt was first added to food to preserve it and enhance taste and thus added salt became the major source of dietary sodium. But whereas salt was originally a precious commodity


with cultural and even religious significance and hence carefully used, increased mining and ease of transportation over the centuries made it gradually more accessible and inexpensive. With the developments in industrialized food processing over the last few hundred years, salt has become an almost ubiquitous additive to food with additional functions in e.g. food texture and appearance. Consumption is now so reliable that table salt is used as the principle vehicle for supplementary micronutrients e.g. iodine to prevent iodine deficiency disorders. Today in modern diets, naturally occurring sodium found in unprocessed foods accounts for less than 12 % of total sodium intake; the balance of sodium intake is attributed to added salt (2). In high income economies, salt is particularly concentrated in commercially manufactured food products (1-3).

The internationally recommended salt intake level is less than 5 g/person/day (4-5). Actual consumption levels are with few exceptions high world-wide, proven for the first time by the INTERSALT Study (1985-87) that determined comparable standardized sodium intake levels from 52 population samples in 32 countries based on 24-hour urine collections. The subsequent INTERMAP Study from 1996-99 in four countries provided similar findings and several recent dietary and urinary sodium data from observational and interventional studies continue to show excess intake (6).

In the Americas, recent dietary and urinary sodium excretion estimates confirm high per capita salt intake: Brazil at 11 g of salt/day (7); Argentina with 12 g of salt/day (8); Chile at 9.8 g of salt/day (9); the United States with 8.7 g of salt/day (10),] and Canada with 7.7 g/day (11). Sources of dietary sodium vary: in the United States (10) and Canada (11), 75% and 77% respectively of sodium consumed comes from processed foods; in parts of Brazil, 70% is attributed to discretionary salt (added in household cooking or at the table) (7).

High Salt Intake Is Unsafe

Salt intake exceeding biologically adequate levels has a causal and direct relationship with greater-than-optimum levels of blood pressure (12-13). Excess salt consumption increases the blood pressure of infants and children, of normotensive and hypertensive adults (14-17) and is attributable to 30% of the prevalence of hypertension (18-19) found in one in four of the world's adult population. Increased blood pressure even within the normal range is a major cause of disability and is the leading risk for premature death in the world (14). Based on it increasing blood pressure and its probable promotion of gastric cancer, high salt intake is estimated to be the 7th leading risk for premature death in the United States and the sec-



ond leading risk in Chile (20-21). High salt intake also causes kidney stones and has a strong pathophysiological basis and association with osteoporosis, increased asthma severity, and obesity (22-24). The prevalence of elevated blood pressure is expected to rise from a combination of people generally living longer and prevention of rising blood pressure being largely ineffective up to now, making the condition almost inevitable with advancing age. In 2001, non-optimal blood pressure and its resulting diseases consumed about 10% of global healthcare expenditures, considered a conservative estimate (25).

Lowering Salt intake Is Cost-Effective and Cost-Saving

Lowering blood pressure by reducing salt intake even a small amount (15%) is estimated to prevent 8.5 million premature deaths in 10 years in low- and middle-economies and can deliver cost-savings in high-income countries (26-28). Three countries have proven that sustained wide-ranging initiatives can reduce salt intake population-wide and in two countries, health benefits have accrued:

- Between 1955-89 average salt consumption in Japan dropped from 13.5 g to 12.1 g per day resulting

in a gradual fall in blood pressure and a marked decline in deaths from stroke (29).

- Finland reduced population level salt intake by 25% over two decades beginning in the 1970s and similarly observed a marked reduction in blood pressure and stroke deaths (24).
- England reduced salt consumption in the population from 9.5 g in 2001 to 8.6 g in 2008 (30).

WHO Platforms for Dietary Salt Reduction

WHO and the UN Food and Agriculture Organization (FAO) issued a joint report in 2003 in which the recommended individual salt intake be less than 5 g/day. Then as part of the implementation of the WHO *Global Strategy on Diet, Physical Activity and Health (DPAS)* and the *2008-2013 Action Plan for the Global Strategy for the Prevention and Control of Non-communicable Diseases*, WHO established a salt reduction strategy with three platforms shown below.

WHO Population Salt Reduction Strategy

- WHO with the UK FSA jointly organized Platform 1 – Create enabling environments – in the summer of 2010. It included an information exchange forum with the private sector and NGOs on population-

based salt reduction strategies followed by a technical meeting. Discussed were interventions for consumer education and reformulation of industrially produced foods to enable consumers to make appropriate choices to reduce the total sodium content of their diet (31).

- In the fall of 2010, PAHO/WHO with the Government of Canada (Health Canada) organized Platform 2 – Evaluation and monitoring – also as an information exchange forum with the private sector and NGOs followed by a technical meeting where monitoring sodium intake levels at population level, assessment of dietary sources of sodium, and knowledge, attitudes and behaviours towards sodium and health were discussed (32).
- Platform 3 will bring together stakeholders in dietary salt reduction and salt fortification with iodine, to facilitate the coordination of the two strategies.

The PAHO Initiative – Cardiovascular Disease Prevention through Population-wide Dietary Salt Reduction

Background

Seminal to PAHO launching a regional dietary salt reduction initiative was the 2007 article in *The Lancet* – *Chronic disease prevention: health effect and financial costs of strategies to reduce salt intake and control tobacco use* (26). It concluded that a voluntary or legislated programme of reformulations by the food industry to use less salt and sodium additives in food products, combined with a public health campaign to raise consumer awareness of the dangers of high salt intake and encourage less salt use in household cooking and at the table, is the most cost-effective public health policy available to countries whether they are low-, middle- or high-income.

In 2008/09, PAHO surveyed the Member States in the central and south sub-regions as to national salt reduction activities underway or intended. The 12 responding countries¹ reported the following main aspects of their approaches:

- five (Argentina, Brazil, Chile, Costa Rica and Uruguay) had national recommendations to limit salt intake (four had the target of less than 5 g/day/person and one recommended less than 6 g);
- four had qualitative recommendations for moder-

ate use of salt (Grenada, Guyana, Panama and St. Vincent);

- five were conducting research on sources of sodium in the diet (Argentina, Bolivia, Brazil, Chile and Ecuador); and
- two had working groups/task forces leading national initiatives for salt intake reduction (Argentina and Chile).

With countries already demonstrating their intentions to reduce dietary salt, PAHO with the Canadian Public Health Agency organized a regional meeting in Miami in early 2009 as a forum to exchange information on action in and outside the region and to define what next steps would be of most assistance to advance the action in the Americas. Discussions identified a number of the key challenges to be taken into consideration as the region moved forward:

- Salt is still seen predominantly as a vehicle for micronutrient fortification (in particular for iodine and fluoride) therefore messages and actions to reduce salt intake must be synchronized with those addressing the prevention of micronutrient deficiency.
- Because many Central American and Caribbean countries import much or most of their foods, it is essential for salt to be on the agendas of regional political / trade bodies.
- National food regulatory agencies are still rare in the region, existing only in Brazil, Canada, Chile, Costa Rica, Mexico and the United States, therefore instruments such as mandatory nutrition labeling and nutrient analysis and monitoring, proven elsewhere to be effective in influencing the food industry to remove or reduce harmful ingredients like trans fats and high salt, are uncommon in the region. The public commitment of many countries to the "right to health" of their populations can be used as an argument while regulatory capacities grow.
- Small and medium enterprises and the informal food sector are very common in the region, with products from the latter virtually uncontrolled. This "atomization" of food provision poses special challenges for engagement, effective collaboration and monitoring.
- Taking action now is critical. Trade liberalization and economic growth in several countries in the region, particularly in Central and South America, are increasing consumer purchasing power and as a result, changing lifestyles. Among the changes is a shift in food preferences from home cooking to greater consumption of commercially processed products, many of them highly salted, that multinational and large national companies are increasingly making available on local markets. At this juncture, reducing the salt content across the food supply in the context of promoting healthy diet and lifestyles is timely, to the benefit of population

¹ Argentina, Bolivia, Brazil, CFNI (Bahamas, Dominica, Grenada, Guyana, St. Lucia, St. Vincent and the Grenadines) Chile, Costa Rica, Ecuador, Honduras, INCAP (Guatemala) Panama, Paraguay, Uruguay.

health and by extension supporting the economic development apparent in the region.

The key outcome of the Miami meeting was the call for PAHO to lead a region-wide initiative to mobilize Member States to reduce the overconsumption of salt. PAHO responded, convening in September 2009 a group of independent international experts on salt and health from within and outside the region to guide and support an initiative – Cardiovascular Disease Prevention through Population-wide Dietary Salt Reduction – encompassing North, Central and South America and the Caribbean, intended as multifaceted and multistakeholder.

The Expert Group and its Sub-groups

The Expert Group comprises academics and researchers, and representatives of international health organizations, consumers, non-government organizations active in health and of health institutions in Member States working on dietary salt reduction and cardiovascular and chronic disease prevention. The Group operated with a Chair and two co-chairs, one each for the Caribbean and Latin America, supported by a secretariat led by PAHO.

The Group was to issue recommendations for strategies and interventions in line with the WHO strategy that can be employed to reduce salt intake levels among people in the Americas. Its terms of reference to serve the region and sub-regions were as follows:

- Estimate the impact of a successful program to reduce salt consumption in the Americas on hypertension and cardiovascular disease prevalence.
- Initiate regional contacts with industry.
- Identify advocates in the region.
- Connect main players in the region.
 - Prepare a Policy Statement on salt reduction to be signed by key players in the region.
- Develop common resources.
 - Sets of common messages to raise awareness among consumers and healthcare professionals.
 - Guidance for surveillance and monitoring.
- Provide guidance and support directly to governments (e.g. national task forces, national awareness raising campaigns, national target setting, etc.).
 - Support establishment of national task forces.
 - Support national efforts to build awareness in governments and the general population in countries regarding salt reduction.
 - Identify necessary technical expertise for development of country targets and timelines setting.

The Group divided itself into sub-groups and with additional technical expertise, developed policy recommendations and resources (products) for the areas that reflected its terms of reference and were seen as most helpful to Member States (already taking action as well

For the governments of Uruguay and Costa Rica, their endorsement of the Policy Statement served as a foundation for launching national initiatives.

For civil society in Latin America, led by Consumers International and the Inter-American Heart Foundation, the Policy Statement was the basis for organizing around information exchange, for mobilizing action and providing support for the regional initiative.

The Healthy Caribbean Coalition has developed culturally tailored material to raise awareness of high salt diets and their harmful effects. The Heart Federation of Jamaica promoted “Drop the Salt – Protect Your Health” as the theme for Salt Awareness Week in 2011.

More than 40 Latin American civil society organizations launched the Healthy Latin American Coalition (Coalición Latino América Saludable -CLAS) in Buenos Aires in early 2011. It declared the grave impact of non-communicable diseases in the region – a deterrent to human development and obstacle in preventing poverty. The Coalition developed an initiative specific to salt and health (ALASS –Asociación Latino Americana Sal o Salud) for the exchange of information, to build proposals for funding and provide different platforms for knowledge dissemination.

The Chilean affiliate of WASH International was launched in June 2011.

as those not yet active): engaging the food industry on product reformulation and new product development; advocacy and communication; surveillance of salt intake, sources of salt in the diet and public knowledge and opinions on salt and health; salt fortification with iodine; and promoting national level health economic impact studies on salt reduction. The Group and various members published the advances made in the re-

Three countries in the region have applied the 24-hour sodium excretion Protocol:

- the SALMEX study, conducted by the National Medical Science and Nutrition Institute Salvador Zubiran over 2010/11 involving a worker cohort, is the first 24-hour urinary sodium excretion study in Mexico;

- the Chronic Disease Research Centre in Barbados determined a national baseline for sodium excretion in 24-hour urine in 2010/11 as part of a national Salt Intake Study;

- the New York City Department of Health and Mental Hygiene measured sodium excretion in 24-hour urine samples from 1700 resident non-institutionalized adults in 2010 (33).

A food frequency questionnaire first developed for the Barbados National Cancer Study in 2000 was adjusted in 2010 to account for changes in the Barbadian diet and at the same time allow a specific evaluation of salt consumption.

The National Household Budget Survey (HBS – Pesquisa de Orçamentos Familiares) administered between July 2002 and June 2003 by the Brazilian Institute of Geography and Statistics (IBGE, Instituto Brasileiro de Geografia e Estatística) provided the data from which the main sources of salt in the Brazilian diet were derived (7).

gion and promoted the initiative at regional and global meetings and conferences. A scientific review committee was also formed to examine and respond to new research released during the period of the Group's mandate.

Purpose of this Report

This report marks the completion of the initial two-year mandate of the initiative. It presents the work of the Expert Group, the progress made to date by countries in the region and the Expert Group's key messages and recom-

mendations for further advancement directed to the major stakeholders in dietary salt reduction: Member States; the food and salt industries; non-government organizations (NGOs) and civil society; PAHO/WHO and other international organizations involved in dietary salt reduction and salt iodization.

Tools, Resources and Achievements

The Expert Group and sub-groups developed tools and resources and formulated guidance to assist Member States to start and strengthen national strategies for dietary salt reduction. In some cases, tools and resources were already available and with assistance from PAHO and Expert Group members, were made culturally appropriate and translated where necessary. In other cases, the Expert Group and sub-groups initiated the development of materials, this through participative processes often involving additional technical expertise, supported by PAHO. Described below are the products prepared for each of the areas to which the Expert Group applied itself plus there are examples of achievements relevant to each.

Advocacy and Communication

The Policy Statement

The Expert Group's first product – a Policy Statement, the road map for the initiative – was launched in November 2009 in Santiago Chile at a symposium organized during a meeting of the Latin American Society of Nutrition (SLAN). It set the goal – the internationally recommended intake of less than 5 g salt per capita per day (in the absence of equivalent or lower national targets) – to be reached by 2020 among people in the Americas.

The intended audience for the Statement is policy and decision makers in government; leaders in non-governmental organizations representing consumers, health, scientific and healthcare professionals; civil society; the food industry (including food processors and distributors) and food importers and exporters; and PAHO. As of May 2011, there are 56 endorsements of the Policy Statement, listed in Appendix 1.

Fact sheets, presentations, core references

The sub-group adjusted a standardized presentation on salt reduction and a set of fact sheets to be culturally appropriate, covering salt and hypertension, salt and osteoporosis and salt and kidney failure. The materials are intended for the general public, health professionals and policy makers. The sub-group also compiled a list of core references that underpin the scientific basis and benefits of dietary salt reduction.

Information dissemination

Over the course of its two-year mandate, the Expert Group issued press releases and offered webinars through PAHO; published articles in peer reviewed journals; members made presentations at international conferences and meetings; and otherwise, experiences, scientific information and the Policy Statement were made available at relevant events. Appendix 2 has a complete list of dissemination activities.

Regional support for the WHO Platform I Meeting on Creating Supportive Environments

Members of the Expert Group participated in the Platform I Meeting in July 2010 in London, joining representatives from other WHO regions. They presented the Pan American initiative, the profile of the Expert Group and its sub-groups, their objectives and achievements to date.

Surveillance

Surveillance of actual salt intake, determining the main sources of salt in the diet and understanding what people know about salt and its effects on health are the critical underpinnings to the design of a national effort to reduce the consumption of salt. They point out where to place the emphasis and what goals to set. With baselines known, ongoing monitoring demonstrates whether goals are being met.

Protocol for Determining Population Level Sodium Intake in 24-hour Urine Samples

With support from the National Institute of Public Health in Cuernavaca (Mexico) and the Salvador Zubiran National Institute of Medical Sciences and Nutrition in Mexico City, the sub-group for surveillance prepared the *Protocol for Population Level Sodium Determination in 24-hour Urine Samples*. The 24-hour urine sampling method applied to a representative sample of the population provides the most valid and reliable data on salt intake in a country. The Protocol also recommends the concurrent assessment of potassium and iodine intakes as they are important dietary constituents to take into account in salt reduction programs.

While secondary sources can supply estimates of salt intake, there should be provision made for 24-hour urine sampling to confirm findings.

A Review of Methods to Determine Main Sources of Salt in the Diet

With support from the Brazilian Ministry of Health, the sub-group developed the Review, a companion document to the *Protocol*. It serves as a guide to selecting the most appropriate method, given country resources and circumstances, to identify the complete profile of dietary sources

of salt as baseline information and for ongoing monitoring:

- foods that people consume and the amounts and frequency of consumption;
- salt content of the most commonly consumed foods;
- the amount of salt added at the table and in cooking;
- intake of high-salt foods that are culturally or regionally-specific.

Literature review of 24-hour urine and spot sampling methods

A systematic review of literature, underway, is assessing the validity and reliability of sodium and iodine intake values derived from spot urine samples to estimate salt and iodine consumption compared to the 24-hour urinary excretion method considered to be the gold standard. The analysis of multiple studies has so far found current spot urine methods to be unreliable in representing daily sodium consumption.

Studying knowledge, attitudes and behaviour

Expert group members initiated two projects directed at consumers' knowledge, attitudes and behaviours (KAB) regarding salt/sodium and food labels. The Consumers International (CI) representative on the Expert Group led the first project, assisted by other Group members, in which a KAB instrument and methodology were developed and then tested in five countries among consumers approached in an ad hoc manner on the street. The findings informed a meeting of the CODEX Alimentarius Committee on Food Labeling (see below). A second phase on KAB, to involve interviews with focus groups applying qualitative research methods, is being led by INCIENSA (Costa Rican Institute for Research and Education on Nutrition and Health) working with researchers in Costa Rica, Ecuador and Argentina.

Building regional capacity for food composition data

Essential to an assessment of the main sources of salt in the diet are up-to-date food composition data. Specific to foods in Latin America is the database of LATINFOODS – the Latin American Network of Food Data Systems (34). In 2011, the Expert Group member from Costa Rica facilitated a survey of LATINFOODS members and related food analysis laboratories in 19 countries on the status of their data on the sodium content of foods (35).

Participation in an international collaborative project to compare and monitor the nutritional composition of processed foods

Ten countries in the Americas region (Argentina, Barbados, Brazil, Canada, Costa Rica, Ecuador, Guatemala, Mexico, Panama, Peru) are participating with eight others elsewhere in an international initiative to survey processed foods in major food categories using a standardized methodology. The objective is to com-

Consumers International together with researchers in Chile, Argentina, Ecuador, Costa Rica and Canada tested the KAB instrument. The results (unpublished) showed poor knowledge on salt levels in food and of the effect of salt on health both in the general population as well as among individuals with hypertension, indicating that more knowledge dissemination on the topic is required.

Knowledge, perceptions and behaviour in relation to salt intake and dietary sodium, its association with health and the nutrition labeling declaration in foods. Multicenter collaborative study sponsored by PAHO/WHO, 2011-2012. In Argentina, Costa Rica and Ecuador, in-depth semi-structured interviews with two focus groups in each country, one rural and one urban, six adults in each (12 total), will explore KAB issues, supplemented by direct observation.

The LATINFOODS members and other laboratories responded to a questionnaire on whether they had data on the sodium content in 14 categories of processed and pre-prepared foods. While 68% of respondents reported having some information on sodium content in all food categories (mainly breads, cereals and snacks), all respondents indicate that data need to be updated and a high majority requires resources and training to do so. All respondents are supportive of the regional initiative to reduce dietary salt.

pare the nutritional composition between countries, between food companies and over time. The George Institute for Global Health in Australia is coordinating and secretariat to the project.

Regional support for the WHO Platform II Meeting on Surveillance for Population Salt Reduction
PAHO with the Government of Canada, assisted by

the Chair of the Expert Group, co-hosted a WHO Platform II meeting on surveillance in October 2010 in Calgary, Alberta. Expert Group members contributed the knowledge and experiences with surveillance in the region and featured the related products – the *Protocol for Population Level Sodium Determination in 24-hour Urine Samples*, the results of the KAB survey by CI and research partners, the planned focus group based research on KAB, and the findings up to that point from the review of literature regarding 24-hour urine sampling versus spot urine methods to determine sodium intake.

The outcome was the WHO publication *Strategies to monitor and evaluate population sodium consumption and sources of sodium in the diet – Report of a joint technical meeting by WHO and the Government of Canada, Canada, October 2010.*

Food Industry Engagement

Industry questionnaire (for multinational and national food manufacturers)

The PAHO secretariat with members of the industry sub-group engaged with several multinational food companies that participated in the PAHO 2009 Partners Forum and had joined a healthy eating working group to continue their involvement. These Forum participants agreed to pilot a questionnaire with two objectives:

- determine a baseline on the nature and extent to which commercial food processors are reformulating food products to reduce their sodium content or are formulating new product lines with low/no sodium; and
- identify the leaders in (re)formulation

Questions were specific to reformulation plans and efforts; the food categories/food items in company portfolios undergoing salt content reductions or low salt items being supplied in the region; and the food companies' interests in various actions to be coordinated with PAHO to support dietary salt reduction.

The questionnaire was also designed as a resource for Member States to facilitate their engagement with and understanding of the intentions and efforts of national and multinational food manufacturers whose products are available on their respective national markets.

Joint PAHO and World Economic Forum on Latin America meeting and the “2011 Statement of Rio de Janeiro”

PAHO with members of the Expert Group collaborated with the World Economic Forum (WEF) on Latin America and representatives of the food industry, governments, regulatory agencies and civil society to produce

Dietary Sodium/Salt Reduction in the Americas 2011 Statement of Rio de Janeiro. It is the first position document created jointly by governments, the food industry and civil society that distills elements in the Policy Statement into a set of priorities and commitments regarding their respective roles and responsibilities in reducing the overconsumption of salt. It is a potential contribution to the PAHO/WEF submission to the high level meeting of the UN General Assembly on the prevention and control of non-communicable diseases in September 2011. It is also expected to facilitate dialogue on salt reduction at regional and national levels beyond the UN meeting.

Informing committees of CODEX Alimentarius

Expert Group members joined the International Workshop on Salt and Health in October 2010 in Santiago Chile to contribute to the CODEX Committee on Nutrition and Foods for Special Dietary Uses (CCNFSDU) dealing with nutrient reference values for nutrients (like sodium) associated with diet-related risks for non-communicable diseases. The Workshop, promoting salt/sodium content of foods to be framed as a food safety issue, provided additional information and evidence to CCNFSDU.

PAHO on behalf of the Expert Group also made submissions to directly inform two meetings of the CODEX Alimentarius Committee on Food Labelling (CCFL) in 2010 and 2011. The Expert Group position was:

- 1) That it be mandatory for sodium/salt content to be declared on food nutrition labels whether labelling is voluntary or mandatory;
- 2) That the decision about whether sodium or salt is declared be determined by nationally recognized competent scientific authorities in each country;
- 3) That it be mandatory for sodium or salt content to be effectively communicated to consumers as determined by nationally recognized competent scientific authorities e.g. front-of-package information;
- 4) That the nutrient reference value for sodium be set at a level as low as possible in keeping with an achievable health promoting diet or be a limit, such as adequate intake, established by nationally recognized competent scientific authorities.

For the CCFL meeting, the results of the KAB survey on salt led by CI, where consumer understanding of salt and sodium on food labels was probed, were also submitted and highlighted in the meeting summary. The CCFL meeting conclusion reflected the Expert Group position – that while sodium is the correct scientific term to include on labels, national authorities can choose to have the total amount of sodium expressed as a salt equivalent on labels in cases where the public demonstrates a better understanding of the term salt than sodium.

Some key findings from the pilot:

- *The more supportive the local government e.g. raising consumer awareness about dietary salt, the easier it is for food companies to undertake salt reduction in their products. An informed consumer helps industry, otherwise a company that unilaterally lowers its salt content risks losing market share.*
- *Important to a multinational is for governments to play a strong role in encouraging local/regional food manufacturers to engage in reducing salt in their products.*
- *Food companies plan and operate on the basis of markets that in the case of multinationals can represent regions or sub-regions (e.g. common markets), not necessarily countries. Therefore engaging them to reformulate or anticipating product distribution may require a multi-country approach i.e. taking the market perspective as defined by industry.*

In mid 2010 the Secretariat interviewed representatives of five national initiatives where voluntary targets and timelines had been established or were in the process of being negotiated with the food industry – in Argentina, Brazil, Chile, Mexico and the US NSRI. The interviews gave insights to the facilitators and challenges to industry engagement and to government action in the context of working with industry. See section III Status of Action.

In the Statement of Rio de Janeiro, governments, the food industry and civil society together strongly commit to the principle of sustainable food consumption; recognize that international organizations have a strong role to play in convening stakeholders; and encourage stakeholders to act now to improve the health of the people of the Americas using a comprehensive multisectoral approach to specifically reduce dietary salt.

The framework for action in the White Paper has seven areas in which collaboration by salt iodization and salt reduction programmes is recommended: common and coordinated messaging; common advocacy platforms; concurrent surveillance; coordinated evaluation; strategic joint research; shared forums with relevant sectors of the food industry; and coordinated mapping of existing and needed resources and mobilizing them.

Researchers in Argentina estimated the burden of acute coronary heart disease and stroke and the cost-effectiveness of preventative population-based and clinical interventions. Two interventions were particularly cost-saving: lowering salt intake in the population by reducing salt in bread and multidrug therapy (36).

Researchers in Canada conducted a preliminary study of the effect of various levels of dietary sodium reduction on cardiovascular disease mortality and morbidity in 18 Latin American countries combined (Argentina, Bolivia, Brazil, Chile, Colombia, Costa Rica, Cuba, Dominican Republic, Ecuador, El Salvador, Guatemala, Mexico, Nicaragua, Panama, Paraguay, Peru, Uruguay and Venezuela). They found that if sodium intake fell by 10% every year from a current average of 3700 mg daily, the optimal intake (1200-1500 mg daily) would be reached in 11 years. There would be about 593,000 fewer CHD and stroke events and about 54,000 fewer deaths in these countries by the end of the period (37).

Synchronizing Salt Iodization and Salt Reduction Programs

White Paper on Improving Public Health in the Americas through Optimal Intake of Sodium and Iodine

A WHO supported technical consultation from 2007, the PAHO Expert Group's Policy Statement (2009) and an international Iodine Network resolution from February 2010 agree that there is no inherent conflict between salt iodization and salt reduction initiatives. Stepping off from this position, PAHO convened two meetings in early 2011: the first in January with a small group of international and regional technical experts on iodine deficiency disorders and dietary salt reduction, among them members of the Expert Group, who confirmed that collaboration between and synchronization of the two programmes to achieve a common goal – the optimal intake of sodium and iodine in the Americas – will be cost effective and of great public health benefit; and a second larger meeting in April where representatives of regional and international agencies working to optimize iodine supplementation along with PAHO and Expert Group members agreed to a *White Paper on Improving Public Health in the Americas through Optimal Intake of Sodium and Iodine*. The White Paper includes a framework for action with recommendations to national governments, civil society, various sectors of the salt and food industries and international organizations active in the region. Participants agreed to an immediate next step – the selection of pilot countries in the region where implementation of the two programs can be coordinated, guided by the framework for action, with the experiences to become case studies for dissemination.

Intervention with the US Institute of Food Technologists (IFT)

The framework for *Improving Public Health in the Americas through Optimal Intake of Sodium and Iodine* was featured during the IFT 2011 annual meeting, specifically the symposium on “The role of food processors world-wide in preventing iodine deficiency disorders”. Participants recognized that: processed foods, the main source of salt in high-income economies, should be the focus for reformulation; culinary and food science collaborations are necessary to address salt reductions in both retail and food service markets; and customized reformulation approaches are needed within and across food categories.

Nutrition Guidance Expert Advisory Group

Expert Group members have contributed to the WHO Nutrition Guidance Expert Advisory Group (NUGAG) as

it reviews the evidence on how varying levels of population salt intake can impact the effectiveness of salt iodization programs.

Economic Studies on the Cost-Effectiveness and Cost-Savings of Population Level Dietary Salt Reduction

The sub-group on health economic analysis identified two readily available models that can demonstrate the health and economic development benefits of population level dietary salt reduction using a minimal data set: the WHO CHOICE model (generalized cost-effectiveness) and the Coronary Heart Disease (CHD) Policy Model. PAHO and University of California in San Francisco are collaborating to prepare countries in the region to evaluate the projected and observed effectiveness and cost-effectiveness of their national dietary salt reduction programs, aiming to maximize impacts within each national context.

Status of Action in the Americas

This section begins with the country-specific summaries of activities relevant to dietary salt reduction collected through a 2011 survey of Member States admin-

istered by the secretariat. A standardized questionnaire was used to determine:

- which Member States have and do not have national initiatives to reduce the overconsumption of salt;
- the nature of the initiatives currently underway, what has facilitated them, the barriers encountered and the kind of assistance needed to make further progress;
- the barriers in countries with no initiatives to reduce dietary salt at the population level and what would facilitate overcoming them.

Following the country status reports is the description of food categories for which salt content reductions are underway or intended in the most active countries in the region. Then there is a compilation of the lessons learned so far and challenges to be addressed. The section concludes with a synthesis of the momentum and potential for specific advancements in the region to inform a next phase of the initiative.

Country-specific Summaries

The table below presents countries that have national scale initiatives underway. This is followed by a synthesis of the information reported by countries not yet active.

Summaries for countries reporting strategies

Strategy	Key activities	Lead and participating organizations
Country: ARGENTINA		
<p>"Menos Sal Más Vida" (less salt, more life) launched in 2010, part of CVD prevention and the National Strategy for the Prevention and Control of NCDs. Lead organizations are the Ministry of Health, National Nutrition Institute, National Institute of Industrial Technology, and National Commission for NCD Prevention and Control (the latter comprising ministries of health, agriculture, social development, science and technology; the National Institute of Industrial Technology, food companies, NGOs, consumers and universities).</p> <p>Recommended intake 6 g salt per food code.</p>	<p>Mass media campaigns; creation/dissemination of education materials; meetings with the food industry and voluntary industry target setting; investigation to determine main sources of salt; assessing cost-effectiveness of reducing overconsumption of salt.</p> <p>"Menos Sal Más Vida" is extending to large food industries with salt content baseline analysis and target setting. As of October 2011, 2 and 4 year targets are set for reductions of salt content (between 5 and 18%) for various products and commitment obtained from 41 leading food industries and signed. Otherwise under development are surveillance plans and monitoring strategies: food analysis; labelling and self-reporting of sodium content data; a national nutrition survey for 2012 with a sub-sample of participants to supply a 24-hour urine sample for sodium analysis; a household expenditure survey in 2012.</p> <p>The salt content of French bread and other bakery products has already been reduced by 25%.</p>	<p>Secretary of Regulatory Institute, Secretary of Health Policy, National Institute of Nutrition.</p>

Strategy	Key activities	Lead and participating organizations
Country: BRAZIL		
<p>National Plan for Reducing Salt Consumption in Brazil, complementing the national nutrition and CVD prevention strategies and the overall National Plan for Tackling Chronic NCDs (2011-2012). Launched in 2011, under the general joint coordination of the Ministry of Health and the national food regulatory agency (ANVISA) and involving the Brazilian Association of Food Industries (ABIA).</p> <p>Recommended intake 5 g salt (2000 mg sodium)/day/person.</p>	<p>Mass media campaigns (“Menos Sal”) with e.g. a website about salt reduction; the food guide for the Brazilian population; Primary Health Books; other materials to educate and sensitize consumers, food industries, health professionals and other partners; nutrition facts labels that include salt; meetings with the food industry and voluntary target setting for specific food categories; Family Budget Survey Food Availability Modules used to estimate current salt intake levels; research to determine main sources of salt; plans to coordinate the salt reduction program with iodine supplementation program.</p>	<p>National food regulatory agency (ANVISA), Brazilian Associations of Food Industries, of Pasta and Dough Industries, of Wheat and Bakeries, Brazilian Association of Supermarkets (ABRAS), National Health Council, National Food and Nutrition Security Council.</p>
Country: CANADA		
<p>2010 Sodium Reduction Strategy for Canada, Recommendations of the Sodium Working Group (SWG) – calls for reduction of population level sodium intake from the current 3400 mg/day to an average of 2300 mg/day by 2016 with an eventual goal of 95% of the population consuming less than 2300 mg/day.</p> <p>Draft sodium reduction targets for 2016 for a number of food categories with interim milestone targets for 2012 and 2014.</p>	<p>Sodium Reduction Strategy has four action areas: structured voluntary sodium reductions in food; public education campaigns; food science and health research related to sodium; planned, periodic monitoring and reporting of sodium intake levels and sodium reduction program evaluation.</p> <p>Detailed actions [38]: estimation of the impact of high dietary sodium on hypertension and CVD and costs of health care; development of tools to educate the public, healthcare professionals and policy makers; clinical and scientific sessions on salt and health at regional and national meetings of national health NGOs; extensive national media campaigns led by NGOs regarding high salt content of processed, manufactured and restaurant foods and negative health effects; government research to determine main sources of dietary salt and population survey of 24-hour urinary excretion; studies on knowledge, attitudes and behaviours of the general population and hypertensives relating to dietary salt; research and development of food procurement policies for public institutions; advocacy to restrict marketing of food and beverages to children; combined funding from Canadian Institute for Health Research, NGO and private sector for the Canada Chair in Hypertension Prevention and Control.</p> <p>Federal, Provincial and Territorial (FPT) Health Ministers agree to a target of sodium intake of less than 2300 mg/day. An FPT Sodium Task Group is reviewing the recommendations of the SWG including the identification of potential FPT regulatory mechanisms to achieve the 2016 target. The first full report of targets and timelines for food reformulations, a monitoring and evaluation program and a social marketing program are expected in the fall of 2011.</p>	<p>Health Canada, Canadian Public Health Agency, Canadian Institute for Health Research, Statistics Canada, provincial and territorial ministries of health, the food industry, health NGOs, scientific organizations.</p>

Strategy	Key activities	Lead and participating organizations
Country: CHILE		
<p>Strategy to Reduce Salt/Sodium Consumption in Chile, complementing the Strategy Against Obesity, the Nutrition Intervention Strategy through the human life cycle, the Cardiovascular Health Program and the Choose Healthy Living Campaign that is focused on a healthy population especially children. Leadership lies with the Ministry of Health, Subsecretary of Public Health, Division of Public Policy, Department of Food and Nutrition.</p> <p>Recommended intake 1200-1500 mg sodium/day, according to age for children under 18.</p>	<p>Mass media campaigns; nutrition facts labels that include salt; front of pack labelling; creation/dissemination of education materials; talks at academic meetings for societies of hypertension, cardiology, nephrology, and paediatrics; meetings with the industry and voluntary target setting; survey of physical measurements to estimate current level of dietary intake; research to determine main sources of intake; question included in National Quality of Life Survey on how often table salt is added to food; and development of a public consultation with the following objectives: understand the population's opinion on salt reduction; inform new feasible strategies to reduce salt intake; and increase the population's perception of the risks of high salt intake.</p>	<p>Medical societies of hypertension, cardiology, pediatrics, and nephrology; Food industry; Ministry of Health; associations of artisanal bread makers and supermarkets.</p>
Country: COSTA RICA		
<p>National Plan to Reduce the Consumption of Salt/Sodium in the Population of Costa Rica within the National Policy on Food Security and Nutrition (2011-2021), led by the Ministry of Health, Institute for Research and Education in Nutrition and Health [39].</p> <p>Recommended intake 5 g salt/day/person.</p>	<p>Action plan based on the WHO Population Salt Reduction Strategy and includes research to gather the evidence base to support the three platforms in Costa Rica: KAB regarding salt consumption and health in one population group and sodium analysis of two specific food groups is ongoing; previous research needs to be completed with other representative populations and food groups; and estimation of the impact of high dietary sodium on hypertension and CVD and costs of health care.</p> <p>In the next 12 months: mass media campaigns; workshops on how to reduce dietary salt; creation/dissemination of educational tools; meetings with the industry including food service establishments (restaurants); survey of physical measurements to estimate current level of dietary intake; identification of sources of salt; coordinating the salt reduction program with the iodine and fluoride supplementation programs; monitoring and evaluation.</p>	
Country: CUBA		
<p>Salt reduction is among the comprehensive actions and governmental policies of the 2010 National Program of Non-Communicable Diseases, led by Ministry of Public Health, NCD Department, National Institute of Nutrition and Food Hygiene.</p> <p>Recommended intake of no more than 2300 mg sodium / day (5 g of table salt).</p>	<p>Mass media campaigns; nutrition facts labels that include salt; workshops on how to reduce dietary salt; national implementation of Cuban dietary guidelines; meetings with the industry; studies on knowledge attitudes and beliefs; coordinate salt reduction program with iodine supplementation program.</p>	<p>Progressive actions are taken for a period of three years by the Ministry of Food Industry of Cuba to complete the global strategy to reduce salt.</p>

Strategy	Key activities	Lead and participating organizations
Country: MEXICO		
<p>Acuerdo Nacional para la Salud Alimentaria, Estrategia para el Sobrepeso y la Obesidad, Programa de Acción en el Contexto Escolar. Specific to salt is “Mas Agua, Menos Sal”.</p> <p>Recommended intake of <2g/day of sodium.</p>	<p>Mass media campaigns; nutrition facts labels that include salt; front of pack labelling; limit foods high in sodium within preschool, primary and secondary schools; creation/dissemination of educational tools; meetings with the industry; voluntary agreement with goals and timeline for specific food categories; survey of physical measurements to estimate current level of dietary intake; other survey (national nutrition and health survey 2012); cost effective analysis.</p>	<p>Secretary of Health/Sub-secretary of prevention and promotion, Secretary of public education, Secretary of the economy.</p>
Country: SURINAME		
<p>National Approach for Reducing salt use in our food/ Een Nationale Aanpak ter Vermindering van het Zoutgebruik in ons Voedsel), launched in 2010, led by the Ministry of Health.</p>	<p>Action to being in the next 12 months: Mass media campaign; nutrition labels that include salt; front of pack labelling; workshops on how to reduce dietary salt, creation/dissemination of educational tools; meetings with the industry; a survey of physical measurements to estimate current dietary intake; research to determine main sources of salt in the diet.</p>	
Country: UNITED STATES		
<p>US actions are not housed under one program. At the federal level activities include national goals (Healthy People 2020), and Dietary Guidelines for Americans (see below), labelling initiatives of FDA and USDA, educational efforts including the National High Blood Pressure Education Program, scientific and monitoring efforts, procurement strategies and funding community efforts. The National Salt Reduction Initiative (NSRI) is a partnership of 70+ city and state health authorities and health organizations focusing on reducing sodium in packaged and restaurant foods. Complete details of government initiatives are given in Appendix B of the IOM Strategies to Reduce Sodium in the United States [10].</p> <p>The 2010 Dietary Guidelines for Americans recommend limiting intake of sodium to 1500 mg per day for people aged 51 years and older, African Americans, and those who have high blood pressure, diabetes, or chronic kidney disease (the majority of US adults). All others should reduce sodium intake to less than 2300 mg per day.</p>	<p>Mass media campaigns; nutrition labels that include sodium; proposed rule by USDA for labelling of enhanced meats and sodium information available upon request in restaurants; workshops on how to reduce dietary salt; creation and dissemination of education tools; meetings with the industry (NSRI set food category sodium reduction targets for 2012 and 2014, received voluntary corporate commitments, and is monitoring progress through the creation of packaged and restaurant food data bases); surveys to determine salt intake (National Health and Nutrition Examination Survey (NHANES) and the NYC 24-hour urinary sodium study associated with NSRI); studies on knowledge attitudes and beliefs regarding salt consumption and health; assessing health benefits and cost-effectiveness of salt reduction programs; coordinating salt reduction with iodine supplementation programs; food procurement policies for public institutions – the Health and Sustainability Guidelines for Federal Concessions and Vending Operations.</p> <p>CDC is currently conducting a 24-hr urine calibration study and examining trends in estimates of 24-hour sodium excretion using spot urines from stored samples. Proposing to conduct 24-hr urine collections on a sub-sample of adult NHANES participants starting in 2013-2014.</p>	<p>FDA, CDC, NIH, USDA, NSRI.</p> <p>Several food processors, manufacturers and retailers have committed to lower sodium in their products. See Appendix 3, US NSRI Corporate Commitments and Comments.</p>
Country: URUGUAY		
<p>No formal title to the initiative, part of the national NCD prevention and national nutrition programs, led by the Ministry of Health.</p> <p>Recommended intake less than 5 g salt/day/person.</p>	<p>Nutrition labels that include salt; nutrition education using the food-based dietary guidelines; workshops on how to reduce dietary salt; creation/dissemination of educational tools; meetings with the industry; other survey method to determine salt intake – National Survey of Expenditures and Household Income.</p> <p>Bakers Industrial Centre of Uruguay plans to decrease the sodium content of baked goods on a voluntary basis.</p>	

Summary for countries reporting no current plan or plans under development

The seven countries of Central America (Belize, Costa Rica, El Salvador, Guatemala, Honduras, Nicaragua and Panama) and the Dominican Republic through El Consejo de Ministros de Salud de Centroamérica (COMISCA) agreed to a common health plan – Plan de Salud de Centroamérica y República Dominicana 2010-2015 – that includes the commitment to set policies to regulate sodium and trans fats levels in food products (40). At the same time Costa Rica (see table above) and Guatemala have national strategies relevant to salt intake reduction. But while Guatemala reported mass media campaigns, nutrition labels that include salt, a survey of physical measurements to estimate current level of dietary intake, application of an inventory method to estimate current salt intake and a recommended per capita salt intake of less than 5 g per day, full implementation is dependent on synchronization with the iodine fortification program.

Bolivia reported no current plan to reduce population level dietary salt while Ecuador and Paraguay have plans in development but no implementation timeline. Nevertheless, countries report activities underway that are relevant to addressing the overconsumption of salt e.g. mass media campaigns, nutrition labels that include salt, workshops on how to reduce dietary salt, creation/dissemination of educational tools and KAB studies regarding salt. Columbia reports that an initiative is under consideration but no decision has been made on how to start, largely due to overconsumption of salt not yet being recognized because of a lack of data to make the case (e.g. on salt intake, the main sources of salt in the diet, nutritional patterns and the household food basket (the latter needed by administrative regions in the country). There is a possibility of adding questions on salt to the national study on nutrition and health. Of benefit would be technical support to design a national strategy including the research to determine baselines for and subsequent monitoring of key indicators. To date, Columbia has nutrition labels and conducts public health mass media campaigns.

Food Categories for which Salt Content Is being Reduced

As of July 2011, national public health authorities have or are in the process of negotiating national scale voluntary targets and timelines with the food industry in six countries in the Region of the Americas – three in the southern cone (Argentina, Brazil, Chile), and three in

the north (Mexico, Canada and the US NSRI). The food categories currently being addressed in each country are shown below divided into two broad groups –packaged and unpackaged foods. At this point, common to all countries except Mexico is government and food sector collaboration to reduce the salt content of bread (artisanal bread, packaged breads or both) and packaged meat products.

Argentina	Packaged	Unpackaged
	meats and derivatives; dairy products; cookies and derivatives; soups and dressings.	artisanal bakery bread
Brazil	Packaged	Unpackaged
	instant pasta (cup noodles and ramen pasta); industrially produced bread and buns; cakes and cake mixes; snacks; processed meat products; condiments and sauces; margarines; mayonnaise; breakfast cereals; dairy products; industrially produced meals; biscuits; cookies; ready meals.	artisanal bakery bread
Canada	Packaged (including restaurant foods)	
	bakery products; cereals and ready-to-eat cereals; dairy products; fats and oils; fish products; combined dishes; processed meat products; sauces; soups; snacks; sauces, dips, gravies, condiments; canned vegetables; fresh meat products; seasoning mixes; pasta and noodles; legumes; drinks and beverages; meat alternatives; baby and infant food; rice; nut butter.	
Chile	Packaged	Unpackaged
	sausages, cheese	artisanal bakery bread and private label supermarket bread
Mexico	Foods supplied to and available in the school environment (limiting salt plus fat and sugar content and the caloric value per portion).	
US NSRI	Packaged	
	bakery products; cereal and other grain products; meats; dairy products and substitutes; fats and oils; sauces, dips, gravies and condiments; snacks; soups; potatoes; mixed dishes; vegetables; legumes; canned fish; seasoning mixes; nut butters. restaurant foods: hamburgers; chicken; seafood; sandwiches; breakfast sandwiches; pizza; Mexican food; potatoes; soup; bakery products.	

All countries have opted for structured gradual voluntary food reformulations (with targets, timelines and monitoring of progress). All except Mexico have at this point schedules of interim targets and timelines for specific food categories. Links to the detailed country-specific food categories, targets and timelines are in Appendix 3.

Common action on bread

Bread is a staple in national diets across the Americas, consumed in consistent quantities by virtually all segments of the population and on the whole has relatively high salt content. Argentina, Brazil, Canada, Chile and the US NSRI all include bread in their salt reduction initiatives. In the three southern cone countries – Argentina, Brazil and Chile – where a limited number of food categories is currently the focus for salt content reductions, bread was either the first or among the first products to be addressed. The experiences in these countries are highlighted below as examples especially relevant to countries that have not yet addressed salt reduction of how a national initiative can begin with bread. For Argentina and Chile in particular, the countries were able to work with existing or few additional resources and in the case of Argentina, have already achieved a 25% reduction in the salt content of bread.

Lessons Learned So Far

The two information gathering exercises conducted by the secretariat (in mid 2010 and mid 2011) provided insights into what has in general terms facilitated and hindered their actions and attempts to reduce the over-consumption of salt on national scales and in specific terms, what the countries have learned in the process of engaging the food industries to reduce their use of salt. Findings are summarized below.

The regional initiative and the activities of the Expert Group have had a positive impact

Countries reported the following about the initiative and the Expert Group:

- has helped advocacy to prioritize the issue as a national public health problem concerning CVD and its relationship to high salt consumption;
- the tools, resources and recommendations are informing and assisting with national plans and strategies;
- assisted with analysis of available evidence and identification of effective interventions;
- has made available a network of accessible experts to provide advice and collaborate on research;

- provided updates on experiences across the Americas;
- has provided the opportunity to explore the current epidemiological situation and promoted a review of policies and interventions and development of recommendations based on evidence;
- the evidence and recommendations for the standardized methodology to measure and monitor salt intake in the population allows national results to be compared with the rest of Latin America permitting an evaluation of national interventions;
- facilitated collaborations between countries and members of the LATINFOODS network, coupled with information on how issues are being handled elsewhere.

Many experiences with industry engagement are transferrable

- Upon entering into negotiations of targets and timelines, most important is information about what food companies have already achieved elsewhere as evidence that reformulation and new product development are feasible. Very relevant in the region are the experiences of low- and middle-income countries (LMIC) with similar resources and capacities. It is understood to be many times easier to implement changes if it can be shown that in other countries the same or similar changes are being made.
- Where regulation to limit salt content is pending, industry has appeared particularly motivated to voluntarily reformulate to reduce the market impact and extent of formulation changes necessary once regulations come into effect.
- Some countries in the region that are actively engaging with the food industry are doing so through national associations that represent various sectors of the industry e.g. artisanal bread makers, meat producers, supermarkets, food and beverage associations, etc. Other countries have found that directly approaching major and progressive food manufacturers has been effective in reaching reformulation target and timeline agreements.
- Positive relationships between public health and the food industry, established e.g. in Chile and Argentina with associations of artisanal bread makers when the use of fortified flour was mandated, facilitated the launch of national dietary salt reduction first through bread products. Bakers were supported by national public campaigns

promoting lower salt intake that featured the changes in bread products. In Chile, small bakery businesses even stated a preference for regulation on an acceptable limit for salt in bread products to “level the playing field”.

- The scale of salt content reductions across food categories is dependent on national capacities to first engage food industries in target and timeline setting and then subsequently monitor progress. Canada and the NSRI have engaged food manufacturers and chain-restaurant for salt content reductions across all food categories with products that contain salt. On the other hand, the southern cone countries have begun with food categories that are the main contributors of salt to national diets and are gradually adding other categories.
- A draft national action plan can be useful in engaging stakeholders to advance dietary salt reduction. The plan can be tabled with groups convened to consider the issue, presented as a template awaiting concrete commitments to emerge from stakeholders through a participatory process of priority and target setting.
- Argentina, Canada and the United States have identified failed experiments with fully voluntary approaches and self-regulation. In Argentina for example, the Ministry of Health was initially willing to accept self-regulation and engaged national food industry associations on this basis. When results were not forthcoming, the government intervened to stimulate action with structured voluntary targets and timelines and monitoring of progress. Similarly in the United States, voluntarily reductions in the use of salt additives in food, relying principally on consumer pressure, without close government oversight and monitoring, produced no meaningful declines in salt additive use or salt intake at the population level (9).

Challenges

- Considered a major barrier in LMIC is the lack of up-to-date objective measures of dietary patterns, the main sources of salt in the diet and levels of salt intake. The information that is available often comes from a few studies that are not representative of the national population or is derived from food consumption methodologies developed for other purposes however scaling up or instituting specific national surveys requires capacity and infrastructures that are not available. Countries have relied on secondary data sources

e.g. household budget surveys and production and sales data from the food industry from which food consumption and salt intake information has been derived. While the information has been used effectively to launch dietary salt reduction initiatives, with regards to salt intake, 24-hour urine sampling is needed to confirm findings.

- There is no centralized up-to-date data source on the salt content of foods as there is no current requirement, at national levels or otherwise, for disclosure.
- In some cases LMIC lack the public sector capacities to uphold their roles in either voluntary reformulation agreements or other aspects considered important to reducing dietary salt intake. For example, if the food industry makes commitments to reach specific salt content targets, and if food-labeling requirements are regulated or even if voluntary, government agencies with the authority to evaluate industry claims on labels and analyze food products need specific and sometimes new institutional capacity to accommodate the tasks.
- In a number of countries, there is uncertainty as to how and where to start population level dietary salt reduction or while plans may be under development, there is uncertainty as to how to coordinate salt reduction with iodine fortification. Some medical sectors advocating for prevention of iodine deficiency disorders are reluctant to accept salt reduction policies.
- The overconsumption of salt is still in some cases not recognized as a public health issue or is recognized but not a priority for action. There is a low overall perception of risk by the general public and health professionals and a low level of recognition as to the role of the food industry and its corporate social responsibility towards dietary salt reduction given that the salt content of processed food products is a food safety issue. And there are competing important nutrition priorities including those related to obesity.
- Having adopted a voluntary approach for food reformulations, industry is proposing conservative goals and the negotiation processes are prolonged. Questions are arising as to whether voluntary agreements are strong enough instruments to have measurable population level impacts in a timely manner.
- Undertaking to standardize regulations e.g. for nutrition labeling across a common sub-region-

al market requires negotiation and consensus building.

- Communicating salt content reductions in food to consumers requires careful and coordinated messaging between the public health sector and the food industry.
- There are difficulties with the transfer of reformulation technology from large to small companies.
- Building and maintaining a multi-stakeholder and multi-sector approach requires sustained commitment on the part of the public health sector.
- In many countries, the technical capacity to evaluate the cost-benefit, cost-effectiveness and/or the cost-savings from reduced dietary salt is lacking.
- In some cases, the food industry has complained about the requirement to modify the original content of their food products, arguing that they lack the necessary infrastructure to reformulate and that there is no enough time allowed to comply with new requirements.
- Where the general population has not been sufficiently sensitized to the issue, in particular the role and responsibilities of the food industry, there has been opposition to reformulation requirements, with the public preferring personal and/or parental choice in food selection.
- For LMIC countries that have not yet launched dietary salt reduction initiatives and are considering where to start, the experiences with bread in the southern cone countries provide lessons and direction. They are important examples of how national associations of artisanal bread makers and of supermarkets with private labels can be mobilized. And there are health economic analyses from the region to support taking action on bread.
- The range of target and timeline commitments made by the multinational and large national food companies in the southern cone countries and to the US NSRI are evidence of what can be achieved, if not of the specific product reformulations or new low/no salt product availability, then of the technical feasibility of reducing the salt content of common food categories that contribute significant amounts of salt to the diet.
- Several of the challenges identified by countries, whether active and not yet, can be addressed through the dissemination and facilitated implementation of the tools and resources collected and developed by the Expert Group e.g. how to determine the main sources of salt in the diet or how to conduct health economic analyses.
- The national experiences with and tools to disseminate information and raise awareness among the public and communities of health professionals are potentially of great value in countries where public health initiatives are under consideration. They supply evidence to policy makers who need to apply pressure for the issue to be recognized and build up the broad public support necessary to secure the issue on political agendas in the midst of competing priorities.
- Research partnerships have and are evolving e.g. between IDD prevention and salt reduction programmes, increasing the potential to respond to calls for and access research grants.

Momentum and Potential in the Region

- The concentration of momentum in the southern cone countries in Latin America – Argentina, Brazil and Chile – suggests the potential for action to spread to other countries, especially to those in the MERCOSUR common market. In general, common markets in the region can facilitate the distribution of reformulated foods from countries where salt reduction policies are being implemented and may contribute to spreading market demand.

References

- 1 Eaton SB, Konner M.** Paleolithic nutrition. A consideration of its nature and current implications. *N Engl J Med.* 1985;312:283-9.
- 2 Mattes RD, Donnelly D.** Relative contributions of dietary sodium sources. *Am J Clin Nutr.* 1991;10:383-93.
- 3 Scientific Advisory Committee on Nutrition (UK).** Salt and Health. Norwich, England: The Stationary Office; 2003. Accessed August 2011 at http://tna.europarchive.org/20090810121540/http://www.sacn.gov.uk/pdfs/sacn_salt_final.pdf.
- 4 Brown IJ, Tzoulaki I, Candeias V, Elliott P.** Salt intakes around the world: implications for public health. *Int J Epidemiol* 2009; 38(3):791-813.
- 5 World Health Organization.** Reducing Salt Intake in Populations: Report of a WHO Forum and Technical Meeting 5-7 October, 2006 Paris, France. Geneva, Switzerland: World Health Organization; 2007.
- 6 World Health Organization.** Creating an enabling environment for population-based salt reduction strategies. Report of a joint technical meeting held by WHO and the Food Standards Agency, United Kingdom, July 2010 2011; 1(1):3-42.
- 7 Sarno F, Claro RM, Levy RB, Bandoni DH, Ferreira SRG, Monteiro CA.** Estimated sodium intake by the Brazilian population, 2002-2003. *Rev Saúde Pública.* 2009;43:219-25.
- 8 Encuesta nacional de nutrición y salud 2004-05.** Accessed June 2011 at <http://www.msal.gov.ar/html/Site/ennys/download/Implementaci%C3%B3n.pdf>.
- 9 Encuesta nacional de salud ENS Chile 2009-2010.** Accessed October 2011 at http://www.minsal.gob.cl/portal/docs/page/minsalcl/g_home/submenu_portada_2011/ens2010.pdf
- 10 Committee on Strategies to Reduce Sodium Intake, Food and Nutrition Board, Institute of Medicine (US); Henny JE, Taylor CL, Boon CS, Editors.** Strategies to Reduce Sodium Intake in the United States. Washington, DC: The National Academies Press; 2010.
- 11 Sodium Working Group.** Sodium Reduction Strategy for Canada, Recommendations of the Sodium Working Group. Ottawa, Canada: Health Canada; 2010. Accessed June 2011 at http://www.hc-sc.gc.ca/fn-an/alt_formats/pdf/nutrition/sodium/strategy/index-eng.pdf
- 12 He FJ, MacGregor GA.** Salt reduction lowers cardiovascular risk: meta-analysis of outcome trials. *The Lancet.* 2011;378:380-2.
- 13 Campbell NRC, Cappuccio FP, Tobe SW.** Unnecessary controversy regarding dietary sodium: a lot about a little. *Can J Cardiol.* 2011;27:404-6.
- 14 World Health Organization.** Global health risks: Mortality and burden of disease attributable to selected major risks. Geneva, Switzerland: WHO; 2009.
- 15 He FJ, MacGregor GA.** Effect of modest salt reduction on blood pressure: a meta-analysis of randomized trials. Implications for public health. *J Hum Hypertens* 2002;16:761-70.
- 16 He FJ, MacGregor GA.** Importance of Salt in Determining Blood Pressure in Children. Meta-analysis of Randomized Controlled Trials. *Hypertension* 2006;48:861-9.
- 17 He FJ, MacGregor GA.** Effect of longer-term modest salt reduction on blood pressure. *The Cochrane Database of Systematic Reviews* 2004;(1):1-64.
- 18 Joffres M, Campbell NRC, Manns B, Tu K.** Estimate of the benefits of a population-based reduction in dietary sodium additives on hypertension and its related health care costs in Canada. *Can J Cardiol* 2007; 23(6):437-443.
- 19 Committee on Public Health Priorities to Reduce and Control Hypertension in the U.S.** Population (IoM). A Population-Based Policy and Systems Change Approach to Prevent and Control Hypertension. Washington DC: National Academy of Sciences; 2011.
- 20 Departamento de Epidemiología Ministerio de Salud.** Chilean Health Report. Gobierno De Chile, Ministerio De Salud; 2003.
- 21 Danaei G, Ding EL, Mozaffarian D, Taylor B, Rehm J, Murray CJ et al.** The preventable causes of death in the United States: comparative risk assessment of dietary, lifestyle, and metabolic risk factors. *PLoS Med.* 2009;6:e1000058.
- 22 Panel on Dietary Reference Intakes for Electrolytes and Water, Standing Committee on the Scientific Evaluation of Dietary Reference Intakes.** Dietary Reference Intakes for Water, Potassium, Sodium, Chloride and Sulfate. Scientific Evaluation of Dietary Reference. Washington, DC: National Academies Press; 2004.
- 23 He FJ, Marrero NM, MacGregor GA.** Salt intake is related to soft drink consumption in children and adolescents: a link to obesity? *Hypertension* 2008;51:629-34.
- 24 He FJ, MacGregor GA.** A comprehensive review on salt and health and current experience of worldwide salt reduction programmes. *J Hum Hypertens* 2009;23:363-84.
- 25 Gaziano TA, Bitton A, Anand S, Weinstein MC for the International Society of Hypertension.** The global cost of nonoptimal blood pressure. *J Hypertens.* 2009;27:1472-7.
- 26 Asaria P, Chisholm D, Mathers C, Ezzati M, Beaglehole R.**

- Chronic disease prevention: health effects and financial costs of strategies to reduce salt intake and control tobacco use. *The Lancet*. 2007;370:2044-53.
- 27 **Cobiac LJ, Vos T, Veerman JL.** Cost-effectiveness of interventions to reduce dietary salt intake. *Heart* 2010; 96(23):1920-1925.
 - 28 **Bibbins-Domingo K, Chertow GM, Coxson PG, Moran A, Lightwood JM, Pletcher MJ et al.** Projected Effect of Dietary Salt Reductions on Future Cardiovascular Disease. *N Engl J Med* 2010;362:590-9.
 - 29 **Meneton P, Jeunemaitre X, de Wardener HE, MacGregor GA.** Links between dietary salt intake, renal salt handling, blood pressure, and cardiovascular diseases. *Physiol Rev* 2005; 85(2):679-715.
 - 30 **He FJ, MacGregor GA.** Reducing population salt intake worldwide: from evidence to implementation. *Prog Cardiovasc Dis*. 2010;52:363-82.
 - 31 **World Health Organization.** Creating an enabling environment for population-based salt reduction strategies. Report of a joint technical meeting held by WHO and the Food Standards Agency, United Kingdom, July 2010. Geneva: World Health Organization, 2010. Accessed July 2011 at http://whqlibdoc.who.int/publications/2010/9789241500777_eng.pdf.
 - 32 **World Health Organization.** Strategies to monitor and evaluate population sodium consumption and sources of sodium in the diet. Report of a joint technical meeting convened by WHO and the Government of Canada, October 2010. Geneva: World Health Organization, 2011. Accessed July 2011 at http://whqlibdoc.who.int/publications/2011/9789241501699_eng.pdf.
 - 33 **New York City Department of Health and Mental Hygiene.** Sodium study confirms that New Yorkers eat too much salt. Accessed August 2011 at <http://www.nyc.gov/html/doh/html/pr2011/pr005-11.shtml>
 - 34 **Tabla de Composición de Alimentos de América Latina.** Accessed October 2011 at <http://www.rlc.fao.org/es/bases/alimento/default.htm>
 - 35 **Blanco-Metzler A, Montero-Campos M, Chan V, Campbell N.** Survey on data of sodium in processed and prepared foods of Latin America. 9th International Food Data Conference. Norwich UK, September 2011.
 - 36 **Rubinstein A, Colantonio L, Bardach A, Caporale J, et al.** Estimation of the burden of cardiovascular disease attributable to modifiable risk factors and cost-effectiveness analysis of preventative interventions to reduce this burden in Argentina. *BMC Public Health*. 2010;10:627 doi:10.1186/1471-2458-10-627.
 - 37 **Joffres M, Alimadad A.** Effect of sodium reduction on cardiovascular disease (Latin American countries). Faculty of Health Sciences, Simon Fraser University, British Columbia. Unpublished paper.
 - 38 **Campbell NRC, Willis KJ, L'Abbé M, Strang R, Young E.** Canadian Initiatives to Prevent Hypertension by Reducing Dietary Sodium. *Nutrients*. 2011;3:756-64.
 - 39 **Oficialización del Plan de Reducción del Consumo de Sal/Sodio de Costa Rica.** Accessed October 2011 at <http://portal.campusvirtualsp.org/virtualcampus/costarica/drupal/?q=node/64>
 - 40 **Plan de Salud de Centroamérica y República Dominicana 2010 – 2015.** Accessed October 2011 at <http://www.sica.int/busqueda/Noticias.aspx?IDItem=45368&IDCat=3&IdEnt=143&Idm=1&IdmStyle=1>
 - 41 **Ferrante D, Apro N, Ferreira V, Virgolini M, Aguilar V, Sosa M, et al.** Feasibility of salt reduction in processed foods in Argentina. *Rev Panam Salud Publica*. 2011;29:69–75.

Appendix 1: Organizations that Have Endorsed the Policy Statement as of September 2011

ORGANIZATION	DATE OF ENDORSEMENT
1. Blood Pressure Canada	November 2009
2. InterAmerican Heart Foundation	November 2009
3. Canadian Stroke Network	November 2009
4. Canadian Diabetes Association	November 2009
5. Canadian Heart Failure Network	November 2009
6. Canadian Hypertension Society	November 2009
7. Canadian Medical Association	December 2009
8. Healthy Caribbean Coalition	February 2010
9. Caribbean Cardiac Society	February 2010
10. World Hypertension League	March 2010
11. Canadian Public Health Association	March 2010
12. Sociedad Ecuatoriana de Cardiología	April 2010
13. PROTESTE-Associação Brasileira de Defesa do Consumidor	April 2010
14. International Diabetes Federation	May 2010
15. Heart Foundation of Jamaica	May 2010
16. Instituto Brasileiro de Defesa de Consumidor	May 2010
17. Asociación Argentina de Dietistas y Nutricionistas Dietistas	May 2010
18. Sociedad Argentina de Nutrición	May 2010
19. American Heart Association	May 2010
20. International Federation of Kidney Foundations	May 2010
21. Asociación Solidaria de Insuficientes Renales	April 2010
22. Colegio de Médicos y Cirujanos de Costa Rica	April 2010
23. Sociedade Brasileira de Medicina de Família e Comunidade	April 2010
24. Sociedad Chilena de Hipertensión	May 2010
25. Ministerio de Salud de Chile	August 2010
26. Ministerio de Salud Venezuela	August 2010
27. Ministry of Health Suriname	August 2010
28. Ministerio de Salud de Uruguay	November 2010
29. Ministerio de Salud de Argentina	September 2011

30. Ministério da Saúde Brasil	confirmed pending letter
31. Ministerio de Salud de Costa Rica	January 2011
32. Asociación Latinoamericana de Diabetes	March 2011
33. Sociedad Latinoamericana de Nefrología e Hipertensión	March 2011
34. Fundación InterAmericana del Corazón Argentina	March 2011
35. Federación Nacional de Operadores de Mercados Fruti hortícolas de la República Argentina	March 2011
36. Federation Argentina de Cardiología	March 2011
37. Sociedad Latinoamericana de Arteriosclerosis	March 2011
38. ForoSalud, Red Peruana de Pacientes, Enlace Red Andina de Pacientes	March 2011
39. Centro de Investigación Epidemia del Tabaco, Framework Convention Alliance	March 2011
40. Asociación Lupus Chaco – Pacientes Online	March 2011
41. Dirección de Salud Mental	March 2011
42. Fundación para Prevención y Control de Enfermedades Crónicas No-Transmisibles para América Latina	March 2011
43. Sociedad Mexicana de Salud Pública / International Union Against Tuberculosis and Lung Disease	March 2011
44. Alianza Internacional al Pacientes	March 2011
45. Union Antitabaquica Argentina	March 2011
46. Del Plata Adventist University, School of Health Sciences	March 2011
47. International Federation Medical Students Association	March 2011
48. Red Familiar, Usarios y Voluntarios	March 2011
49. Sociedad Sudamericana de Cardiología	March 2011
50. Asociación Latinoamericana de Tórax	March 2011
51. American Cancer Society	March 2011
52. Fundacion Cardiologica Correntina	March 2011
53. Unión de Usuarios y Consumidores, Filial Rosario	March 2011
54. LATINFOODS (Latin American Network of Food Data Systems)	July 2011
55. Ministry of Social Protection, Vice Ministry of Public Health, Colombia	September 2011
56. Consumers International South America	March 2011

Appendix 2: Dissemination Activities

Articles

Campbell N, Dary O, Cappuccio FP, Neufeld L, Harding K, Zimmermann MB. A call for action to coordinate programs to improve global health by optimizing salt and iodine intake. Accepted for publication in the World Health Organization Bulletin.

Legetic B, Campbell N. Reducing salt intake in the Americas: Pan American Health Organization actions. J Health Communication. 2011;16:37-48.

Campbell N, Correa-Rotter R, Neal B, Cappuccio FP. New evidence relating to the health impact of reducing salt intake. Nutrition, Metabolism and Cardiovascular Diseases. 2011;21:617-9.

Campbell NRC, Legowski B, Legetic B. Mobilizing the Americas for dietary salt reduction. The Lancet. 2010;377:793-5.

Blanco-Metzler A, Legetic B, Campbell NRC. Los países de las Américas se movilizan para disminuir la hipertensión y las ECV mediante la reducción del consumo de sal en la población. Archivos Latinoamericanos de Nutrición. 2010;60:Artículo No.1.

Campbell NRC, Legowski B, Legetic B, Wilks R, Pinto de Almeida Vasconcellos AB. A new initiative to prevent cardiovascular disease in the Americas by reducing dietary salt. CVD Prevention and Control. 2009;4:185-7.

Campbell NRC, Legowski B, Legetic B, Wilks R, Pinto de Almeida Vasconcellos AB, on behalf of the PAHO/WHO Regional Expert Group on Cardiovascular Disease Prevention through Dietary Salt Reduction. PAHO/WHO Regional Expert Group Policy Statement – Preventing cardiovascular disease in the Americas by reducing dietary salt intake population-wide. CVD Prevention and Control. 2009;4:189-91.

Presentations

Norm Campbell – October 2009, World Hypertension Conference in Beijing, China.

Norm Campbell, Branka Legetic, Ricardo Uauy – November 2009, Congress of the Latin American Society for Nutrition (SLAN) in Santiago, Chile.

Branka Legetic – June 2010, World Cardiology Congress in Beijing, China.

Norm Campbell – September 2010, scientific meeting of the International Hypertension Society in Vancouver, Canada.

Ricardo Correa-Rotter – December 2010, by invitation from the Ministry of Health of Uruguay, Uruguayan Cardiology Association, Hypertension Society and Academy of Medicine, and the PAHO office in Uruguay.

Branka Legetic – September 2010, the II World Congress on Public Health Nutrition and the European Salt Action Network in Portugal.

Branka Legetic, Norm Campbell, Kirsten Bibbins-Domingo, Ricardo Correa-Rotter – March 2011, symposium on salt reduction as a cost-effective method for prevention of cardiovascular disease, 14th Congress on the Investigation in Public Health, Mexico.

Norm Campbell, Ricardo Correa-Rotter, Sonia Angell and Adriana Blanco-Metzler – March 2011-Participación en representación de la Iniciativa de la OPS en la Reunión Técnica de los Participantes en la Consulta Regional de Alto Nivel de las Américas contra las Enfermedades Crónicas NO Transmisibles (ECNT) y la Obesidad, Mexico.

Adriana Blanco-Metzler, Branka Legetic, Norm Campbell – September 2011, the PAHO Initiative on Cardiovascular Disease Prevention through Dietary Salt Reduction, at the 9th International Food Data Conference in Norwich UK.

Adriana Blanco-Metzler, M.A Montero-Campos, Victoria Chan, Norm Campbell – September 2011, Survey on Data of Sodium in Processed and Prepared Foods of Latin America, at the 9th International Food Data Conference in Norwich UK.

Adriana Blanco-Metzler – September 2010, Iniciativa de la OPS: Prevención de las ECV en las Américas mediante la reducción de la ingesta de la sal alimentaria en toda la población at the Workshop for the validation of the “Plan Nacional de Reducción del Consumo de Sal/Sodio en Costa Rica” in Costa Rica.

Adriana Blanco-Metzler – March, 2010, Iniciativa de la OPS: Prevención de las ECV en las Américas mediante la reducción de la ingesta de la sal alimentaria en toda la población/proyecto investigación ITCR-INCIENSA. Expert Panel of Research in Foods and Nutrition, Curso Metodología de la Investigación I. Nutrition Scholl, University of Costa Rica.

Branka Legetic, Adriana Blanco-Metzler – June 2009, Movilización sobre Reducción del consumo de sal en las Américas Workshop de la Red LATINFOODS: Iniciativa regional sobre salud cardiovascular OPS-OMS. LATIN-FOODS Workshop in Argentina.

Appendix 3: Country-specific Targets and Timelines

Argentina / Campaña “Menos Sal, Mas Vida”, accessed August 2011 at http://www.msal.gov.ar/hm/Site/noticias_plantilla.asp?id=274

Brazil / Orientações para redução do consumo de sódio, accessed August 2011 at <http://nutricao.saude.gov.br/sodio.php>


Canada / Draft sodium reduction targets and label data for prepackaged foods, accessed August 2011 at <http://www.hc-sc.gc.ca/fn-an/consult/2011-sodium/append-a-eng.php>

Chile / Estrategia de Reducción de SAL/SODIO en los Alimentos, accessed August 2011 at http://www.redsalud.gov.cl/portal/url/page/minsalcl/g_proteccion/g_alimentos/reduccion_sodio.html

Mexico / Lineamientos técnicos para el expendio o distribución de alimentos o bebidas en los establecimientos de educación básica, accessed August 2011 at <http://www.insp.mx/alimentosescolares/index.php>

US NSRI

- Targets for packaged food, accessed August 2011 at <http://www.nyc.gov/html/doh/html/cardio/cardio-salt-initiative-packagedfood.shtml> and <http://www.nyc.gov/html/doh/downloads/pdf/cardio/cardio-salt-nsri-packaged.pdf>
- Targets for restaurant food, accessed August 2011 at <http://www.nyc.gov/html/doh/html/cardio/cardio-salt-initiative-restaurantfood.shtml>
- Corporate Commitments and Comments, accessed August 2011 at <http://www.nyc.gov/html/doh/downloads/pdf/cardio/cardio-salt-nsri-commitments.pdf>





**Population
Awareness-Building**

Applying a Social Marketing Framework to Salt Reduction

A Social Marketing Framework for Salt Reduction

In 2008, approximately one billion adults worldwide had uncontrolled hypertension with almost a third of all cases attributed to excess dietary sodium. (Campbell et al, 2012). Almost every nation in the world has an average population salt intake above what is required for optimum health. In developed countries, processed and catered foods contribute most of the salt in the diet with the rest added at the table or during cooking. In developing countries, especially in rural areas, most dietary sodium comes from salt added during cooking. For most countries, implementing a national salt reduction program is one of simplest and most cost-effective ways of improving public health.

Since 2007, the World Health Organization (WHO) has been supporting the development of national salt reduction strategies by establishing networks in partnership with regional organizations around the world. In September 2009, the Pan American Health Organization (PAHO) established an expert group (EG) to examine excess dietary salt as a health risk in the Americas, issue evidence-informed policy recommendations for salt reduction in the region, and develop tools and resources to support regional dietary salt reduction (Campbell et al., 2009; PAHO, 2009.) The following social marketing framework, an evidence-based foundation for national, regional, and local social marketing strategies, is one of those resources.

Social marketing is defined by Kotler and Andreasen as the application of commercial marketing principles to influence social behaviors. Marketing principles key to social marketing include: ⁽¹⁾ A Consumer Focus; ⁽²⁾ Audience Segmentation; ⁽³⁾ The Exchange Theory (including an analysis of the 'competition'); and ⁽⁴⁾ The Marketing Mix. These principles will be discussed in detail throughout the framework, with the goal of enabling the reader to apply a social marketing framework to the design of salt reduction initiatives in their country or region.

According to Webster et al (2011), no country is likely to achieve a significant decrease in population salt consumption when a program is restricted to consumer education alone. For maximum impact, national salt reduction efforts must include changes to the environment that make it easier for the population to consume less salt. The framework can be used to design interventions that decrease demand for salt and high sodium products, generate demand for low sodium or salt alternatives, and aim at influencing the policy environment. With the goal of moving beyond public awareness to actually changing behavior, designing programs using a social marketing framework, will link supply and demand side activities, and allow us to integrate best

practices from salt reduction programs gathered since the 1960's.

Following is an outline for a SOCIAL MARKETING STRATEGY. Each component of the strategy includes outputs and a step by step process. Planning tools for select components can be found in the APPENDIX.

Situation Analysis

OUTPUT: A snapshot situation analysis and description of the communication environment as it relates to salt reduction.

The situation analysis allows you to plan your social marketing intervention within a larger context. It will ensure that your strategy complements what is currently going on in the current environment and builds upon lessons learned from previous salt reduction communication, marketing, advocacy, and public education efforts. This section should include a situation analysis and a description of the national, regional, or local communication environment.

To develop a SITUATION ANALYSIS, follow these steps:

STEP A. Describe the Background

You may use these questions as a guide to describing the background and context of hypertension, CVD, and salt/sodium consumption in your country:

- What is the prevalence of hypertension/CVD in your country/region?
- Who is most at risk?
- What are some of the drivers of salt consumption (e.g. processed foods, food preparation, absence of risk perception?).
- What are the current levels of sodium and salt consumption?
- What are country policies relevant to salt reduction? (e.g. food labeling).

STEP B. Describe Previous/Current Salt Reduction Communication, Marketing, Advocacy, and Public Education Efforts

Talk with partners, including local or national media, to answer the following questions:

- What previous efforts have been made to reduce salt consumption in your country? (Describe previous salt reduction interventions).
- Were they evaluated and, if so, what lessons have been learned that could inform future planning?
- What, if any, current salt reduction initiatives are taking place? (This may include mass media, community-based, and or policy advocacy efforts).

STEP C. Do a Gap Analysis

Now that you are aware of the current communication

environment, what interventions could you design that would complement existing communication efforts? For example, if a certain population has not yet been targeted in salt reduction efforts, (e.g. patients with high blood pressure) this may be an ideal time to develop an intervention for them. The reason for a gap analysis is so that you do not duplicate efforts or use valuable resources to reinvent the wheel.

Literature Review and Research Findings

OUTPUT: Key findings from a review of the literature and secondary research on salt reduction interventions, including any documented best practices identified in salt reduction. Note that most previous salt reduction strategies have been led by government organizations, involve industry-led food reformulation of food, have clear daily intake targets, and strong communication strategies aimed at changing consumer behavior. Five countries, including Finland, France, Ireland, Japan, and the UK have demonstrated some impact as a result of their salt reduction initiatives. In four cases, this included evidence of changes in population salt consumption; in another four, changes in the salt levels in food; and in two, changes in consumer awareness. (2011, Webster et al) SEE SALT REDUCTION LITERATURE REVIEW SYNTHESIS IN THE APPENDIX.

Develop this section by following these steps:

STEP A: Conduct a Literature/Secondary Research Review

Analyze all secondary research as it pertains to salt reduction in your country or region. Focus on identifying baseline data regarding target audience knowledge, attitudes, beliefs, and behaviors. In addition, identify demand drivers and barriers to salt reduction. Barriers may be both external/environmental (e.g. policy, fast food popularity) as well as internal/behavioural (attitudes and practices.) If you have conducted your own primary research, synthesize key findings here.

Results of a recent baseline survey in several countries in the Americas revealed that although awareness of excessive salt as a health issue was high, risk perception and self-efficacy were low. (Claro et al, 2012)

STEP B: Assess Need and Plan for Additional Formative Research

Based on your synthesis of key findings from review of the literature and secondary research, determine what additional information or insights you may need to develop an effective social marketing strategy. For example, you may have collected sufficient quantitative data to help drive your strategy, but may be lacking in-depth insight about a particular population, including

their food purchasing, preparation, and consumption patterns. In 2010, the U.S. Institute of Medicine (IOM) Committee identified three areas that require additional research; among them, understanding how salty taste preferences develop throughout the lifespan, and understanding factors that impact consumer awareness and behavior relative to sodium reduction. (*Strategies to Reduce Sodium Intake in the United States, 2010*) It is only by understanding the obstacles (perceived and tangible) that the target audience faces that will enable us to influence their behavior.

It is also recommended that additional consumer research be conducted to pre-test creative concepts and/or materials with the target audience before finalizing a creative strategy or producing materials. Testing is described in more detail in Section VII-Step D of this document.

Target Audience

OUTPUT: Target audience selection, audience profile(s) and segmentation strategy.

Although you may want to educate ‘the public’ about the harm of excess sodium and cast a wide net promoting salt reduction, one size does not fit all. With a limited budget and a need to demonstrate results, it is crucial to select a more homogeneous group or you risk diluting social marketing efforts. International best practices direct us to a more consumer-focused targeted approach in order to impact behavior.

Target audience identification should not be based on epidemiology alone; i.e., who is most at risk, but should include additional considerations. If you need to show more immediate results, you may want to select a population that is also most open and ready to change. A social marketing strategy tailored to a specific target group will be more targeted, more relevant, and in the end, have a better chance of reaching intended goals.

Target audience selection/identification can be based on any number of variables, including the following:

- Who is most open (ready) to change? (*Trans-theoretical Model Stages of Change, Prochaska*).
- Who would benefit the most?
- Who feels most vulnerable?
- Who is easiest to reach?

To identify your target audience(s), follow these steps:

STEP A. Identify a **Primary Audience** based on the variables above.

This is the ‘bulls-eye’ target audience, the person whose behavior you hope to impact.

Keep in mind that, depending on the duration of your program, you may target more than one primary group; for example, female caregivers who make the primary

food purchasing and preparation decisions and people already diagnosed with hypertension. In this case, you may benefit by taking a phased approach where you begin targeting one of these groups and follow with another. Also, keep in mind that target groups may overlap; e.g. female caregivers with consistently high blood pressure.

STEP B. Develop a **target audience profile** that describes who this person is, what he or she values, the behaviors that put him/her at risk, and their current attitudes and behaviors related to salt consumption. Following is an example of a target audience profile for a female caregiver living in an unidentified urban location somewhere in the Americas.

LIZ

Liz is a 60 year old grandmother living in a large urban setting. She has always been a bit overweight, but she considers her health stable. Like most women, she takes care of others first and her own health comes last. Always known as the best cook in the family, Liz has started using more canned and processed foods for the sake of convenience. In fact, she felt a little guilty last holiday season to have used canned soup and packaged seasonings to prepare the family dinner instead of her more traditional recipes made from scratch.

When Liz and her husband eat alone at home, they automatically add salt to everything. In addition, Liz uses salt or other popular seasonings (e.g. Maggi) to prepare their meals. Liz and her husband are big bread lovers and they also enjoy visiting the grandchildren and taking them out for fast food.

Liz is not as active as she knows she should be; in fact, her lifestyle is fairly sedentary. She enjoys watching her favorite TV shows (on the food network), learning new recipes, talking to her grandchildren on the phone, and driving the 30 miles at least once a week to visit them.

Liz has heard that too much salt is bad for health, but doesn't know how much is too much. In addition, she has no idea how much sodium is in the fast food or canned and processed foods that she consumes. Liz hasn't had a physical exam in at least five years because she 'feels fine.' She has no idea what her blood pressure is or even should be. Once in a while, she gets headaches, but attributes that to the noises of the big city.

STEP C. Segment the primary audience based on the variables above or other variables specific to the population.

Once you have identified a population, you may want to **segment** the population further into even more homogeneous sub-groups for more targeted communication. For example, if you are targeting female caregivers, you

may want to segment using any number of variables, including the following:

- Socio-economic Status (SES).
- Heavy salt consumers (and/or consumers of processed foods).
- Urban versus rural.
- New mothers and caregivers of young children.
- Weight.

STEP D. Identify the **Secondary Audience**.

The secondary audience includes those people who have influence over the primary audience. Their input into program planning will ensure relevance and bring buy-in. They should be considered as key messengers in promotional efforts and should be recruited and trained as such. Common examples of influentials in salt reduction efforts include healthcare providers, dietitians, restaurant owners, street food vendors, retailers, and community leaders. If we return to the audience profile above, a secondary audience for Liz might include her daughter, her husband, or even grandchildren. A secondary audience for those diagnosed with hypertension might include their pharmacist as well as their healthcare provider.

You may also want to target and influence the behavior of other influentials, often considered a 'tertiary' audience, including the media, policymakers, educators, and food industry partners

Audience Analysis

OUTPUT: A synthesis of the primary target audience's current knowledge, attitudes, and behaviors related to salt consumption.

A common mistake is to assume that simple lack of awareness or knowledge is the primary barrier to individual salt reduction. As a result, you may develop a strategy which is information-driven and not audience-driven or consumer-centered. Communicating information to people who are already aware is not the best use of available time and resources.

As a case in point, results of a recent survey conducted in several Latin American countries revealed that almost 90% of those surveyed agreed that eating a diet high in salt can cause serious health issues – ranging from 76% in Ecuador to 98% in Argentina – but only 13% of participants believed they personally consumed too much salt. In this case, the gap between knowing and doing appears to be low risk perception. Even if motivated to change consumption patterns, the same survey revealed that only 26% of participants reported knowing the existence of a recommended maximum value of salt intake and only 7% were able to correctly identify this value. (Claro et al, 2012)

Results of recent message testing in Canada also revealed that there is virtually no awareness of how much sodium one currently consumes. In addition, knowledge of appropriate sodium levels and the most effective ways of reducing intake was very low. Based on research findings, document the following as it pertains to the primary target audience:

STEP A. Current knowledge, beliefs and behaviors—Describe the target audience’s current knowledge about safe sodium levels and monitoring sodium intake. Include any misperceptions. In addition, describe current attitudes and beliefs. For example, do they perceive themselves to be at risk for hypertension or other CVD’s? Does he or she feel confident in his/her ability to read and understand food labels correctly? Do they believe that salt reduction is worth the effort? In addition, include a description of the current behavior that puts them at risk. Describe their current use of salt and high sodium products.

STEP B. (Real and perceived) barriers—Describe the current barriers or potential obstacles to the target audience’s salt reduction efforts. This may include real (tangible) barriers such as lack of access to low sodium alternatives as well as perceived barriers such as lack of confidence or low risk perception. Also list any competitive behaviors (e.g. fast food consumption, adding salt at every meal, etc...) Competition may also include enjoyment and gratification; i.e. one’s personal taste and preference for salt which is learned over time and may even have an emotional association with family and tradition.

STEP C. (Real and perceived) benefits—Describe benefits to salt reduction that the target audience may find relevant. These are also referred to as demand drivers. Make sure to include both real/tangible benefits (e.g. Protection against hypertension and CVD) as well as perceived benefits (e.g. showing your family how much you care.)

Overview Of Supply Side

OUTPUT: A summary description of supply side interventions, including agreements with industry partners, progress on reformulation, sales and distribution systems, healthcare provider/retailer training, availability of low sodium and salt alternatives, product labeling, and point of purchase signage.

It is critical to have a functional and high-quality supply side in place before you begin to generate demand. This will include, but not be limited to collaboration/partnerships with the food industry, retailers who serve as im-

portant messengers and champions, a sufficient supply of salt substitutes and/or low sodium alternatives and an environment that facilitates access to information and low sodium options.

Best practices for salt reduction on the supply side include:

- Product reformulation –working with the food industry to reformulate foods and reduce sodium (voluntary or mandatory);
- Collaboration with the Food Industry—One example of successful voluntary collaboration is the partnership in France with the bakery sector to reduce sodium in their products;
- Regulation of the Food Industry—Population-based policies for sodium reduction such as those in Finland that have significantly decreased consumption;
- Food Labeling –consumer-friendly labeling such as warnings, traffic lights, percentage of daily intake (%DI), or healthy choice logos;
- Increased access and availability of low-sodium foods (e.g. Finland’s low sodium, potassium enriched Pansalt).

To document supply side progress and identify gaps, follow these steps:

STEP A. Review and document existing secondary audience (e.g. healthcare providers/retailers) training in interpersonal communication (IPC.) For example, have healthcare providers been adequately trained in delivering salt reduction messages? You may need to develop a new module, refresher training, or new materials/tools if you are designing a new intervention.

STEP B. Review and document national policies relating to salt and sodium reduction. This may include restrictions on advertising to children, mandatory reformulation, labeling, and/or taxation/subsidies. Identify gaps in policy and make recommendations for policy/media advocacy.

STEP C. Identify available salt alternatives or low sodium options. This should include consumer access to both in the marketplace and through restaurants/food vendors. Identify and map the placement of promotional signage (and labeling) where products are sold, specifically at point of sale locations such as supermarkets and restaurants. Identify locations where signage would further generate demand and build campaign recognition.

Demand Side/Communication Goal And Objectives

OUTPUT: Description of communication goal and objectives: In addition to an overarching demand side

communication goal, identify knowledge, belief, and behavioral objectives. You may want to make your objectives measurable (% change) and time-bound (by____.)

The following goals and objectives must be based upon the target audience's current knowledge levels, attitudes, and behaviors. (SEE SECTION II- RESEARCH FINDINGS AND SECTION IV- AUDIENCE ANALYSIS)

STEP A. Identify the **Communication Goal**. Your communication goal may be: To decrease demand for salt and high sodium products by (e.g. female caregivers) in food purchasing, preparation and consumption. You might also state it this way: To increase demand for more natural, less processed foods (with low sodium content and less salt)

STEP B. Identify **Knowledge Objectives** by answering this question: What do you want the target audience to *know* (that they don't already know) as a result of the communication? You may include several knowledge objectives; for example:

They will know:

- Which products are high(est) in sodium;
- The difference between salt and sodium;
- The signs and symptoms of hypertension;
- Where to access low-sodium products or salt substitutes;
- How to read a label;
- How to monitor their intake of salt/sodium;
- How to prepare foods/meals that deliver less sodium.

STEP C. Identify Belief Objectives by answering the question: What do you want the target audience to feel or believe (differently) as a result of the communication? Again, there may be more than one belief objective, including but not limited to objectives like the following:

They will believe:

- That they (and/or their families) are at risk for hypertension and CVD;
- In their ability to read labels, prepare foods with lower sodium content, monitor their sodium intake, and consume a diet lower in sodium;
- That purchasing and preparing foods lower in sodium will protect their loved ones;
- That women who take the initiative to lower the salt/sodium in their family's diet are wise and caring.

STEP D. Identify **Behavioral Objectives** by answering the question: What do you want the audience to do as a result of the communication? Be realistic in setting behavioral objectives, realizing that behavior does not change overnight. Even though a reduction in salt consumption may be the ultimate goal, because behavior change occurs along a continuum, your be-

havioral objective may be linked to an interim behavior; for example:

They will:

- Read labels every time they purchase a canned or packaged product;
- Ask about low sodium alternatives when they eat outside of their homes;
- Prepare foods using less added salt, salt substitutes, or low sodium alternatives;
- Monitor their (family's) daily intake of sodium;
- Talk to a healthcare professional about hypertension and heart health.

Following are tips (suggested behaviors) from the American Heart Association/American Stroke Association for reducing one's overall sodium intake:

- Read the nutrition facts label to compare and find foods that are lower in sodium.
- Choose fresh fruits and vegetables when possible.
- Limit the amount of processed foods you eat and limit your portion size.
- Avoid adding salt when cooking and/or eating, use fresh herbs instead.
- Specify how you want your food prepared when dining out. Ask for your dish to be prepared without salt.
- Try to choose foods with potassium. They counter the effects of sodium and may help lower your blood pressure.

As you look at the behaviors recommended above, put yourself in the shoes of the target audience you hope to reach. Do they know how to read labels? Do they have time to compare foods? Do they understand what is meant by 'processed' foods? Do they know what a portion size looks like? Do they consider ketchup and fries to be a fruit and vegetable? Do they have access to fresh herbs? Will their kids eat fresh herbs? Can you imagine them asking their fast food restaurant to prepare their food without salt? Do they know what foods contain potassium or what that is? Although all of these tips reflect sound public health advice, we need to make sure that we are promoting behaviors that can be clearly understood, performed, and reported.

Marketing Mix Strategies

OUTPUT: A Marketing Mix strategy which includes the 'four P's': PRODUCT, PRICE, PLACE, and PROMOTION strategies.

When one thinks about shifting consumer preference and increasing/decreasing demand, one normally thinks about the fourth P: promotion. Yet the success of any promotional effort is dependent upon effective product, price, and place strategies. Understanding what the target audience values, how they currently

perceive a product; in this case, salt/sodium or foods high in sodium, and designing strategies that go beyond messages to facilitating behavior change are key to social marketing.

In social marketing, a PRODUCT strategy can refer to a tangible product (e.g. salt substitute,) a service (e.g. blood pressure screening,) or a behavior (e.g. monitoring daily sodium consumption.) The PRICE strategy looks at both monetary and non-monetary (e.g. denial, lack of self-confidence, misperceptions) costs. The PLACE strategy refers to the supply side, where the product or service is accessed, in addition to where the behavior is performed or choice is made. PROMOTION goes beyond messaging to promote the 'offering' (PRODUCT) and identify benefits that outweigh the barriers (PRICE) through multiple channels and cues to action (PLACE.) Developing a Marketing Mix Strategy (for all 4 P's) ensures that you have considered potential product innovation or improvement, supply side issues, and costs before designing your promotion. The MARKETING MIX Planning Tool will help guide you developing a marketing mix strategy. (SEE MARKETING MIX PLANNING TOOL IN APPENDIX.)

STEP A. Develop a Product Strategy

Start by developing a PRODUCT strategy based on the 'offering'. In this case, the offering may be alternatives to salt or low sodium and salt-free options. Products are normally designed to meet consumer needs, which explains the development of the new products based on consumer tastes and preferences. In addition, products are branded to make them unique and particularly attractive to certain segments of the population. These are all product strategies.

As you develop a PRODUCT strategy, think about any new products or product improvement that you could introduce. This might include developing a better tasting or healthier salt substitute or product. In Argentina, the Federation of Bakers' Industries has been partnering with government since 2006 to reformulate products to reduce salt content. As consumers in developed countries become more accustomed to high levels of sodium in processed and restaurant foods, it becomes increasingly difficult to introduce and increase demand for foods with healthier levels of sodium.

In social marketing, the PRODUCT also refers to the behavior that you are promoting. If, in fact, one of those behaviors is reading and understanding labels, you may consider designing a product that increases label literacy or helps 'translate' nutritional information into more consumer-friendly language. You may also be promoting self-monitoring; i.e. determining how much sodium one consumes in any given day. In that instance, you may want to work with a tech-savvy partner to design a

product/tool that facilitates self-monitoring (A 'saltometer' perhaps?)

Also integral to product strategy is package design which can either deter or promote usage; for example, a warning on high sodium products, or a healthy family displayed on low-sodium alternatives. The look, taste, and smell of a product also drive sales and consumption. All of these product 'benefits' must be considered when designing product strategies and later, in the promotion.

Labeling has been a key component in global salt reduction efforts. In fact, nutrition labeling has been mandatory on pre-packaged foods for many years. Labeling includes not only nutritional information on sodium content and percentage of Daily Intake (%DI), but also provides a visual identity that helps consumers identify 'heart-healthy' branded products. One example of product identity is Canada's "Health Check" program where products that meet sodium content criteria display a Heart Check logo. Sodium has been reduced in 150 food products to meet the Health Check criteria. (*Sodium Reduction in Canadian Food Products with the Health Check Program*) The American Heart Association has also recently announced an updated, new look to its nationally recognized Heart-Check mark, one of the components of the Heart-Check Food Certification Program. In addition, a heart-healthy logo is been displayed on select restaurant menu items across the United States to help consumers identify and select heart-healthy options. The food industry has hopped on the bandwagon with the Kellogg Co. recently unveiling its new "Heart Healthy Selection" logo on nine of its cereals.

POSITIONING is also a PRODUCT STRATEGY. Discover how the target audience currently perceives sodium and salt use (or processed foods) and think about how you can re-position it in their minds. For example, if you discover that people living with hypertension consider low sodium products to be a 'punishment', you may decide to reposition them as a gift. Instead of focusing on what they're giving up (salt/tradition/taste), focus on what they're getting (being a better mother/new exciting tastes/better health). You may also consider 'repositioning' foods highest in sodium so that consumers think twice about eating them for 'convenience-sake.' Although scare tactics rarely work in public health programs (unless you facilitate a solution at the same time,) positioning foods with high levels of sodium as silent killers might also be effective if executed 'tastefully.' Product strategies also include experiential marketing tactics; i.e. creating opportunities when and where the product can be 'experienced', such as product sampling and product demonstrations. This might include sampling events in supermarkets or street fairs and heart healthy cooking demonstrations.

In short, when developing a PRODUCT strategy for salt/sodium reduction, ask:

- What, if anything can I do to make the product (behavior) more attractive to the target audience?
- What can I do to make the packaging more attractive or easier to understand?
- Are there any new products that I can develop or introduce to facilitate sodium reduction? (e.g. salt substitute, reformulated products, ‘saltometer’)
- How can I ‘position’ or ‘reposition’ high salt use/ sodium consumption in the minds of the target audience to make it less attractive?

STEP B. Develop a Price Strategy

Now that you’ve designed the ‘offering’, (product innovation/positioning behavior) take a look at price. Your PRICE strategy will ensure affordability of salt substitutes and low sodium choices, but in addition, pricing strategies must consider all costs related to decreasing salt and sodium consumption, both tangible and intangible.

Adopting a new behavior or using a new product often comes at a price. It always includes an exchange, giving up one behavior for another and sometimes exchanging money for goods and services. Costs include both tangible monetary costs as well as intangible less direct costs, such as tradition (food preparation), low risk perception, or perceived loss of enjoyment. In monetary terms, asking a low income caregiver to feed her children more fruits and vegetables and less fast food would come at a price. Consider interventions that will decrease/lower the costs of or facilitate consuming less salt and sodium reduction (not only monetary costs, but emotional, psychological and time costs). Building in benefits and rewards that outweigh the costs or competition is referred to in marketing as the **exchange theory**. Think from your audience’s perspective. Don’t ask only what they **need**...Understand what they **want!** A PRICE strategy can also be policy-driven and refer to incentives and disincentives. One example is levying taxes on less healthy products (e.g. alcohol and tobacco) and providing subsidies on other products (e.g. corn, beef.) To date, no country has introduced economic measures to tax unhealthy foods/deter salt consumption and provide discounts or subsidies on healthier foods. That said, coupons issued by the food industry to generate demand and introduce new products are another common PRICE strategy that has been a best practice for decades.

In short, when developing a PRICE strategy,

- Identify costs; i.e. all barriers/obstacles/competition associated with monitoring and consuming/serving less salt and sodium.
- Ask how you can decrease the costs, monetary

and non-monetary, associated with salt reduction. (e.g. vouchers and coupons for free/reduced price samples)

- Ask how you can increase the (perceived) value of a reduced salt/ low sodium diet.
- A key element in developing a pricing strategy, often overlooked, is rewarding people in order to reinforce and sustain their behavior. Can you build in any rewards or reinforcement?

STEP C. Develop a Place Strategy

Think about where and when the audience performs the behavior or accesses the product. How can you make this place more convenient with easier access to the product? Examples include vending machines in and near schools that offer healthier (low sodium) snacks, displaying low sodium products or salt substitutes in a predominant way in the market, branding restaurant menus to identify low sodium choices or partnering with street vendors who serve heart-healthy low sodium foods. Also think about your “sales force” – the people who will take your program to the target audience. Consider the need for community-based health promoters, nutrition counselors, school cafeteria ‘champions’, and others who can make alternatives to salt and low sodium choices more attractive or accessible.

Place, oftentimes supply side strategies, are strategies that ensure access to products and convenience. In addition to product availability, place strategies include point of purchase (POP) branding, linking demand side promotional activities and materials with in-store or health facility-based communication activities. Place strategies also recognize the crucial role that providers and retailers play as key messengers in demand generation. Place strategies include training healthcare providers and retailers as ‘lo so’ salt reduction ambassadors, offering free delivery or mobile services that distribute new products and offer services (e.g. blood pressure screening,) and extending clinic or retailer hours for convenience. A place strategy might include working with a small neighborhood convenience store or family business to ensure availability of healthy low sodium choices.

Place strategies also include determining where the target audience thinks about or makes choices that impact their salt or sodium consumption and then determining how you can intervene at those moments of choice. This might include placing reminders or identifying healthier choices strategically in bakeries, in fast food restaurants, or posting point of sale materials in supermarkets. Another place strategy might be offering tastes of lower sodium recipes in street fairs, food stalls, or supermarkets.

Once again, policy often drives and supports a Place Strategy. For example, Chile has plans to prohibit the

sale of unhealthy foods, including those high in salt, in areas where children convene, including schools and children's events. Chile is also considering a ban on advertising of unhealthy foods to children. (Legowski and Legetic, 2011)

STEP D. Develop a Promotion Strategy

Now that you have designed product, price, and place strategies, you are ready to develop the **promotion** strategy. As part of your promotion strategy, be sure to promote the product, price and place features that you want the audience to know about. And follow these international best practices borrowed from commercial marketing:

D1. Begin with a Creative Brief (SEE CREATIVE BRIEF FORM IN APPENDIX) which will serve as a foundation and strategic direction for your promotional strategy. It will also serve as a consensus building tool, enabling input from key stakeholder. The Creative Brief allows you to identify and agree upon target audiences, synthesize target audience research, and translate findings into a creative strategy. It enables you to summarize audience insights, including barriers, benefits, and communication openings. In short, beginning the process with a **Creative Brief** ensures that your promotional strategy is audience research-driven and that all partners begin 'on the same page.'

As part of your Creative Brief, you will outline key messages and communication channels. Use audience research to determine the **communication channels** and activities that will best reach your audience. Whom do they listen to (and trust?) Where are they when they are making food purchasing decisions? (Consider in-store advertisements/ point of purchase materials) What channels do they access and where do they learn how to cook? (Think about radio, TV, Internet, and community-based channels.) What special promotional items would they use? (e.g., water bottles, refrigerator magnets, recipe cards) What special events might they attend? (e.g. concerts, health fairs, cooking demonstrations) How and when can you include the secondary audiences as influential key messengers?

In terms of communication channels, we know that mass media works well to influence social norms on a large scale, build brand recognition; i.e. popularize a brand or category, whereas interpersonal communication (IPC) is most effective in building skills and reinforcing product use. Interpersonal communication allows for feedback and questions, but messengers must be carefully trained. Healthcare professionals are considered valuable in promoting salt and sodium reduction and counseling patients about the risks of excessive consumption. Sodium reduction strategies should

be incorporated into a professional training curriculum that focuses on CVD prevention. (Mohan et al, 2009) On the supply side, retailers can also be key agents of behavior change as witnessed in many social marketing campaigns, including interventions that deter youth from purchasing tobacco products. In reaching harder to reach populations such as those living in rural areas or lower income populations with less access to healthy food choices, community health workers (*promotoras*) are a very popular and trusted channel. Community organizations and community leaders, considered influential as well, often need to be addressed as separate target audiences in advocacy strategies. When targeting children, who are known to have an influence on parent's purchasing decisions, school is another important communication channel, especially schools with a school lunch program or nutrition education curricula. Other popular communication channels include radio talk shows where target audience members can call in and ask questions, cooking shows or demonstrations where caregivers can learn new recipes, and social media such as mobile phone reminders and text message tips. A social networking sight where mothers and caregivers congregate might also be an effective way of engaging a higher income segment of the population.

D2. Develop Creative Concepts.

Before you go directly to 'messaging'; i.e. crafting information or telling people what they should do, it is useful to develop concepts or 'hooks' that will grab the audience's attention and help provide instant recognition for all campaign materials and activities. Concept development often requires striking a balance between expert recommendations and consumer perceptions and preferences.

Concepts, otherwise known as 'big ideas' may take the form of a theme, slogan, or a call to action, such as *Just do it* (Nike). Or they might take the form of a spokescharacter such as the Pillsbury Dough Boy who sells baked goods or Tony the Tiger who sells sweetened cereal to children. These examples represent creative strategies based on audience research and preference and are not simply random ideas. If you learn through your research, for example, that awareness of excessive salt consumption as a health issue is high, but knowledge of the sources of excess sodium is low (as was the case in Canada), you may decide to develop a creative concept that focuses on the hidden sources of sodium. If research reveals that high awareness does not translate into personal risk, you may want to develop a concept or creative strategy that increases risk perception and addresses that gap. Once again, in Canada, testing sodium reduction messages revealed that the most popular messages were also the most surprising; i.e.

those that warned of hidden sodium or the amount of sodium that children were consuming. Regardless of form, a concept will always include an element that can be repeated across every communication channel, branding the campaign and ensuring that one channel reinforces another.

Concept development is a talent, oftentimes best left to creative professionals. If you cannot afford the services of a full service advertising agency or one is not available, look for a local creative partner who has 'a way with words' or artistic talent. If you are purchasing these creative services and playing the role of consumer, use the BRAVO criteria (found in the APPENDIX) to evaluate the creative concepts presented to you.

Examples of creative concepts that have been designed to promote salt reduction include Argentina's *Less Salt. More Life. (Menos sal. Mas vida.)* Also, spokescharacters such as the UK's Sid the Slug or USDA's Power Panther have been created to counter popular food industry cartoon characters that sell processed foods to children. The marketplace is definitely ripe for big ideas that motivate consumers to think about salt, sodium, and processed foods in a whole new way.

D3. Test Creative Concepts.

Concept testing provides an important opportunity to gather more information about the target audience, specifically how they think and talk about salt or sodium and what, if anything might motivate them to reduce their salt or sodium intake. Testing three or four different concepts (creative approaches) with the target audience will serve as a catalyst for discussion that helps you to understand what ideas are most relevant, believable, and motivational. In addition, a wealth of audience verbatims, language that the audience uses when they speak to one another about salt or processed foods normally surfaces during concept testing. By using *their* language instead of expert language, you will have an easier time making sodium 'top of mind' and shifting tastes and social norms.

Concept testing is conducted most effectively in focus group discussions where participant's responses build upon one another and rich discussion can take place. There is not always a clear 'winner' or favorite concept; in fact, there are times when a new concept surfaces based on something that a participant suggests. Make sure to listen carefully and watch body language as participants react to each concept. And remember, sometimes the concept that makes them most uncomfortable leads to the most discussion and thought and ultimately, has the biggest impact on behavior change.

D4. Develop materials

Based upon results of concept testing, work with your creative partner(s) to develop materials. Refer back to

the Creative Brief to verify communication channels. If the target audience has low literacy levels or prefers getting their information through more verbal or visual sources, consider using audio or live demonstrations and, if affordable, video.

Although mass media works well to shift norms and popularize products, services, and even behaviors, you may not have the budget to produce TV spots. In that case, you may consider forging a partnership with the media to jointly develop and air public service announcements. In addition, partnering with the media will allow you to use 'earned' (versus paid) media, or public relations, placing newsworthy stories and featuring role models and spokespeople from the community. In addition, you may consider forging a partnership with an association or private sector advertiser (e.g. produce association or fruit juice manufacturer) to piggy back on their resources and promotional budget.

New media, with the ability to send targeted reminders and messages through SMS (mobile phone text messaging) is an efficient way to tailor messages to specific populations such as caregivers of school aged children. Developing materials with some degree of interactivity works well for two-way discussion and helps facilitators understand more about audience needs. Whenever possible, create a mechanism to elicit consumer-generated content such as real person stories, new recipes, and testimonials.

Often successful in shifting attitudes and influencing behavior are satisfied user testimonials, real people narratives with high emotional content. Understanding how this person overcame the obstacles and was rewarded as a result can often drive demand. Both behavioral science and marketing have provided a wealth of evidence that popularity of products drives demand.

The secondary audience, including restaurant owners, educators, healthcare providers, and retailers, plays a key role as campaign messengers. During this step, you will also develop materials for them that will help build their interpersonal communication (IPC) skills and ensure that the messages they communicate reflect those being heard, seen, or read through all other communication channels.

D5. Pre-test materials

Before producing final materials, it is always best to pre-test them with members of the target audience. You will need to work with your creative partner(s) to develop pre-test or prototype materials which are close to, but not yet, final. This is fairly simple to do for print and collateral materials, but a bit more difficult for radio/audio as the quality of the final production plays a large role in likeability and comprehension. If you are testing video, you can develop storyboards (frame by

frame visuals) with creative partners for pre-testing purposes.

This stage of pre-testing can be done through in-depth interviews (IDI) which are conducted one on one. Develop a pre-test interview guide with questions that help you determine whether the materials are relevant, understandable, likeable, and credible, and whether there are any problems which have been overlooked.

If you cannot afford to pre-test or don't have the time, you may combine concept testing (D3) with pre-testing by developing initial concepts that are closer to final materials (e.g. concept boards that include body copy/ text.)

D6. Produce Materials

You've developed concepts based on the Creative Brief. You've tested materials with the target audience. You've finalized the materials based on their input. Now it is time to produce materials. This means bringing in new partners, outside vendors such as printers, video production, audio production, multi-media production partners, or even organizations that convene the community or get hotlines up and running.

Keep in mind that, when it comes to print and collateral products, the more you print, the more economical, so anticipate your future needs now. In addition, you may want to leave space on materials for partners to add their logos or contact information. In addition, request a version of the artwork that can be easily reproduced in your own office setting so that you are able to create your own materials as needed, such as flyers.

If you cannot afford to contract a full service production company, you may want to capture some of your own images or record your own real people stories digitally. Digital stories, in either audio or video format are economical and can be used across communication channels, through mass media, played in clinic waiting room, used as a catalyst in group discussions, and at community-based events such as health fairs.

D7. Train key messengers

Before launching any promotional campaign, it is important to train key influentials as messengers and 'salt reduction ambassadors'. This can be done by sponsoring ambassador workshops for secondary audience members, inviting healthcare providers, community outreach workers, retailers, and the media. This will be the first 'launch' of the campaign, familiarizing these important messengers with the campaign strategy and theme, raising awareness about the important role that each one of them plays, and providing them with materials, including badges and point of service/sale signage. This will ensure that once the campaign is launched and materials disseminated to the target population, the people providing access to salt substitutes, low so-

dium products, and/or information, are well-versed in the promotional strategy and campaign messages.

D8. Disseminate materials

You will need to develop a dissemination plan. This will include taking a look at the communication channels selected based on audience preference and habits to determine when and how each channel is introduced. You may want to begin with a campaign launch, inviting the media to a community-based event. This will create excitement and 'buzz' around the campaign and get people talking (also known as 'word of mouth advertising' or viral marketing.)

When developing your dissemination plan, keep in mind that you will need a separate budget to purchase radio and television time, as well as newspaper, magazine, and billboard space. To save resources, you may want to consider leveraging community-based partners who have direct contact with and access to the primary target audience.

STEP E. Identify Gaps in Policy and Partnerships

Often considered important 'fifth P's' in social marketing, policy and partnerships are both critical to population-based change. Policy is crucial to reduced salt consumption, driving product strategies aimed at reformulating foods with lower sodium content, nutritional labeling, and regulating advertising to children. Partnerships with the food industry have also been key. As consumer demand for lower sodium alternatives is generated, products are available to meet those demands. Identify gaps in policy and consider interventions (e.g. media advocacy) and partners that will help you advocate for new policy measures. In addition, consider interventions for strengthening/reinforcing existing partnerships with industry and recruiting new, less traditional partners (e.g. street food vendors.)

Monitoring and Evaluation Plan

OUTPUT: An M & E Plan to monitor the implementation process and measure outcomes. Approximately 10% of your social marketing budget should be dedicated to M & E. Note that the effectiveness of countries with evidence of impact resulting from salt reduction initiatives was measured by population salt levels, salt content of foods, and/ or consumer knowledge.

STEPS: Answer the following questions to develop your M & E Plan.

- What techniques and methodologies will be used?
- When will measurements be taken?
- How will measurements be reported and to whom?
- How will M & E data be used to inform revisions?

Remember to refer back to the objectives (knowledge, belief, and behavioral) which you identified in your CREATIVE BRIEF and in STEP VI (GOALS AND OBJECTIVES) and make sure that your indicators reflect these objectives.

For ease in collection, you may want to link social marketing activities and outcomes back to existing country or program indicators if relevant indicators exist. Or you may want to develop your own indicators similar to the following examples:

PROCESS INDICATORS:

- Percentage of retailers displaying low sodium/salt substitute promotional posters, shelf danglers, and/or brochures;
- Number and percent of respondents who report having seen and/or heard the messages promoting salt/sodium reduction;
- Number of healthcare providers/community outreach workers trained in salt reduction communication.

OUTCOME INDICATORS:

- Percentage of caregivers who know the primary sources of sodium;
- Percentage of caregivers who know a place where they can get low sodium products or salt substitutes;
- Percentage of all respondents who, in response to prompted questions, say that a person can reduce their risk of CVD by reducing salt/sodium intake;
- Percentage of people reporting consistent reduction in salt and products high in sodium in the last year.

Since increase in demand must be met with an adequate supply, the following are offered as examples of useful supply side indicators:

SUPPLY SIDE INDICATORS:

- Total number of low sodium products available for distribution nationwide during the last 12 months;
- Number and percent of storage facilities that experienced a stock out at any given time period;
- Proportion of randomly selected retail outlets that have salt substitutes and/or low sodium options in stock at the time of a survey, of all retail outlets selected for survey.

Work Plan and Timeline

OUTPUT: A work/implementation plan and timeline for social marketing activities.

STEPS: Answer the following questions to develop your work plan and timeline:

- Will you roll out your program in phases? For example, if it includes a supply side intervention, will you

start there to ensure that supply can meet demand and that your offering is of quality?

- How will the phases be organized (i.e., by location/audience, objectives, activities?)
- For each phase, what will be done, who will be responsible, and when will it be done?

(SEE WORKPLAN TEMPLATE IN APPENDIX.)

Budget/Funding Sources

OUTPUT: National, Regional, or Local Social Marketing Budget

STEPS: Answer the following questions to develop your budget:

- What financial costs will be associated with the program? (Consider staff labor, outside costs such as advertising agency and research partner costs, materials development and production costs, media buying costs, etc...)
- If costs exceed available funds, what additional funding sources will be explored?
- What strategies will you use to appeal to potential donors?
- What potential cost-share or in-kind contributions can be made by partners; i.e. a media partner?

Section VII-STEP D (PROMOTIONAL STRATEGY) describes partners who are most relevant to promotional efforts. These partners add value by offering creative talents; an ability to ‘translate’ audience research into creative strategies; writing, design, and production skills; and a venue for disseminating messages and materials. **Cost-saving partnerships** include, but are not limited to:

- Local media partners can donate free time (radio and TV) and space (newspapers and magazines) for your promotional materials and include relevant characters, storylines, and products in their current programming.
- Local musicians and even celebrities are oftentimes happy to share their talents and skills by producing a hit song that targets the population you want to reach. Norms are created, reinforced, and shifted by popular culture. Remember that 1 ‘pop’ piece of music can go far with a music video played on TV and in restaurants, radio programming including talk shows with celebrities and ‘satisfied customers’, and live events and appearances.
- Working with real people who can share their stories and serve as role models. Remember that consumer-generated content (CGC) has become very popular and people are increasingly turning to the Internet to post and share stories, strategies, and recommendations with peers.

- Community-based partnerships, including partnerships with NGO's that serve the population you want to reach; clinics, pharmacies, and other non-traditional outlets that can post your promotional materials and provide access to low sodium and sodium alternatives.

Postscript: How Can Social Marketing Strengthen Current Efforts?

It is clear that governments have been proactive, thinking comprehensively about salt reduction for quite some time. Using the same terminology as private sector commercial marketing professionals, governments and NGO's are looking at both supply and demand. Supply side strategies have been developed through partnerships with the food industry and shifts in policy to increase availability and access to healthier choices. At the same time, public education efforts have been aimed at raising awareness and educating consumers in an attempt to increase demand for salt reduction and low sodium choices.

As you reviewed this framework, you may have observed that several social marketing principles are already reflected in current population-based efforts; i.e. product reformulation and labeling are a PRODUCT STRATEGY. Increased access to healthy alternatives and decreased access to high sodium products reflect a PLACE STRATEGY. Countries are also looking at PRICE STRATEGIES that provide incentives for healthier choices and deter consumers from high sodium products. Best practices in current salt reduction efforts are referenced throughout this framework, most of them focusing on the supply side.

Where social marketing can really make a difference is in helping us rethink demand side strategies. Taking a new look at current salt reduction efforts through a social marketing lens will help us strengthen efforts to influence individual choice, bridging the gap between what consumers know and what they actually do. Applying a social marketing framework to salt reduction will also help us tie together population-based and individual initiatives and create even more integrated and holistic programs.

In summary, here are select international best practices borrowed from social marketing that may help us achieve even greater impact in current salt reduction efforts:

1. *Identify one segment of the 'public' whose behavior you hope to influence*-Current efforts to target a broad general public only reach those who are motivated to listen. And even if they become more knowledgeable, they may never change their behavior. Start small. Be focused. Select a smaller homogeneous segment of the population and target your efforts for greater impact.
2. *Think like a consumer.* Understand what drives the target audience to consume as much salt and high sodium products as they do. Take a close look at the obstacles they face and an even closer look at the competition. Design interventions that address those drivers and knock out the competition. Make sodium reduction and consuming less salt as easy and convenient as fast food.
3. *What do they get in return?* Remember that they are giving up something in exchange for what you're offering. Ask yourself (or better yet, ask them) '*what's in it for them?*' Isolate a single and immediate benefit that will make it all worthwhile. Promoting prevention can't just be about what *won't* happen several years from now.
4. *Change doesn't occur overnight.*-Think about the multiple behaviors along an individual's road to salt reduction. *Read the label. Compare products. Monitor your daily sodium intake. Use less salt. Eat more produce and potassium.* Even though the list is long from a public health perspective, we need to make it more achievable for them. Select one behavior at a time, facilitate that behavior, and evaluate the outcome.
5. *Surprise them.* Evidence that surprising messages are the most memorable is not so surprising. Consumers are overwhelmed with public health and nutritional messages, many of them competing for mindshare and even contradictory. Our job is to help them see the same old thing in a whole new way. Changing behavior is voluntary and there's nothing like an 'ahah' moment to bring it on.

Appendix 1: Reviews, Outlines and Forms

- Global Literature Review Synthesis.
- Social Marketing Plan Outline.
- Marketing Mix Planning Tool.
- Creative Brief Form (Blank).
- Bravo Criteria Checklist.
- Workplan Template.
- Links to Additional Resources.

Global Literature Review Synthesis

Social Marketing Strategies For Salt/Sodium Reduction

Following is a summary of findings related to social marketing and related communication and public education strategies for salt/sodium reduction that come out of a global literature review.

Literature Review Findings

A review of the literature shows that there are numerous public education and behavior change strategies implemented by governments and health organizations to help people reduce their intake of excessive dietary salt, which is associated with increased blood pressure and other health risks. Successful interventions include both behavioral and population-based initiatives (Campbell et al., 2012). Notably, one article identified 32 different country salt reduction initiatives, and 28 “countries working with the food industry to reduce salt in foods. Ten had front-of-pack labeling schemes and 28 had consumer awareness or behavior change programs” (Webster et al., 2011). Since the 1970s, Finland has had a population-based policy for sodium reduction based on partnership and regulation of the food industry and consumer education via mass media (Mohan et al., 2009).

In the literature, common types of interventions include:

- Media campaigns.
- Regulations for standardized information of packaged food products.
- Internet-based lecture series (Campbell et al., 2010).
- Education and skills development aimed at improving the cardiovascular health of entire families (Johnson et al., 1995).
- Telehealth counseling (Nolan et al., 2011).
- School-based programs (Luepker et al., 1988; McKay et al., 1985).

In North America, Canada has implemented large, multi-sectoral interventions, such as the 2009 Canadian Hy-

pertension Education Program (CHEP) (Shi et al., 2011; Campbell et al., 2011). The CHEP includes “an extensive education program for the public and people with hypertension, a program to reduce dietary salt and a funded leadership position” (Campbell et al., 2010). In the United States, the Institute of Medicine produced recommended strategies for implementation by the food industry, government, and public outreach (2010). Other interventions have targeted specific populations, such as older adults, with diet-education materials to improve dietary status (Francis et al., 2009). To increase cardiovascular disease (CVD) prevention among African-American and low-income adults, the People Reducing Risk and Improving Strength through Exercise, Diet and Drug Adherence (PRAISEDD) was pilot-tested and showed significant decreases in blood pressure, but no changes in physical activity, sodium intake, medication adherence, or self-efficacy and outcome expectations (Resnick et al., 2009). From 2005 through 2008, Project HEART (Health Education Awareness Research Team) aimed to promote behavior changes to decrease CVD risk factors in a high-risk Hispanic border population in El Paso, Texas (Balcazar et al., 2010).

Interventions in South America also include education campaigns and/or regulations for standardized information on the sodium content of packaged food products. In Chile, the Ministry of Health and the food industry are working to gradually decrease the concentration of salt in bread (Valenzuela et al., 2011). Argentina is also working toward placing limitations on the salt content of food (Legowski et al., 2011).

Based on this initial literature review, best practices and lessons learned have been highlighted in the social marketing framework.

Resources

APPEL, L. J., CHAMPAGNE, C. M., HARSHA, D. W., COOPER, L. S., OBARZANEK, E., ELMER, P. J., ET AL. (2003). EFFECTS OF COMPREHENSIVE LIFESTYLE MODIFICATION ON BLOOD PRESSURE CONTROL: MAIN RESULTS OF THE PREMIER CLINICAL TRIAL. *JAMA*, 289(16), 2083-2093.

AYALA, C., TONG, X., VALDERRAMA, A., IVY, A., AND KEENAN, N. (2010). ACTIONS TAKEN TO REDUCE SODIUM INTAKE AMONG ADULTS WITH SELF-REPORTED HYPERTENSION: HEALTH-STYLE SURVEY, 2005 AND 2008. *J CLIN HYPERTENS (GREENWICH)*, 12(10), 793-799.

BALCAZAR, H. G., DE HEER, H., ROSENTHAL, L., AGUIRRE, M., FLORES, L., PUENTES, F. A., ET AL. (2010). A PROMOTORES DE SALUD INTERVENTION TO REDUCE CARDIOVASCULAR DISEASE RISK IN A HIGH-RISK HISPANIC BORDER POPULATION, 2005-2008. *PREV CHRONIC DIS*, 7(2), A28.

CAMPBELL, N. R. (2007). CANADA CHAIR IN HYPERTENSION PREVENTION AND CONTROL: A PILOT PROJECT. *CAN J CARDIOL*, 23(7), 557-560.

CAMPBELL, N. R., JOHNSON, J. A., AND CAMPBELL, T. S. (2012).

SODIUM CONSUMPTION: AN INDIVIDUAL'S CHOICE? INT J HYPERTENS, 2012, 860954.

CAMPBELL, N. R., KACZOROWSKI, J., LEWANCZUK, R. Z., FELDMAN, R., POIRIER, L., KWONG, M. M., ET AL. (2010). 2010 CANADIAN HYPERTENSION EDUCATION PROGRAM (CHEP) RECOMMENDATIONS: THE SCIENTIFIC SUMMARY - AN UPDATE OF THE 2010 THEME AND THE SCIENCE BEHIND NEW CHEP RECOMMENDATIONS. CAN J CARDIOL, 26(5), 236-240.

CAMPBELL, N. R., WILLIS, K. J., L'ABBE, M., STRANG, R., AND YOUNG, E. (2011). CANADIAN INITIATIVES TO PREVENT HYPERTENSION BY REDUCING DIETARY SODIUM. NUTRIENTS, 3(8), 756-764.

CORSINO, L., SVETKEY, L. P., AYOTTE, B. J., AND BOSWORTH, H. B. (2009). PATIENT CHARACTERISTICS ASSOCIATED WITH RECEIPT OF LIFESTYLE BEHAVIOR ADVICE. N C MED J, 70(5), 391-398.

ENGSTROM, A., TOBELMANN, R. C., AND ALBERTSON, A. M. (1997). SODIUM INTAKE TRENDS AND FOOD CHOICES. AM J CLIN NUTR, 65(2 SUPPL), 704S-707S.

FORTMANN, S. P., WINKLEBY, M. A., FLORA, J. A., HASKELL, W. L., AND TAYLOR, C. B. (1990). EFFECT OF LONG-TERM COMMUNITY HEALTH EDUCATION ON BLOOD PRESSURE AND HYPERTENSION CONTROL. THE STANFORD FIVE-CITY PROJECT. AM J EPIDEMIOL, 132(4), 629-646.

FRANCIS, S. L., AND TAYLOR, M. L. (2009). A SOCIAL MARKETING THEORY-BASED DIET-EDUCATION PROGRAM FOR WOMEN AGES 54 TO 83 YEARS IMPROVED DIETARY STATUS. JOURNAL OF THE AMERICAN DIETETIC ASSOCIATION, 109(12), 2052-2056.

GLASCOFF, M. A., TAYLOR, S., GLASCOFF, D. W., AND RAFF, L. G. (1986). A SOCIAL MARKETING APPROACH TO REDUCING SALT INTAKE. HEALTH EDUC, 17(2), 11-14.

IOM (INSTITUTE OF MEDICINE). (2010). STRATEGIES TO REDUCE SODIUM INTAKE IN THE UNITED STATES (2011/01/07 ED.). WASHINGTON, DC: NATIONAL ACADEMIES PRESS.

JOHNSON, C. C., AND NICKLAS, T. A. (1995). HEALTH AHEAD--THE HEART SMART FAMILY APPROACH TO PREVENTION OF CARDIOVASCULAR DISEASE. AM J MED SCI, 310 SUPPL 1, S127-132.

KUMANYIKA, S. K., COOK, N. R., CUTLER, J. A., BELDEN, L., BREWER, A., COHEN, J. D., ET AL. (2005). SODIUM REDUCTION FOR HYPERTENSION PREVENTION IN OVERWEIGHT ADULTS: FURTHER RESULTS FROM THE TRIALS OF HYPERTENSION PREVENTION PHASE II. J HUM HYPERTENS, 19(1), 33-45.

LEGOWSKI, B., AND LEGETIC, B. (2011). HOW THREE COUNTRIES IN THE AMERICAS ARE FORTIFYING DIETARY SALT REDUCTION: A NORTH AND SOUTH PERSPECTIVE. HEALTH POLICY, 102(1), 26-33.

LUEPKER, R. V., PERRY, C. L., MURRAY, D. M., AND MULLIS, R. (1988). HYPERTENSION PREVENTION THROUGH NUTRITION EDUCATION IN YOUTH: A SCHOOL-BASED PROGRAM INVOLVING PARENTS. HEALTH PSYCHOL, 7 SUPPL, 233-245.

MCKAY, R. B., LEVINE, D. M., AND BONE, L. R. (1985). COMMUNITY ORGANIZATION IN A SCHOOL HEALTH EDUCATION PROGRAM TO REDUCE SODIUM CONSUMPTION. J SCH HEALTH, 55(9), 364-366.

MOHAN, S., CAMPBELL, N. R., AND WILLIS, K. (2009). EFFECTIVE POPULATION-WIDE PUBLIC HEALTH INTERVENTIONS TO PROMOTE SODIUM REDUCTION. CMAJ, 181(9), 605-609.

PAPADAKIS, S., PIPE, A. L., MOROZ, I. A., REID, R. D., BLANCHARD, C. M., COTE, D. F., ET AL. (2010). KNOWLEDGE, ATTITUDES AND BEHAVIOURS RELATED TO DIETARY SODIUM AMONG 35- TO 50-YEAR-OLD ONTARIO RESIDENTS. CAN J CARDIOL, 26(5), E164-169.

PAPADAKIS, S., PIPE, A. L., REID, R. D., AND BLANCHARD, C. (2010). SHORT TERM IMPACTS OF THE CHAMPLAIN 'GIVE YOUR HEAD A SHAKE' SODIUM REDUCTION CAMPAIGN ON KNOWLEDGE, ATTITUDES AND BEHAVIOURS. CANADIAN JOURNAL OF CARDIOLOGY, 26, 96D-96D.

PIANE, G. (1990). A COMPARISON OF THE EFFECT OF A HYPERTENSION EDUCATION PROGRAM AMONG BLACK AND WHITE PARTICIPANTS. J HEALTH CARE POOR UNDERSERVED, 1(2), 243-253.

VALENZUELA LANDAETA, K., AND ATALAH SAMUR, E. (2011). [GLOBAL STRATEGIES TO REDUCE SALT INTAKE]. ARCH LATINO-AM NUTR, 61(2), 111-119.

WEBSTER, J. L., DUNFORD, E. K., HAWKES, C., AND NEAL, B. C. (2011). SALT REDUCTION INITIATIVES AROUND THE WORLD. JOURNAL OF HYPERTENSION, 29(6), 1043-1050.

WHELTON, P. K., HE, J., APPEL, L. J., CUTLER, J. A., HAVAS, S., KOTCHEN, T. A., ET AL. (2002). PRIMARY PREVENTION OF HYPERTENSION: CLINICAL AND PUBLIC HEALTH ADVISORY FROM THE NATIONAL HIGH BLOOD PRESSURE EDUCATION PROGRAM. JAMA, 288(15), 1882-1888.

Social Marketing Plan Outline

Social Marketing Plan

I. Situation Analysis

Background

Describe problem incidence and severity in addition to drivers and determinants.

Previous/Current Efforts

What previous efforts have been made to promote the desired product, service, or behavior? Describe previous and/or current programs. What lessons have been learned that could inform future planning?

Gap Analysis

Based on the current communication environment, what is missing and/or would complement existing communication efforts?

SWOT Analysis

- Internal Forces: *What organizational Strengths can be maximized?* • *What Weaknesses should be minimized?*
- External Forces: *What Opportunities can you capture?* • *What potential Threats can you prepare for?*

II. Target Audience

Primary

This is the 'bulls-eye' target audience, oftentimes those most at risk.

Develop an audience profile that includes demographics, psychographics, and current behaviors.

Secondary

Who are the people that influence your primary audience? Common examples include healthcare providers, family members, and community leaders.

III. Research Findings

- List research available and summarize key findings. (In place of a summary in the strategy document itself, reference a separate report or attach as an appendix).
- Describe any additional formative research required or planned. [Note: if the need for additional formative research is significant, this will be a preliminary strategy or outline at this point, to be completed after research results are available].

IV. Audience Analysis

- Current knowledge, beliefs and behaviors.
- (Real and Perceived) Barriers.
- (Real and Perceived) Benefits.

V. Overview Of Supply Side

Describe key supply side issues currently being addressed, sites to be included in the program, main activities (provider/retailer training, branded and non-branded condom availability, pricing), and approximate timeline of supply-side interventions.

VI. Demand Side/Communication Goals And Objectives

Describe the overall communications goal and specific communications objectives.

- *Knowledge Objectives*
(PROVIDE EXAMPLES BASED ON AUDIENCE'S CURRENT KNOWLEDGE LEVELS/PERCEPTIONS).
- *Belief Objectives*
(PROVIDE EXAMPLES BASED ON AUDIENCE'S CURRENT ATTITUDES/BELIEFS).
- *Behavioral Objectives*
(LIST ACHIEVABLE AND MEASURABLE DESIRED BEHAVIORS).

VII. Marketing Mix Strategies

The Marketing Mix (or 4 P's) ensures that you have considered potential product innovation, supply side issues, and pricing in addition to the promotional or demand generation strategy. Attached is a Marketing Mix Planning tool to help you in your strategic planning.

Product

This can be a tangible product or service or a desired behavior. How will you 'position' or modify it to make it

more appealing to the target audience? Are there any new products or product innovation you could design?

Price

'Costs' associated with the product, service, or behavior can be monetary or non-monetary (perceived). Identify costs associated with exiting current behavior and adopting or maintaining the desired behavior. What will you do to lower exit costs and reward the new behavior?

Place

Where does the audience perform or think about the desired behavior, use the product, or access the service? How will you make this place more appealing or conducive? How will you intervene at the moment of choice?

Promotion

Your promotion strategy should include a Creative Brief which will serve as strategic direction for your creative strategy. In addition, outline key messages and communication channels in this plan.

VIII. Monitoring and Evaluation Plan

- What objectives will be measured?
- What techniques and methodologies will be used?
- When will measurements be taken?
- How will measurements be reported and to whom?
- How will M & E data be used to inform revisions?

IX. Timeline and Implementation Plan

- Will you roll out your program in phases? For example, if it includes a supply side intervention, will you start there to ensure that your offering is of quality?
- How will the phases be organized (i.e., by market/audience, objectives, activities?)
- For each phase, what will be done, who will be responsible, when will it be done, and what is the estimated cost?

X. Budget/Funding Sources

- What financial costs will be associated with the program?
- If costs exceed available funds, what additional funding sources will be explored?
- What strategies will you use to appeal to potential donors?
- What potential cost-share or in-kind contributions can be made by partners; i.e. a media partner?

Marketing Mix Planning Tool

The Marketing Mix (4 P's) Strategic Planning Tool

TARGET AUDIENCE	PRODUCT	PRICE	PLACE	PROMOTION
Primary	Tangible product, service, or desired behavior. <i>How will you 'position' or modify it to make it more appealing to the target audience? Also consider new product ideas.</i>	'Costs' (monetary and non-monetary) associated with the product, service, or behavior. <i>What will you do to decrease costs and reward/reinforce desired behavior?</i>	Where the audience performs or thinks about the desired behavior. <i>How will you make this place more convenient and/or intervene at a key moment?</i>	Outline key messages, messengers, and communication channels as part of a CREATIVE BRIEF before developing CREATIVE CONCEPTS .
Secondary				

Creative Brief Form

CREATIVE BRIEF	PROJECT/DATE:
1. Target Audience(s) <i>Describe the person that you want to reach with your communication. Include a primary and secondary (influencer) audience if appropriate. Include a summary of any relevant audience research.</i>	
2. Objective(s) <i>What do you want your target audiences to think, feel, or do after experiencing the communication?</i>	
3. Obstacles <i>What beliefs, cultural practices, pressure, misinformation, etc. stand between your audience and the desired behavior?</i>	
4. Key Promise <i>Select one single benefit that will outweigh the obstacles in the mind of your target audience. Suggested format: If I (desired behavior), then (immediate benefit).</i>	
5. Support Statements <i>This is the substantiation for the key promise; i.e.; the reasons why the promise is true. Oftentimes, this will begin with the word 'because'.</i>	
6. Tone <i>What feeling should your communication have? (e.g. authoritative, humorous, emotional, etc)</i>	
7. Communication Channels <i>What channel(s) will you employ? TV spots/drama? Radio? Print Ads? Point-of-purchase? Promotional giveaways? Earned Media (PR)? Interpersonal? Internet or SMS? Community events? All of the above?</i>	
8. Openings <i>What opportunities (times and places) exist for reaching your audience? When is your audience most open to getting your message? (e.g. Mothers Day, Heart Health Month)</i>	
9. Creative Considerations -Any additional considerations i.e. branding guidelines, language, visual identity?	

Bravo Criteria Checklist

IS YOUR BEHAVIOR CHANGE COMMUNICATION BRAVO?

Use this checklist to evaluate the creative before the target audience does.

- B** Are you promoting a clear **BEHAVIOR**? ✓
If so, what is it? If not, what, if anything could you do to revise?
- R** Is it **RELEVANT/RESEARCH-DRIVEN**? ✓
If the materials were driven by audience research, what were the insights?
- A** Is it **ATTENTION-GETTING**? ✓
Would you stop to look or listen? More importantly, would your target audience?
- V** Are you offering anything of **VALUE**? ✓
What are you offering the audience in exchange for adopting the behavior or accessing the product or service?
- O** Does it meet your **OBJECTIVES**? ✓
Does it meet the objectives identified in your Creative Brief?

Implementation Plan Template

IMPLEMENTATION PLAN																
Activity List each activity. Be as specific as possible. Use additional pages if necessary.	Target Audience Who is the audience for each activity?	12 Month Timeline Place an X in the appropriate box to indicate the first and last month of each activity.												Person(s) Responsible Who is responsible for implementing the activity?	Evaluation Indicators How will you measure the success of each activity?	
		1	2	3	4	5	6	7	8	9	10	11	12			



**Monitoring and Evaluation
of Salt Consumption,
Sources of Salt and
Knowledge and Behaviors**

**Protocol for Population Level
Sodium Determination
In 24-Hour Urine Samples**

Section 1: Introduction

Overview of the PAHO/WHO Protocol for Population Level Sodium Determination in 24-Hour Urine Samples

The PAHO/WHO Protocol for Population Level Sodium Determination in 24-hour Urine Samples is a resource to countries that want to start, contribute to and share information on dietary salt reduction initiatives. It will assist with:

- Planning and preparing the scope and environment for a survey study to estimate dietary salt intake.
- Recruiting and training field staff for data collection.
- Reporting and disseminating the results.

While the substance of concern to health is sodium, strategies to reduce its intake are aimed at its main source in the diet – salt (sodium chloride) – used in the household at the table or in cooking and as an additive in industrially-manufactured foods.

Primary aims

- Estimate the average intake of dietary salt in men and women in the Americas in the age stratum 25 to 64 through measurement of 24 hour urinary sodium excretion.
- Provide information for designing and implementing interventions aimed at reducing population level dietary salt.
- Determine subsequent estimates of salt intake in the same population in aid of monitoring intake over time.
- Provide trends in salt intake against which to monitor and evaluate the effectiveness of interventions aimed at population level dietary salt reduction.

Additional aims

- Estimate the average intake of dietary potassium through joint measurement of 24-hour urinary potassium excretion.
- Estimate the average intake of iodine through joint measurement of 24-hour urinary iodine excretion.
- Determine creatinine excretion.

Other possible aims

- Estimate intake of sodium, potassium and iodine in populations otherwise differentiated e.g. by ethnicity, socio economic status, geographic location, other target age groups, etc.

- Support health economic analysis by estimating salt intake for specific age strata.
- Estimate fluoride excretion as well.

Intended audience

The protocol is primarily intended for principle investigator (s) of studies of sodium, potassium and iodine intake. Parts of the manual are also intended for field staff who are to interact with survey participants.

Structure

The protocol has seven Sections following a sequence that helps to implement population level sodium, potassium and iodine determination in 24-hour urine samples. Section 8 shows the full dataset required for health economic analysis of sodium reduction strategies.

There is both general information and specific instructional material that can be extracted and used for:

- Training.
- Data collection.

Important conversions

5g salt (NaCl)=2,000 mg sodium=87 mmol sodium=87 mEq sodium
23 mg sodium = 1 mmol sodium
39.1 mg potassium = 1 mmol potassium
126.9 mg iodine = 1 mmol iodine
113.12 g creatinine = 1 mol creatinine

Rationale for Population Level Sodium Determination in 24-Hour Urine Samples

Background

In Latin America and the Caribbean, chronic non-communicable disease (CNCD) is the main cause of disability and premature mortality (1). Hypertension, a principal risk factor for a number of CNCD, in particular cardiovascular (CVD) and renal diseases, affects up to a third of adults in the Pan American Region (2). There is compelling evidence (epidemiological, clinical and animal-experimental) of the direct relationship between salt consumption and blood pressure (BP) and that current levels of salt intake are a major factor increasing BP (3,4,5). If people reduce dietary salt, whether they are normotensive or hypertensive, raised blood pressure can be avoided, hypertension better controlled, thousands of deaths from stroke, heart and renal disease prevented (6) and healthcare systems spared substantial treatment and health-related costs (7,8,9,10,11).

PAHO is spearheading an initiative, guided by an Experts Group, to reduce dietary salt intake at the population level across the Americas. Its first product, a Policy Statement, has the goal – reduce salt intake to the internationally recommended target of <5g per adult per day by 2020 (12).

Rationale for surveillance of salt intake

Fundamental to the PAHO initiative is for Member States to estimate a baseline of population level dietary salt intake, and from there, to monitor trends in intake and the effectiveness of any interventions within and between populations.

The best estimate of the population profile distribution and average level of dietary salt intake is provided by measuring 24-hour urinary sodium excretion in a representative sample of individuals (13).

Rationale for complementary food consumption information

To guide policy development and associated population level interventions aimed at reducing dietary salt, not only is information needed on salt intake but also on the main food sources of salt in the diet and the typical frequency of their consumption. There are several methods available to collect information on food consumption, among them 24-hour food recall. The INTERMAP Study is an international, cross-sectional, epidemiologic study where in-depth 24-hour dietary recall was used to identify foods that account for most dietary sodium intake (14).

While the instruments that collect food consumption information are typically very detailed in terms of the food products listed in order for survey participants to be able to select the specific products they consume, it is recommended to group the products into a smaller number of broad categories. They become the basis for raising awareness among consumers as to the food categories that contribute the most salt to the diet, and are also the basis for policies and interventions with industry that include target setting per category. If a category is too wide and varied, it is difficult to set a target; if there are too many categories, target setting and monitoring can become unmanageable.

There are a number of examples of food categories to consider, among them the 12 food categories used in the Salt Campaign of the European Commission (15) and the 19 basic product groups and 8 non-basic groups in the Choices Programme (16).

Rationale for joint surveillance of potassium

Low dietary potassium is associated with hypertension (17) and stroke (18) and supplementing potassium to hypertensive individuals lowers blood pressure (19) and reduces the use of anti-hypertensive medications (20). Increased potassium intake also reduces the hypertensive response to high dietary sodium. Some populations are deficient in dietary potassium if they rely on processed foods, however there is a deficiency in data on intake of potassium in most populations. Estimating potassium and sodium intake at the same time can inform the design of potential population interventions to improve both sodium and potassium intakes.

Rationale for joint surveillance of iodine

To address the concern regarding the possible detrimental effect of dietary salt reduction on programs to prevent Iodine Deficiency Disorder (IDD) that rely on salt as a carrier of iodine, it is recommended that iodine intake be assessed along with salt. The inclusion of this variable in studies of salt intake that use 24-hour urine samples would in fact benefit IDD-prevention programs. The method provides the most accurate and appropriate indicator of whether populations, regardless of age, gender or climatic environment, are receiving the recommended amounts of this nutrient, which, judging from current salt intake and salt iodization levels, may be insufficient, sufficient and even excessive (21).

Use of spot- or timed urine testing

Collecting 24-hour urine samples has been considered difficult, and therefore the use of the spot-urine method has been proposed as an alternative. To estimate intake of sodium, potassium and iodine, the use of spot urine is not recommended unless the following conditions are met:

- A baseline estimate of these analytes has been conducted using the recommended methods for 24-hour urine assessment.
- A calibration study for use of spot urine has been done in the specific population of interest.

Once the above conditions are met, ‘timed’ urine collections (over three or more hours with provision of water) are preferred over non-timed (‘spot’) samples as they reduce the errors due to residual urine in the bladder.

Even if the above conditions are met, the results are likely to be unreliable especially for population subgroups or time trends.

See Section 7 for further information and advice on calibration.

Section 2: Field Protocol

Overview of the Field Protocol

Components

The protocol for Sodium Determination in 24-Hour Urine Samples can stand-alone or be an additional module to an existing CNCD risk factor instrument (e.g. PanAmerican STEPS – the Pan American Version of the WHO STEPwise Approach to Risk-Factor Surveillance) (22). If stand-alone, the following are the required components of the protocol:

	Description	Purpose
1	Questionnaire on demographic and behavioral information.	To obtain data on: Socio-demographic information. Tobacco and alcohol use. Dietary habits. Physical activity. Knowledge, attitudes and behavior towards dietary salt.
2	Questionnaire on personal medical history, including drug treatment.	To determine the proportion of adults that: Currently suffer from CNCD, and their complications. Are under daily long term medical treatment for any condition.
3	Physical measurements with simple methods.	To determine the proportion of adults who: Are overweight and obese, and Have high blood pressure.
4	24-hour urine sample collection.	To determine sodium, potassium and iodine excretion. To determine creatinine excretion.
5	A 50-100 g sample of household salt.	To determine the iodine content of household salt.

If performed as part of another risk factor study that collects the data described in components 1 to 3, only components 4 and 5 of the protocol are required. The data elements for components 1 to 3 are provided below. They were developed with reference to the framework for risk factor surveillance in PanAmerican STEPS and an instrument from the University of Warwick WHO Collaborating Centre for Nutrition. The PAHO/WHO Expert Group for Cardiovascular Disease Prevention through Population-wide Dietary Salt Reduction developed the questions on knowledge, attitudes and behavior towards dietary salt.

Core and expanded data

Each of the first three components of the protocol has a minimum core of required data and a set of expanded desirable data for collection, shown below. Whether core or core plus expanded data are collected depends

on what can realistically be accomplished in each country setting (financially, logistically and in terms of human and clinical resources).

	Core	Expanded
1	<ul style="list-style-type: none"> Basic demographic information including: <ul style="list-style-type: none"> Country and region of origin (if relevant). Age. Sex. Tobacco use. Alcohol consumption. Physical activity. Sedentary behavior. Fruit and vegetable consumption. Knowledge, attitudes and behavior towards dietary salt. 	<ul style="list-style-type: none"> Expanded demographic information including: <ul style="list-style-type: none"> Ethnicity. Highest level of education. Employment. Household income. History of tobacco use. Patterns of alcohol drinking. Oil and fat consumption. History of raised blood pressure. History of diabetes.
2	Current drug treatment used. Personal medical history.	Family medical history.
3	Height (cm) and weight (kg). Waist circumference (cm). Systolic and diastolic blood pressures (mmHg) and heart rate (bpm).	Hip circumference (cm).

Planning and Conducting a 24-Hour Urine Collection Study

Below are the recommended tasks to plan and conduct a 24-hour urine collection study. The timeframes will be situation specific, to be estimated to support the planning process.

Intended audience

This information is primarily intended for those fulfilling the following roles:

- Site coordinator.
- Coordinating committee.

Tasks and timeframes	
Tasks	Timeframe
Develop implementation plan.	
Identify scope of study.	
Gain ethical approval.	
Schedule data collection.	
Adapting and translating the Field Protocol Questionnaire.	
Pilot test.	

Selecting the Sample

Sample population

The sample size is determined by precision, variability within and between subjects, statistical power, play of chance, representativeness, feasibility and cost. Below is a matrix showing the relationship between sample size, precision in the difference in excreted sodium to be detected and variations in measurements.

In general, to detect approximately 1 g reduction in salt intake over time using 24-hour urinary sodium excretion, with a standard deviation of 75 mmol/day (alpha = 0.05, power = 0.80), a minimum sample of 120 individuals per age and sex stratum is recommended. To account for attrition (e.g. non participation, incomplete collection or implausible values), which may be as high as 50%, up to 240 people per age and sex stratum should be invited to participate.

Requirements for sample selection

- Random or otherwise probabilistic sample.
- Sample selected using culturally appropriate methods.

- Stratification by age group and sex with a minimum of four groups i.e. men and women each in two age groups 25-44 and 45-64 (or men and women each in four age groups 25-34, 35-44, 45-54, 55-64).
- If a sentinel site is selected, must be justifiable and feasible for long term monitoring.
- Age and sex of respondents and non-respondents are noted.
- If sodium excretion data from 24-hour urine samples are to inform health economics analysis of changes in sodium intake, see the table below for the full dataset required.

Exclusion criteria

- People unable to provide informed consent.
- Those with known history of heart or kidney failure, stroke, liver disease.
- Those who recently began therapy with diuretics (less than two weeks).
- Any other conditions that would make 24-hour urine collection difficult.

If pregnant women are included in the sample, their results must be analyzed separately from those of other adult participants.

Matrix to Determine Sample Size

Minimum difference in sodium excretion to be detected δ (mmol/day)	Standard deviation s (SD)	Sample size n (for each age stratum)
10	10	16
10	15	35
10	20	63
10	25	98
10	30	141
10	35	192
10	40	251
10	45	318
10	50	392
10	55	475
10	60	565
10	65	663
10	70	769
10	75	883
10	80	1005
20	10	4
20	15	9
20	20	16
20	25	25
20	30	35
20	35	48
20	40	63
20	45	79
20	50	98
20	55	119

$$n = 2 \frac{\left[z \frac{(1-\alpha)}{2} + z(1-\beta) \right]^2}{\Delta^2}$$

where $\alpha = 0.05$ and $(1-\beta) = 0.80$, 1.96 and 0.8416 respectively

$\Delta = \delta/s$

where $\Delta =$ standardized difference i.e. $(\mu_1 - \mu_2) / s$

$\delta =$ clinically important difference to be detected

$s =$ standard deviation

Minimum difference in sodium excretion to be detected δ (mmol/day)	Standard deviation s (SD)	Sample size n (for each age stratum)
20	60	141
20	65	166
20	70	192
20	75	221
20	80	251
30	10	2
30	15	4
30	20	7
30	25	11
30	30	16
30	35	21
30	40	28
30	45	35
30	50	44
30	55	53
30	60	63
30	65	74
30	70	85
30	75	98
30	80	112
40	10	1
40	15	2
40	20	4
40	25	6
40	30	9
40	35	12
40	40	16
40	45	20
40	50	25
40	55	30
40	60	35
40	65	41
40	70	48
40	75	55
40	80	63
50	10	1
50	15	1
50	20	3
50	25	4
50	30	6
50	35	8
50	40	10
50	45	13
50	50	16
50	55	19
50	60	23
50	65	27
50	70	31
50	75	35
50	80	40

$$n = 2 \frac{\left[z_{\frac{(1-\alpha)}{2}} + z_{(1-\beta)} \right]^2}{\Delta^2}$$

where $\alpha = 0.05$ and $(1-\beta) = 0.80$, 1.96 and 0.8416 respectively

$\Delta = \delta/s$

where $\Delta =$ standardized difference i.e. $(\mu_1 - \mu_2) / s$

$\delta =$ clinically important difference to be detected

$s =$ standard deviation

Implementation Plan

A detailed implementation plan for the 24-hour Urine Sample study is needed for all stakeholders involved in the surveillance process.

Purpose

The implementation plan is to:

- Set out the scope of the surveillance and desired goals.
- Identify required resources.

- Lay out an action plan.
- Develop a communication strategy.
- Provide a budget as the basis for funding.

Core parts of the implementation plan

Below are the core parts needed for the implementation plan. Some have references to Sections within this document where there is information to assist with preparation.

Core part	Detail	References
Executive summary	High level summary of main points including: Current situation. Goals and objectives. Scope. Resources. Budget.	Section 1
Current situation	Specify: Whether the study will determine a baseline of sodium intake or assess change in intake. If to assess change in intake, reference the baseline study. If a risk factor survey has already been conducted. If there is an existing infrastructure (human capacity, equipment, other studies) on which the 24-hour urine sample collection could be built.	
Goals and objectives	Identify planned goals and use of the information collected to: Describe the current level of dietary salt intake in populations (if available). Track the direction and magnitude of trends in salt consumption. Plan and evaluate a health promotion or preventive campaign. Collect data from which to predict likely future demands for health services. Specify objectives that support gathering 'essential' information only. Describe broad timeframes.	
Scope	Specify the scope of surveillance to be conducted (coverage of core and expanded data). Specify if future sodium determination surveillance can be assured.	Section 2
Sampling method	Identify the sample size and sample frame that will be used. Identify geographical coverage. Describe sample design.	Section 2
Resources	Specify the resources in terms of all personnel and equipment required for sodium determination in 24-hour urine sampling study. Describe resources that have been committed or expected, including support from PAHO/ WHO. Specify resources from other organizations.	
Action plan	Prepare a chart of the main tasks with estimated start date and time-frame for completion of each.	Section 2
Communication strategy	Specify the methods for informing and involving all stakeholders relevant to the sodium determination project, including community leaders, members of the public, and media.	
Budget	Provide a detailed budget that includes: Total funds required for each year planned to implement all sodium determination activities as identified in the scope (including future surveys). Sources of funding. Funding gaps.	

Applying for Ethical Approval

Studies that are to use the PAHO/WHO Protocol for Sodium Determination in 24-hour Urine Samples must undergo technical and ethical review and approval. This is to ensure that the study:

- Is conducted in a technically and ethically sound manner;
- Recognizes and protects the rights of participants; and
- Ensures wide access to the information collected in the study.

Process

Usually, ethical approval should be sought by submission of a proposal and application to a national ethics review committee or other equivalent body. However, if such a body is not institutionalized, it is recommended that an application for ethical review be prepared and submitted through an ad hoc local mechanism within the Ministry of Health.

Informed consent

The informed consent must be obtained from every survey participant before conducting any interviews or collection of any samples.

Making a submission

Use the existing templates for proposals supplied by the appropriate ethics committee or equivalent body. If such a template does not exist, identify and contact the relevant bodies, seek guidance on rules, the submission process and any procedures to follow.

Timeframes and Data Collection Considerations

Data collection should be carefully planned to take place over a defined period of time and during appropriate seasons.

General timeframes

The following table shows the recommended phases of a sodium determination study. Timeframes are situation specific:

Phase	Timeframes
Planning and scoping	
Recruiting and training	
Data collection	
Data analysis and reporting	

Data collection

Some key factors to consider when identifying an appropriate time to conduct the study:

Factors to consider	Guidelines
Seasons	Confine the study period to one season to avoid dietary changes Avoid festive seasons (E.g. Ramadan, Christmas, Holy Week, and other national or religious holidays) Avoid seasons when food is in unusually short supply.
Calendar year	Confine the study to one calendar year
Major events	Avoid data collection during periods prior to local, regional, or national elections to avoid confusion with political campaigns.
Civil unrest, turmoil, famine, etc.	Avoid conducting a study at any time when pressing matters occupy the minds and lives of the population.
Collection timeframe	Keep the timeframe as close as possible (within reason) to the recommended timeframe.

Data collection locations

It is recommended that all components of the study be conducted/administered in the household setting. Ideally participants/respondents are to collect all their urine samples at home and otherwise, they are to bring home any urine passed away from home. The total urine passed in the 24-hour period is to be picked up at the household within one day of the 24-hour collection period. It is recommended that if food consumption information is collected, this is done during the second visit to the household.

Adapting the PAHO/WHO Protocol for Sodium Determination in 24-Hour Urine Samples

Using a standardized protocol for Sodium Determination in 24-hour Urine Samples enables comparisons between countries. However, some adaptations may be required to account for differences in cultures or settings.

When to adapt the protocol

Adaptations may be needed to provide valid data from the surveillance. The following are often what need adaptation: terminology, providing additional information, deleting questions on behaviors that do not apply.

Process

The process of adapting the protocol may involve the following:

- Identifying the instructions or questions that require local adaptation.
- Adding or deleting questions.
- Adding other forms as appropriate.

- Seeking feedback and advice.
- Translating and back translating the adapted instructions or questionnaires.
- Pilot testing the questionnaires.

Documents to translate

Below are some of the documents that may need translating, including where they can be found:

Documents	References
Component 1 questionnaire	PanAmerican STEPS
Component 2 questionnaire	PanAmerican STEPS
Guidelines for field work	Section 3
Consent forms	PanAmerican STEPS
Knowledge, attitudes and behavior questionnaire	Section 4
Instructions to participants	Section 5

Pilot Testing

A pilot test of the entire data collection process must be conducted among a limited number of people with a broad range of backgrounds prior to implementing the actual full study. Pilots should involve all aspects of the survey including:

- Approaching potential participants.
- Seeking and obtaining informed consents.
- Making arrangements/appointments for second visits after the participant-led 24-hour urine sample collection.
- Site preparation and set-up.
- Collecting all data needed.
- Identifying participants who may need a follow-up.
- Basic analysis.

Test group

Identify and approach willing participants to be part of the pilot test. The test group should include the following:

- Both men and women.
- Cover the age range 25-64.
- More than one ethnic group (if appropriate).
- Participants with different levels of education.
- Participants from a range of socio-economic groups.
- Participants from distinctly different regions in the same country.

Test environment

Where possible conduct the pilot test under the field conditions expected for the final full study i.e. the household setting.

Timeframe

When planning the pilot test, allow sufficient time for adjustments to be made prior to starting full data collection.

Section 3: Data Collection Guide

Guidelines for data collection for components 1 through 3 of the protocol can be obtained from the Pan American STEPS Manual, Part 3, Sections 1 through 4, except for the core questions on knowledge, attitudes and behavior towards dietary salt, which are in Section 4 of this manual.

The information below serves the field staff/survey team involved in components 4 and 5 of the protocol for 24-hour urine sample collection.

Instructions for Field Staff, Equipment and Analytic Methods

Instructing participants

Field staff must explain the collection protocol, obtain informed consent and provide the record sheet on which participants note the start and finish times of their 24-hour urine collection, any missed urine collections, and any medication taken during the collection.

In the morning of the start of the 24-hour period, the participant must void the bladder and note the time. **This “first-pass urine” is discarded.** All urine passed thereafter is collected in the container provided, including the first urine of the following morning, with the final time recorded. Respondents are given detailed written instructions (see Section 5).

At the time of the first visit to the household, field staff must inform the participant of the second visit.

The second visit must be made within one day of the completion of the 24-hour collection period. A sample of household salt is taken during the second visit.

If food consumption information is required, it is collected during the second visit.

Equipment supplied to participants

- A 5 liter capacity screw cap container to store the collected urine.
- A 1 litre container with a wide opening into which urine is voided, with or without the use of a funnel.
- Optional 2 liter capacity screw cap container for temporal collections of urine made away from the home.
- Funnel for women to be used during urine collection, kept inside a re-sealable plastic bag when not used.
- Plastic carrier bags for transporting the equipment away from home.
- An aide-memoire to help participants remember to collect their urine e.g. a safety pin to pin the under- and outer garments.

ments together during the period of the collection as a reminder that the urine about to be passed should be collected.

The use of PABA to assess completeness of the urine collection is not recommended. It requires that each participant take a PABA pill three days prior to the start of collection thereby increasing the risks of non-compliance and attrition. In addition, laboratory facilities for the testing of PABA in the urine are limited and where they exist, will increase the costs of the study.

At the completion of the collection

- Field staff measure the total volume of urine, mix it thoroughly in its container and withdraw three 10-ml aliquots into separate labelled tubes for storage and shipping for analysis. The rest of the urine is discarded.
- Sodium, potassium, iodine and creatinine content in the urine are to be measured in certified laboratories, as is the iodine content of the household salt.

Analytic methods

- Sodium and potassium content in the urine may be determined through Ion Selective Electrode (indirect) with a Beckman Coulter Synchron CX5PRO System.
- Creatinine content may be determined through the Creatinine (urinary) Jaffe kinetic method, standardized, also to be measured by Beck Coulter synchron CX5PRO System.
 - Iodine in urine may be determined with the traditional kinetic method of Sandell-Kolthoff (23) or by Inductively Coupled Plasma (ICP) Spectrometry.
 - Iodine content of household salt can be determined quantitatively with the titration method. In addition to the titration method, there are possibilities of using potentiometry or spectro-photometry. (23)

Guide to Physical Measurements

Component 3 of the PAHO/WHO Protocol for Sodium Determination in 24-hour Urine Samples requires that selected physical measurements be taken to determine the proportion of participants in the study who:

- Have raised blood pressure.
- Are overweight and/or obese.

Below is a description of:

- The physical measures and what they mean.
- The equipment needed.
- How to assemble and use the equipment.
- How to take the measurements and accurately record the results.

Physical measurements

Blood pressure is measured to determine the proportion of participants with raised BP. Heart rate, measured at the same time as BP with automated devices, is a common independent cardiovascular risk factor. Height and weight measurements are taken to calculate the body mass index (BMI), needed to determine the prevalence of overweight and obesity in the population. Waist circumference measurements provide additional information on overweight and obesity. Hip circumference is an expanded data option to measure overweight and obesity.

Units of measurement

Below are the standard units for the physical measurements in component 3 of the protocol, including their upper and lower limits for data entry purposes.

Physical Measure	Unit	Minimum	Maximum
Systolic blood pressure (SBP)	mmHg	40	300
Diastolic blood pressure (DBP)	mmHg	30	200
Height	cm	100	270
Weight	kg	20	350
BMI (Body Mass Index)	kg/m ²	11	75
Waist circumference	cm	30	200
Hip circumference	cm	45	300
Heart rate	beats/minute	30	200

Sequence of questions and measurement

As is the case with many risk factor studies, physical measurements are to be taken immediately after the personal medical history. Physical measurement results are to be recorded on the same participant instruments as personal medical history.

Participant instructions

Prior to taking physical measurements, explain to the participant that the following measurements will be taken:

For core

- Blood pressure.
- Heart rate.
- Height.
- Weight.
- Waist circumference.

For expanded, additional

- Hip circumference.

Measuring Blood Pressure and Heart Rate

Equipment needed

Validated digital automatic blood pressure monitor e.g. OMRON. For the choice of validated blood pressure measuring devices see http://www.bhsoc.org/bp_monitors/automatic.stm.

- Appropriate size cuffs.

Preparing the participant

Prior to measuring blood pressure, ask the participant to sit in a quiet comfortable place for at least 5 minutes with back support and his/her legs uncrossed. If the questions in components 1 and 2, on behavior and personal medical history, have been asked just before the physical measurements are to be taken, the participant should rest for at least 5 minutes before blood pressure measurement is started. Do not talk to the participant whilst BP is being taken.

Three measurements

WHO recommends taking three blood pressure measurements. During the data analysis, the mean of the second and third readings is calculated. The participant must rest for one minute between each of the readings. The measurement and recording of heart rate should be done three times along with the measurement and recording of blood pressure. Heart rate and blood pressure results are displayed simultaneously with automated equipment.

Recording the blood pressure measurements

The following steps are required:

- after each of the three measurements, record the result in the participant's instrument;
- after all three readings are taken, double-check that all three results are correctly recorded in the instrument;
- inform the participant of their blood pressure readings only after the whole process is completed.

OMRON procedure

The instructions below apply to the use of an OMRON blood pressure monitor. However, more detailed operating instructions are included with the device and should be reviewed before taking any blood pressure measurements.

Note that if a different digital automatic blood pressure monitor is used, instructions should be read carefully.

Applying the OMRON cuff

Follow the steps below to select an appropriate size of cuff and apply it:

Step	Action								
1	Place the left arm* of the participant on the table with the palm facing upward.								
2	Remove or roll up clothing on the arm.								
	Select the appropriate cuff size for the participant using the following table:								
	<table border="1"> <thead> <tr> <th>Mid Arm Circumference (cm)</th> <th>Cuff Size</th> </tr> </thead> <tbody> <tr> <td>17-22</td> <td>Small (S)</td> </tr> <tr> <td>22-32</td> <td>Medium (M)</td> </tr> <tr> <td>>32</td> <td>Large (L)</td> </tr> </tbody> </table>	Mid Arm Circumference (cm)	Cuff Size	17-22	Small (S)	22-32	Medium (M)	>32	Large (L)
Mid Arm Circumference (cm)	Cuff Size								
17-22	Small (S)								
22-32	Medium (M)								
>32	Large (L)								
3	If the cuff is the correct size, the marker at the end of the cuff will fit between two other markers in the mid section of the cuff. The cuff is the wrong size if the end is outside the markers. It is advisable to select the larger size cuff if there is a question of which size is best. Some Omron cuffs are not marked in which case they must be labeled with markers.** Otherwise, use the mid arm circumference of each arm to select the correct cuff size.								
4	Position the cuff above the elbow and aligning the mark ART on the cuff with the brachial artery.								
5	Wrap the cuff snugly onto the arm and securely fasten with the Velcro. Note: The lower edge of the cuff should be placed 1.2 to 2.5 cm above the inside of the elbow joint.								
6	Keep the level of the cuff at the same level as the heart during measurement.								

*If the right arm is used, indicate this in the right hand side margin of the participant's Instrument.

**Even if cuffs are marked by the manufacturer to indicate the acceptable range of arm circumference for the size of cuff, the markings may not agree with the current recommended range and need to be checked and possibly remarked. [24] Marking can be performed easily using a ruler and permanent marker. The ideal arm circumference for a cuff is 2.5 times the cuff's bladder width. Cuffs can be used on arms that have a circumference ± 4 cm of 'ideal'. To mark or remark the cuff, start the measurement at the end that contains the bladder. Permanently mark the cuff at the ideal arm circumference then draw a line across the cuff at 4 cm on either side of the ideal (ie draw two lines). The cuff is the right size if when wrapped around the mid arm, the end is between the two marked lines.

Taking the BP measurement with an OMRON

Follow the instructions below to take the blood pressure measurements:

Step	Action
1	Switch the monitor on (dark purple button) and press START (light purple button).
2	The monitor will start measuring when it detects the pulse and the "heart" symbol will begin to flash. The systolic and diastolic blood pressure readings should be displayed within a few moments (systolic above and diastolic below). The heart rate will also be displayed.
3	Record the reading in the participant's instrument.
4	Switch the monitor off, but leave the cuff in place.
5	Wait one minute, then repeat steps 1-4 two more times.
6	Inform the participant of the blood pressure readings only after the whole process is completed.

When to use a sphygmomano-meter

The sphygmomanometer is generally not recommended, but may be used in the following circumstances:

- the OMRON is not functioning;
- the OMRON display shows multiple errors;
- to cross check OMRON blood pressure readings in various clinical states such as irregular pulse, peripheral circulatory disturbance, extreme hypotension;
- when systolic BP is >200 mmHg (appropriate measurement of systolic BP requires inflating the cuff to a pressure of 40 mmHg above the systolic BP; OMRON maximum inflation pressure seldom exceeds 240 mmHg);
- for calibration of the OMRON Monitor.

Procedure for sphygmomano-meter

Follow the steps below and refer to the operating instructions included with the device to measure the blood pressure of a participant using the sphygmomanometer.

Step	Action
1	Apply the cuff (as detailed above).
2	Put stethoscope earpieces in ear and set to bell.
3	Palpate pulse at either brachial or radial artery. Take a pulse on count for one full minute.
4	Pump up pressure and inflate cuff until unable to feel pulse.
5	Continue to inflate cuff 40 mmHg beyond this point.
6	Apply the bell of the stethoscope to the right antecubital fossa.
7	Listen for pulse sounds while deflating the cuff slowly.
8	Record the systolic blood pressure (SBP) when a pulse is first audible.
9	Record the diastolic blood pressure (DBP) when the pulse sound disappears.
10	Deflate the cuff fully and let the arm rest for one minute (between each reading).
11	Repeat Steps 2-10 twice to obtain three readings. Record the readings to the nearest 2 mmHg.*
12	Check that all readings are correctly filled in on the instrument.
13	Inform the participant of the blood pressure readings only after the whole process is completed.

* Analyze blood pressure readings by 2 mmHg to test for terminal digit preference as a quality assurance method. (Terminal digit preference is the tendency to record to 10 mmHg rather than 2 mmHg.)

Measuring Height

Equipment needed

Portable height/length measuring board.

Assembling the measuring board

Follow the steps below to assemble the measuring board:

Step	Action
1	Separate the pieces of board (usually 3 pieces) by unscrewing the knot at the back.
2	Assemble the pieces by attaching each one on top of the other in the correct order.
3	Lock the latches in the back.
4	Position the board on a firm surface against a wall.

Measuring height

Follow the steps below to measure the height of a participant:

Step	Action
1	Ask the participant to remove their: -footwear (shoes, slippers, sandals, etc.) -head gear (hat, cap, hair bows, comb, ribbons, etc.) Note: If it would be insensitive to seek removal of a scarf or veil, the measurement may be taken over light fabric.
2	Ask the participant to stand on the board facing you.
3	Ask the participant to stand with: feet together heels against the back board knees straight.
4	Ask the participant to look straight ahead and not tilt their head up.
5	Make sure eyes are the same level as the ears.
6	Move the measuring arm gently down onto the head of the participant and ask the participant to breathe in and stand tall.
7	Read the height in centimeters at the exact point.
8	Ask the participant to step away from the measuring board.
9	Record the height measurement in centimeters in the participant's Instrument.

Measuring Weight

Equipment needed

- portable electronic weighing scale;
- a stiff wooden board to place under the scales, if you are likely to have problems with uneven surfaces (such as dirt or mud floors or carpet);
- a generator, if electronic scales are being used and electricity is not guaranteed in all survey areas (check if scale can work with batteries).

Set up requirements

Make sure the scales are placed on a firm, flat surface. Do not place the scales on:

- carpet;
- a sloping surface;
- a rough, uneven surface.

Electronic scales

Follow the steps below to put electronic scales into operation:

Step	Action
1	Put the scale on a firm, flat surface.
2	Connect the adaptor to the main power line or generator.
3	Turn on the scale.
4	Switch the scale on and wait until the display shows 0.0.

Measuring weight

Follow the steps below to measure the weight of a participant:

Step	Action
1	Ask the participant to remove their footwear (shoes, slippers, sandals, etc) and socks.
2	Ask the participant to step onto scale with one foot on each side of the scale.
3	Ask the participant to: <ul style="list-style-type: none">▶ stand still▶ face forward▶ place arms on the side and wait until asked to step off.
4	Record the weight in kilograms on the participant's instrument. If the participant wants to know his/her weight in pounds, convert by multiplying the measured weight by 2.2.

Measuring Waist Circumference

Equipment needed

- constant tension tape (for example, Figure Finder Tape Measure);
- pen;
- chair or coat stand on which the participant will place their clothes.

Privacy

A private area is necessary for this measurement. This could be a separate room, or an area that has been screened off from other people within the household.

Preparing the participant

This measurement should be taken without clothing, that is, directly over the skin.

If it is not possible, the measurement may be taken over light clothing. It must not be taken over thick or bulky clothing. This type of clothing must be removed.

How to take the measurement

This measurement should be taken:

- at the end of a normal expiration;
- with the arms relaxed at the sides;
- at the midpoint between the lower margin of the last palpable rib and the top of the iliac crest (hip bone).

Measuring waist circumference

Follow the steps below to measure the waist circumference of a participant:

Step	Action
1	Standing to the side of the participant, locate the last palpable rib and the top of the hip bone. You may ask the participant to assist you in locating these points on their body.
2	Ask the participant to wrap the tension tape around themselves and then position the tape at the midpoint of the last palpable rib and the top of the hip bone, making sure to wrap the tape over the same spot on the opposite side. Note: Check that the tape is horizontal across the back and front of the participant and as parallel with the floor as possible.
3	Ask the participant to: <ul style="list-style-type: none">▶ stand with their feet together with weight evenly distributed across both feet;▶ hold the arms in a relaxed position at the sides;▶ breathe normally for a few breaths, then make a normal expiration.
4	Measure waist circumference and read the measurement at the level of the tape to the nearest 0.1 cm, making sure to keep the measuring tape snug but not tight enough to cause compression of the skin.
5	Record the measurement on the participant's instrument.

Measuring Hip Circumference

Equipment needed

- constant tension tape (for example, Figure Finder Tape Measure);
- pen;
- chair or coat stand on which the participant will place their clothes.

Privacy

A private area is necessary for this measurement. This

could be a separate room, or an area that has been screened off from other people within the household. Hip measurements are taken immediately after waist circumferences.

Preparing the participant

This measurement should be taken without clothing, that is, directly over the skin.

If it is not possible, the measurement maybe taken over light clothing. It must not be taken over thick or bulky clothing. This type of clothing must be removed.

How to take the measurement

This measurement should be taken:

- with the arms relaxed at the sides;
- at the maximum circumference over the buttocks.

Measuring hip circumference

Follow the steps below to measure the hip circumference of a participant:

Step	Action
1	Stand to the side of the participant, and ask them to help wrap the tape around themselves.
2	Position the measuring tape around the maximum circumference of the buttocks.
3	Ask the participant to: ▶ stand with their feet together with weight evenly distributed over both feet; ▶ hold their arms relaxed at the sides.
4	Check that the tape position is horizontal all around the body and snug without constricting.
5	Record the measurement on the participant's Instrument. Note: measure only once and record.

Section 4: Questionnaire on Knowledge, Attitudes, and Behavior toward Dietary Salt

1. Do you add salt to food at the table?

- a) never
- b) rarely
- c) sometimes
- d) often
- e) always

2. In the food you eat at home salt is added in cooking

- a) never
- b) rarely

- c) sometimes
- d) often
- e) always

3. How much salt do you think you consume? (READ LIST)

- a) Far too much
- b) Too much
- c) Just the right amount
- d) Too little
- e) Far too little
- f) Don't Know
- g) Refused

4. Do you think that a high salt diet could cause a serious health problem?

- a) Yes
- b) No
- c) Don't know
- d) Refused

5. If Yes in 4 above, what sort of problem?

- a) high blood pressure
- b) osteoporosis
- c) stomach cancer
- d) kidney stones
- e) none of the above
- f) all of the above
- g) don't know
- h) refused

6. How important to you is lowering the salt/sodium in your diet?

- a) Not at all important
- b) Somewhat important
- c) Very important

7. Do you do anything on a regular basis to control your salt or sodium intake?

- a) Yes
- b) No (SKIP to QX)
- c) Don't know
- d) Refused

8. If answer is Yes in 7 above, what do you do?

- a) Avoid/minimize consumption of processed foods
- b) Look at the salt or sodium labels on food
- c) Do not add salt at the table
- d) Buy low salt alternatives
- e) Buy low sodium alternatives
- f) Do not add salt when cooking
- g) Use spices other than salt when cooking
- h) Avoid eating out
- i) Other (specify) _____

Section 5: Detailed Instructions for Participants in 24-Hour Urine Collection

We are interested in measuring the dietary intake of certain nutrients – sodium, potassium and iodine. The best way to get this information is by analyzing the urine sample you collect during a 24-hour period.

We cannot get this essential information in any other way!

We are not testing for drugs or viruses.

Your cooperation is very much appreciated.

Why 24 hours?

The content of some nutrients in urine fluctuates according to what we last ate, how much fluid we drink, how much we exercise and also on the weather. Collecting urine over 24 hours gives much more reliable information than a single sample about the typical intakes of these nutrients in a person's diet.

Equipment provided

You have the following equipment provided for making your collections. All equipment is disposable and used only for this study.

1. A sheet to record the essential information about the collection.
2. Urine-collecting equipment for the home:
 - a. 5 litre screw-capped plastic collection bottle to store the collected urine during the day. This bottle contains a preservative for keeping the urine at room temperature.
 - b. a 1 litre plastic jug and funnel for temporal reception of the urine samples.
 - c. a funnel to help women collect urine, which may also help participants in transferring urine samples from the 1-L plastic jug to the 5-L plastic bottle.
 - d. a safety pin (to attach to your underclothes or nightwear simply as a reminder for you to make your collection)
3. Urine-collecting equipment for outside the home:
 - a. a 2 liter screw-capped plastic collection bottle (without preservative)
 - b. two plastic bags for carrying the equipment outside the home



Don't forget to take the jug and 2 liter bottle with you if you leave your home during the day.

Before making the urine collection

The health professional will help you choose the day on which you would like to make the 24-hour urine collection. You may prefer to choose a day when you will be mostly at home or only going out for a short time.

If you are female, you should not make your collection during menstruation.

How to make your collection for the whole day (24 hours)

You have been asked to collect all the urine you pass in one day into the container you have been given. It is not difficult; here is how you do it.

- On the day that you start your collection, you will pass urine – DISCARD this urine, DO NOT put it into the container. Collect from the second time you pass urine. Record the date and time on the Collection Sheet as follows:

Date started Day Month Year
Time started Hour Minutes

- From then onwards until the next day, ALL urine you pass in the next 24 hours, both during the day and night, must be collected.
- The last collection is the urine you pass on the second day at approximately the same time you started the day before.
- This completes the 24-hour collection. Record the following on the Collection Sheet:

Date finished Day Month Year
Time finished Hour Minutes

Note: do not worry if you have not collected for 'exactly' 24 hours, as long as you record exact time of start and finish.

- You should pass all urine directly into the 1 litre plastic jug, then pour the urine into the large container, using the funnel if necessary. If you need to open your bowels, always remember to pass urine first before you pass a stool.
- Each time you add a new urine specimen to the large container, screw the lid tight and swirl the urine around a few times, to mix it with the preservative.
- Any urine collected in the small bottle must be transferred to the large bottle as soon as possible e.g. after returning home.

If you miss a sample

If during the 24-hour collection period a sample is missed for any reason, such as because of a bowel movement, record this on the Urine Collection Sheet.

Once you have completed your collection

As soon as possible after you have completed your 24-hour urine collection, the health professional will arrange a time for him/her to pick up the large container with the total volume of collected urine. In the meantime, store your complete collection in a cool, dark place.

If you have any other questions

We hope this leaflet answers the questions you may have. If you have any other questions, contact the health professional. You are free to withdraw from this study at any point.

Section 6: Household Salt Collection and Iodine Determination

This protocol requires assessments of the iodine content of table and cooking salt. It is therefore important to ask participants for large samples of both types of salt (50-100 gm) where both are used in the household. Because this amount of salt might represent the whole supply in the household, field staff should bring sufficient amounts of both types of salt to replace the samples taken.

In the laboratory, both salt samples should be thoroughly mixed using the same procedure of dry samples to ensure homogeneity. Then, the presence of iodate in the salt should be first identified using a qualitative test kit. For samples that produce a positive reaction (usually a change in color), the quantity of iodine in the samples should then be determined by titration, solubilising not less than 10 gm for refined and small crystal-size salt, and not less than 50 gm for raw or large crystal-size salt. Samples that are negative with the test kit should be analyzed for the quantitative content of iodide using an appropriate method with the same amounts of salt as specified above for the positive samples.

Section 7: Use of Spot Urine to Estimate 24-Hour Excretion of Sodium, Potassium and Iodine

Some researchers have used spot-urine samples to determine the daily excreted amounts of either sodium, potassium or iodine. The sample is only one urine pass collected during the day, frequently not the first pass of the morning made just after awakening (25).

However, the content of sodium, potassium or iodine would depend on the volume of urine, which may be very variable among individuals of the same population, and highly affected by age, sex, ethnic background, weather and body mass index and physical activity. Some “correction” has been proposed by dividing the analyte concentration by the creatinine concentration, based on the fact that creatinine excretion is more constant during the day within an individual, as it mainly depends on lean body mass. However, this correction has been found even less precise than expressing the absolute content by volume, especially in populations with undernutrition.^[26] Although the use of spot-urine is discouraged as a method to determine sodium, potassium or iodine intake because of the limitations and uncertainty inherent in the method, for some populations it may be used to approximate 24-hour excretion of these analytes if a “calibration” is carried out. This “calibration” could be made based on the expected 24-h volume of urine or the 24-h total excretion of creatinine, by applying one of the two following equations:

Approximate 24-h analyte excretion = [analyte] (mg or µg/L) x 24-h urine volume (L) (A)

or

Approximate 24-h analyte excretion = [analyte/Creatinine] (mg or µg/g creatinine) x expected 24-h creatinine excretion (g) (B)

With either equation, the “correction factors” should be calculated in a subsample of individuals from the same population subjected to the same environmental conditions and studied in a 24-hour period. Although equations associated to general parameters, such as body weight and height, age and gender have been published (27,28,29,30), they are specific to certain populations and cannot be reliably extrapolated from one site/population group to another. Thus, in many instances the calculation of these “correction factors” is as difficult as determining directly the 24-hour total excretion of the analytes of interest. Finally, it has been suggested that a spot urine in the afternoon/early evening could provide advantages when compared to a morning one (31). Here, it is important to point out that **even if the above conditions are met, the results are likely to be unreliable especially for population subgroups or time trends. Until more studies are carried out to assess simpler but reliable methods of urine collection for the purpose of estimating daily excretions of these analytes, 24 hour urine collections are recommended.**

Section 8: Dataset for Health Economic Analysis

Chronic disease risk factor variable	Required breakdown				
		25-34	35-44	45-54	55-64
1 Salt intake (NaCl as g per day) Mean	By sex and (adult) age group	Male			
		Female			
2 Smoking (prevalence) Mean	By sex and (adult) age group		25-34	35-44	45-54
			55-64		
3 Systolic blood pressure (mmHg) Mean	By sex and (adult) age group	Male			
		Female			
4 BMI (kg/m2) Mean	By sex and (adult) age group		25-34	35-44	45-54
			55-64		
5 Total blood cholesterol (mmol/L) Mean	By sex and (adult) age group	Male			
		Female			
		Male			
		Female			
			25-34	35-44	45-54
			55-64		
		Male			
		Female			
		Male			
		Female			

References

- Pan American Health Organization.** Regional Strategy and Plan of Action on an Integrated Approach to the Prevention and Control of Chronic Diseases. Washington, DC: PAHO; 2007. Available at: <http://www.paho.org/english/ad/dpc/nc/reg-strat-cncds.pdf>. Accessed 29 April 2010.
- Pan American Health Organization.** Health in the Americas. Volume I. 2002 ed. Washington, DC: PAHO; 2002. Available at: http://www.paho.org/English/DBI/MDS/HIA_2002.htm. Accessed 29 April 2010.
- Law M.** Salt, blood pressure and cardiovascular diseases. *J Cardiovasc Risk* 2000; 7:5-8.
- Penney S.** Dropping the salt: Practical steps countries are taking to prevent chronic non-communicable diseases through population-wide dietary salt reduction. Canadian Public Health Agency, Revised version: August 2009. Available at: <http://www.paho.org/English/AD/dpc/nc/salt-mtg-phac-paper.pdf>. Accessed 29 April 2010.
- He FJ, MacGregor GA.** A comprehensive review on salt and health and current experience of worldwide salt reduction programmes. *J Human Hypertension* 2009; 23:363-84.
- Strazzullo P, D'Elia L, Kandala N-B, Cappuccio FP.** Salt intake, stroke and cardiovascular disease: a meta-analysis of prospective studies. *BMJ* 2009; 339: b4567.
- Sacks FM, Svetkey LP, Vollmer WM et al.** Effects on blood pressure of reduced dietary sodium and the Dietary Approaches to Stop Hypertension (DASH) diet. DASH-Sodium Collaborative Research Group. *New Engl J Med* 2001; 344:3-10.
- Kumanyika SK, Cook NR, Cutler JA, Belden L, Brewer A, Cohen JD, Hebert PR, Lasser VI, Raines J, Raczynski J, Shepek L, Diller L, Whelton PK, Yamamoto M** for the Trials of Hypertension Prevention Collaborative Research Group. Sodium reduction for hypertension prevention in overweight adults: further results from the Trials of Hypertension Prevention Phase II. *J Human Hypertension* 2005; 19:33-45.

9. **Asaria P, Chisholm D, Mathers C, Ezzati M, Beaglehole R.** Chronic disease prevention: health effects and financial costs of strategies to reduce salt intake and control tobacco use. *Lancet* 2007; 370:2044–53.
10. **Murray CJ, Lauer JA, Hutubessy RC, Niessen L, Tomijima N, Rodgers A, Lawes CM, Evans DB.** Effectiveness and costs of interventions to lower systolic blood pressure and cholesterol: a global and regional analysis on reduction of cardiovascular disease risk. *Lancet* 2003; 361:717–25.
11. **Bibbins-Domingo K, Chertow GM, Coxson PG, Moran A, Lightwood JM, Pletcher MJ, Goldman L.** Projected effect of dietary salt reductions on future cardiovascular disease. *N Eng J Med* 2010; 362:590–99.
12. **Pan American Health Organization.** Policy Statement – Reducing dietary salt in the Americas as a population-based approach to cardiovascular disease prevention. 2009. Available at: http://new.paho.org/hq/index.php?option=com_content&task=view&id=2022&Itemid=1766.
13. **MRC – Human Nutrition Research, National Centre for Social Research.** An assessment of dietary sodium levels among adults (aged 19–64) in the UK general population in 2008, based on analysis of dietary sodium in 24 hour urinary sodium samples. June 2008. Available at: <http://www.food.gov.uk/multimedia/pdfs/08sodiumreport.pdf>. Accessed 8 April 2010.
14. **Anderson CAM, Appel LJ, Okuda N, Brown IJ, Chan Q, Zhao L, Ueshima H, Kesteloot H, Miura K, Curb JD, Yoshita K, Elliott P, Yamamoto ME, Stamler J.** Dietary Sources of Sodium in China, Japan, the United Kingdom, and the United States, Women and Men Aged 40 to 59 Years: The INTERMAP Study. *J Am Dietetic Association* 2010; 110:736–45.
15. **European Commission.** EU Framework for National Salt Initiatives. Available at: http://ec.europa.eu/health/nutrition_physical_activity/high_level_group/nutrition_salt_en.htm. Accessed 20 May 2010.
16. **Choices International Foundation.** The Choices Programme. Available at: http://choicesprogramme.org/en/about_the_choices_programme/product_criteria. Accessed 21 May 2010.
17. **Elliott P, Stamler J, Nichols R, Dyer AR, Stamler R, Kesteloot H, Marmot M.** Intersalt revisited: further analyses of 24 hour sodium excretion and blood pressure within and across populations. *BMJ* 1996; 312:1249–53.
18. **Khaw K-T, Barret-Connor E.** Dietary potassium and stroke-associated mortality. A 12-year prospective population study. *N Engl J Med* 1987; 316:235–40.
19. **Cappuccio FP, MacGregor GA.** Does potassium supplementation lower blood pressure? A meta-analysis of published trials. *J Hypertens* 1991; 9:465–73.
20. **Siani A, Strazzullo P, Giacco A, Pacioni D, Celentano E, Mancini M.** Increasing the Dietary Potassium Intake Reduces the Need for Antihypertensive Medication. *Ann Intern Med* 1991; 115:753–9.
21. **Andersson M, de Benoist B, Rogers L.** Epidemiology of iodine deficiency: Salt iodisation and iodine status. *Best Pract Res Clin Endocrinol Metab* 2010; 24:1–11.
22. **Pan American Version of the WHO STEPSwise Approach to Chronic Disease Risk Factor Surveillance.** Available at: www.paho.org/English/AD/DPC/NC/panam-steps.htm. Accessed 20 May 2010.
23. **World Health Organization.** Assessment of iodine deficiency disorders and monitoring their elimination: a guide for program managers. Third edition (updated 1st September 2008). Available at: http://www.who.int/nutrition/publications/micronutrients/iodine_deficiency/9789241595827/en/index.html. Accessed 20 May 2010.
24. **Campbell NRC, McKay DW, Chockalingam A, Fodor JG.** Errors in Assessment of Blood Pressure: Sphygmomanometers and Blood Pressure Cuffs. *Can J Public Health* 1994; Supplement 2: S22–S25.
25. **Rasmussen LB, Ovesen L, Christiansen E.** Day-to-day and within-day variation in urinary iodine excretion. *Eur J Clin Nutr* 1999; 53:401–7.
26. **Furnée CA, van der Haar F, West CE, Hautvast JGAJ.** A critical appraisal of goiter assessment and the ratio of urinary iodine to creatinine for evaluating iodine status. *Am J Clin Nutr* 1994; 59:1415–17.
27. **IOM, National Academy of Sciences of the USA.** Dietary Reference Intakes, Iodine. 2000:258–89. Available at: http://www.nap.edu/openbook.php?record_id=10026&page=258. Accessed 29 April 2010.
28. **Tanaka T, Okamura T, Miura K et al.** A simple method to estimate populational 24-h urinary sodium and potassium excretion using a casual urine specimen. *J Hum Hypertens* 2002; 16:97–103.
29. **Kawano Y, Tsuchihashi T, Matsuura H, Ando K, Fujita T, Ueshima H.** Report of the Working Group for dietary salt reduction of the Japanese Society of Hypertension: (2) Assessment of salt intake in the management of hypertension. *Hypertens Res* 2007; 30:887–93.
30. **Knudsen N, Christiansen E, Brandt-Christensen M, Nygaard B, Perrild H.** Age- and sex-adjusted iodine/creatinine ratio. A new standard in epidemiological surveys? Evaluation of three different estimates of iodine excretion based on casual urine samples and comparison to 24 h values. *Eur J Clin Nutr* 2000; 54:361–63.
31. **Mann SJ, Gerber LM.** Estimation of 24-hour sodium excretion from spot urine samples. *J Clin Hypert* 2010; 12:174–80.

Acknowledgements

Norm Campbell

Lubin Cardiovascular Institute of Alberta, University of Calgary, Canada.

Francesco Cappuccio

WHO Collaborating Centre for Nutrition, Warwick University, England.

Anselm Hennis

Chronic Disease Research Centre, Barbados.

Simon Barquera

Institute of Public Health, Cuernavaca, Mexico.

Ricardo Correa Rotter

National Medical Science and Nutrition Institute Salvador Zubiran, Mexico.

Omar Dary

Academy of Educational Development, Washington DC, USA.

Rainford Wilks

University of West Indies, Kingston, Jamaica.

Daniel Ferrante

Health Promotion and Chronic Disease Control, Ministry of Health, Argentina.

Roxana Buscaglione

Department of Chronic Diseases, Ministry of Health, Chile.

Barbara Legowski

PAHO Secretariat, Washington DC, USA.

Branka Legetic

PAHO Secretariat, Washington DC, USA.

PAHO/WHO Sodium Detection in 24-Hour Urine Samples Questionnaire

Questionnaire Information

Location and Date		Response
1	Center/City name	
2	Interviewer ID	
3	Date of completion of the instrument (dd/mm/yyyy)	
Consent, Interview Language and Name		Response
4	Consent has been read and obtained	Yes <input type="checkbox"/> No <input type="checkbox"/> If NO,END
5	Interview Language <i>[insert language]</i>	English <input type="checkbox"/> <i>[add others]</i> <input type="checkbox"/> <i>[add others]</i> <input type="checkbox"/> <i>[add others]</i> <input type="checkbox"/>
6	Time of Interview (24 hour clock)	Hrs: mins
7	Family Surname	
8	First Name	
Additional Information that may be helpful		
9	Contact phone number where possible	

1 - Demographic Information

CORE: Demographic Information		
Question	Response	
10	Sex (<i>Record Male/Female as observed</i>)	Male <input type="checkbox"/> Female <input type="checkbox"/>
11	What is your date of birth? <i>Don't know (leave blank)</i>	/ / dd/mm/yyyy
12	How old are you?	years
EXPANDED: Demographic Information		
Question	Response	
13	What is the highest level of education you have completed	Not formal schooling <input type="checkbox"/> Less than primary school <input type="checkbox"/> Primary school completed <input type="checkbox"/> Secondary school completed <input type="checkbox"/> High school completed <input type="checkbox"/> College/university completed <input type="checkbox"/> Post graduate degree <input type="checkbox"/> Refused <input type="checkbox"/>
14	What is your <i>[insert relevant ethnic group /racial group/cultural subgroup /others]</i> background?	<i>[locally defined]</i> <input type="checkbox"/> <i>[locally defined]</i> <input type="checkbox"/> <i>[locally defined]</i> <input type="checkbox"/> <i>[locally defined]</i> <input type="checkbox"/> <i>[locally defined]</i> <input type="checkbox"/> Refused <input type="checkbox"/>

15	What is your marital status?	Never married <input type="checkbox"/>
		Currently married <input type="checkbox"/>
		Separated <input type="checkbox"/>
		Divorced <input type="checkbox"/>
		Widowed <input type="checkbox"/>
		Cohabiting <input type="checkbox"/>
		Refused <input type="checkbox"/>
16	Which of the following best describes your main work status over the past 12 months? <i>[insert country-specific categories]</i>	Government employee <input type="checkbox"/>
		Non-government employee <input type="checkbox"/>
		Self-employed <input type="checkbox"/>
		Non-paid <input type="checkbox"/>
		Student <input type="checkbox"/>
		Homemaker <input type="checkbox"/>
		Retired <input type="checkbox"/>
		Unemployed (able to work) <input type="checkbox"/>
		Unemployed (unable to work) <input type="checkbox"/>
		Refused <input type="checkbox"/>
17	Taking the past year, can you tell me what the average earnings of the household have been? <i>[record only one, not all three]</i>	Per week <input type="checkbox"/>
		OR per month <input type="checkbox"/>
		OR per year <input type="checkbox"/>
		Refused <input type="checkbox"/>

1 – Behavioral Measurements

CORE: Tobacco Use

Now I'm going to ask you some questions about various health behaviors. This includes things like smoking, drinking alcohol, eating fruits, and vegetables and physical activity. Let's start with tobacco.

Question	Response
18 Do you currently smoke any tobacco product, such as cigarettes, cigars or pipes?	Yes <input type="checkbox"/> No <input type="checkbox"/>
19 Do you currently smoke tobacco products daily?	Yes <input type="checkbox"/> No <input type="checkbox"/>
20 How old were you when you first started smoking daily?	Age (years) <input type="checkbox"/> Don't know <input type="checkbox"/>
21 On average, how many of the following do you smoke each day? <i>[record for each type; if don't know leave blank]</i>	Manufactured cigarettes <input type="checkbox"/>
	Hand-rolled cigarettes <input type="checkbox"/>
	Pipes full of tobacco <input type="checkbox"/>
	Cigars, cheroots, cigarillos <input type="checkbox"/>
	Other <input type="checkbox"/>

EXPANDED: Tobacco Use

Question	Response
22 In the past, did you ever smoke daily?	Yes <input type="checkbox"/> No <input type="checkbox"/>
23 How old were you when you stopped smoking daily?	Age (years) <input type="checkbox"/> Don't know <input type="checkbox"/>
24 Do you currently use any smokeless tobacco such as <i>[snuff, chewing tobacco, betel]</i> ?	Yes <input type="checkbox"/> No <input type="checkbox"/>
25 On average, how many times a day do you use <i>[record for each type; if don't know leave blank]</i>	Snuff, by mouth <input type="checkbox"/>
	Snuff, by nose <input type="checkbox"/>
	Chewing of tobacco <input type="checkbox"/>
	Betel, quid <input type="checkbox"/>
	Other <input type="checkbox"/>
26 In the past, did you ever use smokeless tobacco such as <i>[snuff, chewing tobacco, or betel]</i> daily?	Yes <input type="checkbox"/> No <input type="checkbox"/>

CORE: Alcohol Consumption

The next questions, ask you about the consumption of alcohol

Question	Response
27 Have you ever consumed any alcoholic drinks such as beer, wine, spirits, fermented cider, or <i>[add other local examples]</i> ?	Yes <input type="checkbox"/> No <input type="checkbox"/> <i>If NO, go to Diet Core Questions page 5</i>
28 Have you consumed an alcoholic drink within the past 12 months?	Yes <input type="checkbox"/> No <input type="checkbox"/>
29 During the past 12 months, how frequently have you had at least one alcoholic drink?	Daily <input type="checkbox"/> 5-6 days per week <input type="checkbox"/> 1-4 days per week <input type="checkbox"/> 1-3 days per month <input type="checkbox"/> Less than once a month <input type="checkbox"/>
30 Have you consumed an alcoholic drink within the past 30 days?	Yes <input type="checkbox"/> No <input type="checkbox"/> <i>If NO, go to Diet Core Questions page 5</i>
31 During the past 30 days, on how many occasions did you have at least one alcoholic drink?	Number <input type="checkbox"/> Don't know <input type="checkbox"/>
32 During the past 30 days, when you drank alcohol, on average, how many standard alcoholic drinks did you have during one drinking occasion?	Number <input type="checkbox"/> Don't know <input type="checkbox"/>
33 During the past 30 days, what was the largest number of standard alcoholic drinks you had on a single occasion, counting all types of alcoholic drinks together?	Largest number <input type="checkbox"/> Don't know <input type="checkbox"/>
34 During the past 30 days, how many times did you have for men: five or more For women: four or more Standard alcoholic drinks in a single drinking occasion?	Number of times <input type="checkbox"/> Don't know <input type="checkbox"/>

EXPANDED: Alcohol Consumption

Question	Response
35 During the past 30 days, when you consumed an alcoholic drink, how often was it with meals? Please do not count snacks	Usually with meals <input type="checkbox"/> Sometimes with meals <input type="checkbox"/> Rarely with meals <input type="checkbox"/> Never with meals <input type="checkbox"/>
36 During each of the past 7 days, how many standard alcoholic drinks did you have each day?	Monday <input type="checkbox"/> <input type="checkbox"/> Tuesday <input type="checkbox"/> <input type="checkbox"/> Wednesday <input type="checkbox"/> <input type="checkbox"/> Thursday <input type="checkbox"/> <input type="checkbox"/> Friday <input type="checkbox"/> <input type="checkbox"/> Saturday <input type="checkbox"/> <input type="checkbox"/> Sunday <input type="checkbox"/> <input type="checkbox"/> Don't know <input type="checkbox"/>

CORE: Diet

The next questions ask about the fruits and vegetables that you usually eat.

Question	Response
37 In a typical week, on how many days do you eat fruit?	Number of days <input type="checkbox"/> <input type="checkbox"/> Don't know <input type="checkbox"/>
38 How many servings of fruit do you eat on one of those days?	Number of days <input type="checkbox"/> <input type="checkbox"/> Don't know <input type="checkbox"/>
39 On a typical week, on how many days do you eat vegetables?	Number of days <input type="checkbox"/> <input type="checkbox"/> Don't know <input type="checkbox"/>
40 How many servings of vegetables do you eat on one of those days?	Number of days <input type="checkbox"/> <input type="checkbox"/> Don't know <input type="checkbox"/>

EXPANDED: Diet		
Question	Response	
41	What type of oil or fat is most often used for meal preparation in your household? <i>[select only one]</i>	Vegetable oil <input type="checkbox"/> Lard or suet <input type="checkbox"/> Butter or ghee <input type="checkbox"/> Margarine <input type="checkbox"/> Other <input type="checkbox"/> None in particular <input type="checkbox"/> None used <input type="checkbox"/> Don't know <input type="checkbox"/> Other ----- <input type="checkbox"/>
42	On average, how many meals per week do you eat that were not prepared at a home? By meal, I mean breakfast, lunch and dinner.	Number <input type="checkbox"/> <input type="checkbox"/> Don't know <input type="checkbox"/>
Diet: Knowledge, Attitudes, and Behavior towards Dietary Salt		
The next set of questions is about the knowledge, attitudes and behavior towards dietary salt. Please answer the following even you consider yourself to eat a low sodium diet.		
Question	Response	
43	Do you add salt to food at the table? <i>[select only one]</i>	Never <input type="checkbox"/> Rarely <input type="checkbox"/> Sometimes <input type="checkbox"/> Often <input type="checkbox"/> Always <input type="checkbox"/>
44	In the food you eat at home salt is added in cooking...	Never <input type="checkbox"/> Rarely <input type="checkbox"/> Sometimes <input type="checkbox"/> Often <input type="checkbox"/> Always <input type="checkbox"/>
45	How much salt do you think you consume? <i>[select only one]</i>	Far too much <input type="checkbox"/> Too much <input type="checkbox"/> Just the right amount <input type="checkbox"/> Too little <input type="checkbox"/> Far too little <input type="checkbox"/> Don't know <input type="checkbox"/> Refuse <input type="checkbox"/>
46	Do you think that a high salt diet could cause a serious health problem?	Yes <input type="checkbox"/> No <input type="checkbox"/> Don't know <input type="checkbox"/> Refused <input type="checkbox"/> <i>If NO, DON'T KNOW or REFUSED, go to Q 48</i>
47	What sort of serious health problems do you think a high salt diet could cause? <i>[mark as many as it applies]</i>	High blood pressure <input type="checkbox"/> Osteoporosis <input type="checkbox"/> Stomach cancer <input type="checkbox"/> Kidney stones <input type="checkbox"/> None of the above <input type="checkbox"/> All of the above <input type="checkbox"/> Don't know <input type="checkbox"/> Refused <input type="checkbox"/>
48	How important to you is lowering the salt/sodium in your diet?	Not at all important <input type="checkbox"/> Somewhat important <input type="checkbox"/> Very Important <input type="checkbox"/>
49	Do you anything on a regular basis to control your salt or sodium intake?	Yes <input type="checkbox"/> No <input type="checkbox"/> Don't know <input type="checkbox"/> Refused <input type="checkbox"/> <i>If NO, DON'T KNOW or REFUSED, go to Q 51</i>

50	What do you do on a regular basis to control your salt or sodium intake? <i>[mark as many as it applies]</i>	Avoid/minimize consumption of processed foods <input type="checkbox"/> Look at the salt or sodium labels on food <input type="checkbox"/> Do not add salt at the table <input type="checkbox"/> Buy low salt alternatives <input type="checkbox"/> Buy low sodium alternatives <input type="checkbox"/> Do not add salt when cooking <input type="checkbox"/> Use spices other than salt when cooking <input type="checkbox"/> Avoid eating out <input type="checkbox"/> Other (specify) _____
----	---	--

CORE: Physical Activity

The next questions ask about the time you spend doing different types of physical activity in a typical week. Please answer these questions even if you don't consider yourself to be a physically active person.

Think about the time you spend doing work. This includes paid and unpaid work, like study/training, household chores, harvesting food/crops, fishing or hunting for food, seeking employment, etc.

Remember that when answering the following questions:

- Vigorous-intensity activities require hard physical effort and cause large increases in breathing or heart rate
- Moderate-intensity activities require moderate physical effort and cause small increases in breathing or heart rate.

Question	Response
Work	
51	Does your work involve vigorous-intensity activity that causes large increases in breathing or heart rate like <i>[carrying or lifting heavy loads, digging or construction work]</i> for at least 10 minutes continuously? Yes <input type="checkbox"/> No <input type="checkbox"/>
52	In a typical week, on how many days do you do vigorous-intensity activities as part of your work? <input type="text"/> <input type="text"/>
53	How much time do you spend doing vigorous-intensity activities at work on a typical day? Hours: minutes _____
54	Does your work involve moderate-intensity activity that causes small increases in breathing or heart rate such as brisk walking <i>[or carrying light loads]</i> for at least 10 minutes continuously? <i>[insert examples]</i> Yes <input type="checkbox"/> No <input type="checkbox"/>
55	In a typical week, on how many days do you do moderate-intensity activities as part of your work? <input type="text"/> <input type="text"/>
56	How much time do you spend doing moderate-intensity activities at work on a typical day? Hours: minutes _____

Travel to and from places

The next questions exclude the physical activities at work that you have already mentioned. These questions are about the usual way you travel to and from places. For example to work, for shopping, to the market, to place of worship, etc.

57	Do you walk or use a bicycle (pedal cycle) for at least 10 minutes continuously to get to and from places? Yes <input type="checkbox"/> No <input type="checkbox"/>
58	In a typical week, on how many days do you walk or bicycle for at least 10 minutes continuously to get to and from places? <input type="text"/> <input type="text"/>
59	How much time do you spend walking or bicycling for travel on a typical day? Hours: minutes _____

Leisure activity

The next questions exclude the work and transport activities that you have already mentioned. Now I would like to ask you about sports, fitness, and recreational activities (leisure), etc.

60	Do you do any vigorous-intensity sport, fitness or recreational (leisure) activities that cause large increases in breathing or heart rate like <i>[running or football]</i> for at least 10 minutes continuously? Yes <input type="checkbox"/> No <input type="checkbox"/>
61	In a typical week, on how many days do you do vigorous-intensity sports, fitness or recreational (leisure) activities? <input type="text"/> <input type="text"/>

62	How much time do you spend doing vigorous-intensity sports, fitness or recreational (leisure) activities on a typical day?	Hours: minutes
63	Do you do any moderate-intensity sports, fitness or recreational (leisure) activity that causes small increases in breathing or heart rate such as brisk walking [<i>cycling, swimming, or volleyball</i>] for at least 10 minutes continuously? <i>[insert examples]</i>	Yes <input type="checkbox"/> No <input type="checkbox"/>
64	In a typical week, on how many days do you do moderate-intensity activities as part of your work?	Number of days <input type="checkbox"/>
65	How much time do you spend doing moderate-intensity sports, fitness or recreational (leisure) activities on a typical day?	Hours: minutes

Travel to and from places

The next questions exclude the physical activities at work that you have already mentioned. These questions are about the usual way you travel to and from places. For example to work, for shopping, to the market, to place of worship, etc.

66	Do you walk or use a bicycle (pedal cycle) for at least 10 minutes continuously to get to and from places?	Yes <input type="checkbox"/> No <input type="checkbox"/>
67	In a typical week, on how many days do you walk or bicycle for at least 10 minutes continuously to get to and from places?	Number of days <input type="checkbox"/>
68	How much time do you spend walking or bicycling for travel on a typical day?	Hours: minutes

EXPANDED: Physical Activity

Sedentary Behavior

These questions are about the amount of time you spend sitting or reclining at work, at home, getting to and from places, or with friends including time spend sitting at the desk, sitting with friends, traveling in car, bus, train, reading, playing cards or watching television, but do not include time spent sleeping.

Question	Response	
69	How much time do you usually spend sitting or reclining on a typical day	Hours: minutes

2 – Personal Medical History

CORE: Personal Medical History

Question	Response	
70	Have you ever been told by a medical doctor that you have, or have had, heart failure ?	Yes <input type="checkbox"/> No <input type="checkbox"/>
71	Have you ever been told by a medical doctor that you have, or have had, heart attack ?	Yes <input type="checkbox"/> No <input type="checkbox"/>
72	Have you ever been told by a medical doctor that you have, or have had, other heart trouble ?	Yes <input type="checkbox"/> If yes, please specify No <input type="checkbox"/>
73	Have you ever been told by a medical doctor that you have had a stroke ?	Yes <input type="checkbox"/> No <input type="checkbox"/>
74	Have you ever been told by a medical doctor that you have, or have had, kidney trouble ?	Yes <input type="checkbox"/> No <input type="checkbox"/>
75	Have you ever been told by a medical doctor that you have, or have had, peptic ulcer ?	Yes <input type="checkbox"/> No <input type="checkbox"/>
76	Have you ever been told by a medical doctor that you have, or have had, liver disease ?	Yes <input type="checkbox"/> No <input type="checkbox"/>
77	Have you ever been told by a medical doctor that you have a cancer or malignant tumor ?	Yes <input type="checkbox"/> No <input type="checkbox"/>

CORE: History of Raised Blood Pressure		
Question	Response	
78	Have you ever had your blood pressure measured by a doctor or other health workers?	Yes <input type="checkbox"/> No <input type="checkbox"/> <i>If NO, go to Q 67</i>
79	Have you ever been told by a doctor or other health worker that you have raised blood pressure or hypertension?	Yes <input type="checkbox"/> No <input type="checkbox"/> <i>If NO, go to Q 67</i>
80	Have you been told in the past 12 months?	Yes <input type="checkbox"/> No <input type="checkbox"/>
81	Are you currently receiving any of the following treatments/advice for high blood pressure prescribed by a doctor or other health worker?	
	Drugs (medication) that you have taken in the past 2 weeks?	Yes <input type="checkbox"/> No <input type="checkbox"/>
	Advice to reduce salt intake?	Yes <input type="checkbox"/> No <input type="checkbox"/>
	Advice or treatment to lose weight?	Yes <input type="checkbox"/> No <input type="checkbox"/>
	Advice or treatment to stop smoking?	Yes <input type="checkbox"/> No <input type="checkbox"/>
	Advice to start or do more exercise?	Yes <input type="checkbox"/> No <input type="checkbox"/>
82	Have you ever seen a traditional healer for raised blood pressure or hypertension?	Yes <input type="checkbox"/> No <input type="checkbox"/>
83	Are you currently taking any herbal or traditional remedy for your raised blood pressure?	Yes <input type="checkbox"/> No <input type="checkbox"/>
CORE: History of Diabetes		
84	Have you ever had your blood sugar measured by a doctor or other health workers?	Yes <input type="checkbox"/> No <input type="checkbox"/>
85	Have you ever been told by a doctor or other health worker that you have raised blood sugar or diabetes?	Yes <input type="checkbox"/> No <input type="checkbox"/>
86	Have you been told in the past 12 months?	Yes <input type="checkbox"/> No <input type="checkbox"/>
87	Are you currently receiving any of the following treatments/advice for diabetes prescribed by a doctor or other health worker?	
	Insulin	Yes <input type="checkbox"/> No <input type="checkbox"/>
	Drugs (medication) that you have taken in the past 2 weeks?	Yes <input type="checkbox"/> No <input type="checkbox"/>
	Special prescribed diet?	Yes <input type="checkbox"/> No <input type="checkbox"/>
	Advice or treatment to lose weight?	Yes <input type="checkbox"/> No <input type="checkbox"/>
	Advice or treatment to stop smoking?	Yes <input type="checkbox"/> No <input type="checkbox"/>
	Advice to start or do more exercise?	Yes <input type="checkbox"/> No <input type="checkbox"/>
88	Have you ever seen a traditional healer for raised blood glucose or diabetes?	Yes <input type="checkbox"/> No <input type="checkbox"/>
89	Are you currently taking any herbal or traditional remedy for your raised blood glucose or diabetes?	Yes <input type="checkbox"/> No <input type="checkbox"/>

3 – Physical Measurements

CORE: Height and Weight		
Question	Response	
90	Device ID for height and weight	Height device ___ ___ Weight device ___ ___
91	Height	In Centimeters (cm) ___ . ___
92	Weight <i>If too large for scale, write 666.6</i>	In Kilograms (kg) ___ . ___
93	For women: Are you pregnant?	Yes <input type="checkbox"/> No <input type="checkbox"/>
CORE: Waist		
94	Device ID for waist	Device ___ ___
95	Waist circumference	In Centimeters (cm) ___ . ___
CORE: Blood Pressure		
96	Device ID for blood pressure	Device ___ ___
97	Cuff size used	___ ___
98	Reading 1	Systolic (mmHg) ___ ___ Diastolic (mmHg) ___ ___
99	Reading 2	Systolic (mmHg) ___ ___ Diastolic (mmHg) ___ ___
100	Reading 3	Systolic (mmHg) ___ ___ Diastolic (mmHg) ___ ___
EXPANDED: Physical Measurements		
Question	Response	
101	Hip circumference	In Centimeters (cm) ___ . ___
102	Heart rate	
	Reading 1	Beats per minute ___ ___
	Reading 2	Beats per minute ___ ___
	Reading 3	Beats per minute ___ ___

4 – 24-Hour Urine Sample

24-Hour Urine Samples Collection

[At this point, the participant will be given the “Participant’s Guide for the 24-Hour Urine Samples Collection. The participant will be guided through the guide and choose a day to begin the collection. The Interviewer will then make an appointment to pick up the samples within 24 hours of then end of collection. These questions will be questions obtained from the log sheet given to the participant, after the 24-hour urine sample collection. It’s recommended that it be filled in with the participant present.] Now I will like to review a few your log sheet and go over any comments or concerns you might have.

Question	Response	
103	Device ID for 24–hour urine sample collection <i>[each equipment set must be labeled with participant’s ID; therefore, the Participant ID and Device ID are the same]</i>	
104	Date collection began	dd/mm/yyyy ___/___/_____
105	Time collection started	hh:mm ___:___
106	Date collection finished	dd/mm/yyyy ___/___/_____
107	Time collection finished	hh:mm ___:___
108	Total volume of urine collected	In milliliters (ml) ___ ___

**A Review of Methods to
Determine the Main Sources
of Salt in the Diet**

Section 1: Introduction

For countries to effectively direct their policies and interventions at lowering dietary salt, they need to monitor how much salt people are eating and identify the main food sources of salt. To determine total salt intake, there is a Protocol for Population Level Sodium Determination in 24-hour Urine Samples, prepared by the PAHO/WHO Regional Expert Group for Cardiovascular Disease Prevention through Population-wide Dietary Salt Reduction. For the Protocol, see http://new.paho.org/hq/index.php?option=com_content&task=view&id=3072&Itemid=2376.

This Review is the companion to the Protocol, intended to assist governments and other public health agencies in determining the main sources of dietary salt, to inform their decisions on how best to reduce salt intake. The description of methods is a compilation of general information and the specific experiences of a number of nutrition science experts in North, Central and South America and the Caribbean, identified in the list of acknowledgements.

Primary aims

The Review features the combination of methods and data sources that will deliver a complete profile of the dietary sources of salt by identifying:

- foods that people eat and the amounts and frequency of consumption;
- sodium content of the most commonly consumed foods;
- the amount of salt added at the table and in cooking.
- intake of high-sodium foods that are culturally or regionally-specific.

The variety of methods employed to assess the sources of salt in the diet reflect the complexity and dynamic nature of food supplies. This Review is designed to assist countries in selecting what is appropriate to their resources and circumstances. Once baseline data on the of sources and amounts of dietary salt are determined, ongoing monitoring is needed to keep up with changes e.g. the introduction of new and reformulated food products and population shifts in eating habits.

Other factors to consider

- Data on several nutrients of concern to public health nutrition policy, in addition to sodium, can be captured with the methods described here, for example, saturated fats, trans fatty acids, fibre and total sugar.
- If the individual food products in food consumption surveys and in the food composition tables are grouped into broad categories such as breads, processed meats, etc, they can become

the basis for raising awareness among consumers as to the food categories that contribute the most salt to the diet.

- Grouping food products according to common formulations, functions or processes can also serve as a useful basis for setting reduced-sodium targets with the food industry.
- Questions on knowledge, attitudes and behaviour (KAB) can be added to a food consumption survey to gain more information about consumers' perspectives on salt that can inform interventions and communication strategies. A sample set of KAB questions is provided in the Protocol mentioned above.
- Countries with fairly common food cultures in a particular region can explore opportunities for joint approaches or harmonized tools that may provide economies of scale to determining the main sources of salt in diets.

Intended audience

Governments, public health agencies and principle investigator(s) for studies of nutritional habits to determine the sources of sodium in the diet.

Section 2: Food Consumption Surveys

Overview

Ideally, a national food consumption survey is cross-sectional in nature, capturing food consumption data from which nutrient intakes are calculated as well as collecting height/weight measurements of respondents and sometimes their nutritional and health status. Below are considerations when designing a nutrition survey.

Who to survey

Which ages – infants, children, adolescents, adults, elderly people.

What to survey

All foods consumed by the population whether at home, in restaurants, canteens, cafeterias in schools or workplaces, etc. (depends on who is being surveyed and where they eat).

Possible exclusions

People in remote locations, the military, institutions such as nursing homes, hospitals etc. (depends on who is being surveyed)

Representativeness

- Nationally (and/or state/provincially) represen-

tative, or representative sentinel sites, based on census data for each age/gender group (groups should be consistent with age groups used for national nutritional recommendations).

- Where census data are lacking, different sampling methods can be applied such as convenience samples in urban and rural settings.

Other data to collect

May want to collect information on selected health conditions, socioeconomic or demographic characteristics.

When and how often to survey

- Ideally continuously with national commitment for cycle repeats e.g. every 3, 5, 10 or 20 years, or
- Periodically as national resources permit.

General challenges with surveys

Need to systematically use probing questions and if possible observe existing foods and preparations, to ensure that respondents accurately record all foods eaten and amounts

- Need to measure serving sizes, portions eaten (and frequency of consumption if applicable).
- Need to calculate ingredients and amounts in recipes.
- Consumption of mixed dishes can make it difficult for participants to recall and assess items and amounts of food consumed during a meal.
- Store-bought versus restaurant versus home-made foods – are not always the same in terms of ingredients, serving size, or nutrients.
- Anticipate specific challenges when surveying young children and the frail elderly.
 - need to question parents, daycares, caregivers.
 - often not sure how much is served versus eaten.
- Households eating from a communal pot during meals can make it difficult to assess food consumption of individuals.
- Some diets may include complex stews and soups, in which water and extra ingredients are added to an existing dish over a day or more, and portions are taken as needed.
- Can be difficult to estimate sodium losses with some food preparation practices e.g. soaking or rinsing of foods prior to preparation or discarding salted cooking water (1).
- Food terminology can vary.
- Household measures can vary.
- Foods consumed must be linked to relevant

food composition databases/tables to determine sodium content (see Section 3).

Challenges specific to accounting for foods with salt/sodium additives

- Once changes and reformulations start occurring, the composition of similar food products may vary greatly between manufacturers.
- In some circumstances, a variety of sodium additives can be used interchangeably.

Methods that Provide Direct or Primary Data on Food Consumption

A number of methods can provide direct or primary data on food consumption, requiring participants to provide personal information on what they typically eat and drink in a defined period of time (2,3,4). The main methods used currently for dietary assessment are described below with examples following.

Descriptions and considerations

24-hour food intake recall

A systematic questionnaire/survey designed to capture all foods consumed in a defined 24-hour period.

- Ideal method commonly used by government agencies for national surveys.
- Foods with a certain threshold of intake and sodium content per serving can be added (15).
- A number of governments have extensive experience with this survey methodology (see below the US National Health and Nutrition Examination).
- Survey (NHANES) and the Canadian Community Health Survey (CCHS) 2.2) may be expensive.
- Requires considerable analysis to categorize foods and extensive follow-up questions to ensure complete recalls.
- May not be representative of usual consumption patterns if limited to a single day recall.
- Distribution of single day intakes shows larger variance than usual intake distribution; requires a repeat recall on a different day (ideally a week day and a weekend day) from a sub-sample of respondents to estimate usual intakes.
- Difficult to model usual intake for episodically consumed foods e.g. seasonally available.
- Can have large variations by geographic area e.g. rural versus urban statistical challenges: need to account for days without food consumption; allow for consumption-day amounts that are positively skewed.
- Recent improvements include computer assisted questionnaires (see the multi-pass method below).

Food frequency questionnaires (FFQ)

Participants indicate their usual dietary intake – how frequently certain foods and food groups on a pre-defined list are consumed during a specified period of time.

- Suitable for population surveys.
- Relatively inexpensive.
- Most amenable to web-based administration.
- recent improvements include a number of on-line and web-based versions which are often easier and quicker to complete.
- Can capture intake of all nutrients of interest to national nutrition policy and can inform on overall dietary adequacy.
- Can be expanded or contracted to include or isolate high salt/sodium food items.
- Can add questions on salt added at table or in cooking.
- Can add questions on respondents' willingness to reduce dietary sodium.
- Can add questions on consumption away from home to monitor the nutrition transition.
- In lower resource situations, can consider sentinel food surveillance.
- Usually requires validation food diaries done in a sub-sample of the same population.
- Represents usual nutrient intake, as diet is assessed over long periods of time (e.g. previous 12 months).
- Field staff require extensive training to ensure data consistency.
- Are typically long because of the need to be comprehensive of common food products.

3-day, 7-day food diaries

Written records of all foods and beverages consumed and amounts over 3 or 7 days.

Duplicate food collection or weighed food consumption

The preparation of two identical plates of every food consumed one of which is sent to a laboratory for chemical analysis of nutrient content.

Accurate weighing of all foods consumed combined with analysis or calculation of nutrient composition.

- Usually used only for research projects.
- Very time consuming with heavy burden on participants.
- Relatively expensive.

Example 1: US NHANES 24-hour food recall

NHANES is a program of studies designed to assess the health and nutrition status of adults and children in the United States, combining interviews with physical examinations. For survey questionnaires, exami-

nation components and laboratory components for 2009-2010, see http://www.cdc.gov/nchs/nhanes/nhanes2009-2010/questexam09_10.htm. A dietary screener module is available at http://www.cdc.gov/nchs/data/nhanes/nhanes_09_10/mi_dtq_f.pdf and a dietary data tutorial is available at <http://www.cdc.gov/nchs/tutorials/dietary/index.htm> to promote broader and more proficient use of NHANES data.

NES also includes some FFQ modules. English and Spanish versions of the NHANES FFQ are available at <http://riskfactor.cancer.gov/diet/usualintakes/ffq.html>

Example 2: CCHS 2.2 24-Hour food recall

The survey provides national nutrition data about the food and nutrient intakes of Canadians and the relationship between diet and a wide range of health correlates. The 24-hour recall instrument is described at <http://www.statcan.gc.ca/cgi-bin/imdb/p2SV.pl?Function=getSurvey&SDDS=5049&lang=en&db=imdb&adm=8&dis=2>.

The last CCHS survey also included a series of short questions on health, demographics and socio-economic status, and had several targeted FFQs, for example on use of salt, supplements, fruit and vegetable consumption, etc. A guide to help with assessing and interpreting the data is available at http://www.hc-sc.gc.ca/fn-an/alt_formats/hpfb-dgpsa/pdf/surveill/cchs-guide-escs-eng.pdf

Example 3: Automated multiple-pass method for 24-Hour recall

Automated multi-pass method (AMPM) is a computerized method for systematically collecting interview-administered 24-hour dietary recalls either in person or by telephone. The field staff follow five research-based steps designed to enhance complete and accurate food recall and reduce respondent burden (see below). Information about the AMPM is available at the USDA web site – <http://www.ars.usda.gov/Services/docs.htm?docid=7710>.

Step 1	Quick List <ul style="list-style-type: none">• Everything you had to eat or drink yesterday
Step 2	Forgotten Foods <ul style="list-style-type: none">• List of foods often forgotten e.g. coffee, tea, soft drinks, milk, juice, water
Step 3	Time and Occasion <ul style="list-style-type: none">• Breakfast, lunch, dinner, snacks
Step 4	Details <ul style="list-style-type: none">• Types – e.g. bread (white, brown, rye, etc)• Amounts – number, size of unit (use pictures, models)
Step 5	Final Probe <ul style="list-style-type: none">• Anything else, even small amounts at e.g. meetings, while shopping, cooking, etc.

Sample Screen Viewed by Field Staff

Global Data Entry - C:\TQB\GHD2\INTAKE\INTAKE

Forms: Advance Navigator Options Help

SANDY SMITH (35, F), N1.001.IN.01.001

Please tell me everything you had to eat and drink all day yesterday, Tuesday, from midnight to midnight. Include everything you had at home and away, even snacks, coffee, soft drinks, alcoholic beverages and water. I'll ask you for specific details and amounts of the foods in a few minutes. At this time, just tell me what you had.

[ENTER THE NAME OF EACH FOOD ON A SEPARATE LINE. USE COMMENT, TIME, AND OCCASION FIELDS ONLY IF RESPONDENT PROVIDES DETAILS.]

REC	FoodName	RECComent	RECTime	RECO	RECOccasion	OS
[1]	Loose					
[2]						
[3]						
[4]						
[5]						
[6]						
[7]						

Example 4: National Cancer Institute diet history questionnaire

Diet History Questionnaire II (DHQ II) is a food frequency questionnaire developed by the US National Cancer Institute. It consists of 134 food items and includes both portion size and dietary supplement questions. It takes about one hour to complete and was designed to be easy to use. Web based versions are likely to take less time to complete. Full information about the DHQ (forms, web versions, data analysis) is available at <http://riskfactor.cancer.gov/dhq2/about/>.

The original DHQ was validated against a number of other food frequency questionnaires, with response rates varying from 70-85%, not statistically different from shorter FFQs. Diet*Calc PC software can be used to analyze DHQ data for nutrient intakes. A Dietary Assessment Resource Manual is available in Spanish, published in 2006 by the Instituto de Nutrición de Centro América y Panamá (INCAP).

Using Indirect or Secondary Data Sources

Indirect or secondary data on food consumption are typically collected by departments of finance or agriculture or by market research companies for purposes other than nutrient intakes or nutrition policy. While these data sources cannot provide precise estimates of individual food intake, they have long been used to track information on national supply and household availability of sometimes thousands of food items, summarized into broad categories of food commodities such as cereals, grains, fruits and vegetables to correspond to dietary guidelines. Estimating the intake of high salt/sodium products from secondary sources like these requires additional set of information to be included in the household budget surveys and careful

setting of assumptions. For an example of sodium intake estimated from a household budget survey, refer to the experience in Brazil (5).

Descriptions and considerations

Household income and expenditure (budget) surveys

Designed to provide information on the consumption patterns and economic conditions of private households in a specific time period, typically income, savings as well as consumption.

- With regards to food, detailed information on household level purchases of food items (including beverages) is recorded (brand information is generally not available).
- Average market prices are used to estimate the total amount acquired of each item (if not originally available).
- Inexpensive in that secondary use of budget data (to determine food consumption) is usually free.
- Additional information (such as brand information or total acquired amount of each item) can be included in the study.
- Can rely on periodicity.
- Is generally representative of the national population and the large sample size permits deductions for different strata (urban and rural, socio-economic levels, regions, provinces).
- Comparable methodologies applied across countries.
- Household consumption is divided by the number of inhabitants.
 - Individual intake is refined by estimating the proportional intake of energy by each member of the household using the adult male as the reference (i.e. the adult equivalent concept where for example an adult woman is 0.7 adult equivalents or in other words consumes 70% of the energy of an adult male) (16).
- May be able to add modules to gather data on individual food consumption with specificity for high salt/sodium products e.g. table salt, bouillon cubes, high salt condiments (see also Section 5).
- Foods purchased and consumed away from home are not usually captured, therefore may be less useful if these foods are a main source of salt/sodium.
- Non-monetary food acquisitions (such as gifts, home produced foods) are considered.

- Family member participation in household food consumption may be unclear.
- Household food stocks at beginning and end of survey are not assessed (assumed to be equal).
- Does not account for losses or product uses for other than food preparation.
- Individual food and salt intake may differ from the estimated proportion of energy intake for each household member.
- Can expect to overestimate salt intake.

Household food disappearance data

Estimates total salt used by the household and approximates the per capita (or per adult equivalent) intake.

- Typically, direct measurement is limited to a subsample of a population that is participating in a national or regional survey on food consumption or is resident in a sentinel site that is being studied.
- Relatively easy to implement to capture the total salt used from additions at the table and in cooking, bouillon cubes or dried soup mixes, fish and soy sauces and other highly salt products by counting and/or weighing the disappearance of these salt sources during a specific period of time (e.g. 7 days), and dividing that amount by the number of days and by the number of inhabitants in the household. (See also Section 5.)
 - Individual intake is refined by estimating the proportional intake of energy by each member of the household using the adult male as the reference (i.e. the adult equivalent concept where for example an adult woman is 0.7 adult equivalents because she requires 70% of the energy of an adult male).
- Requires before-and-after measurement of all salt sources in the household.
- Does not account for losses or product uses for other than food preparation.
- Individual food and salt intake may differ from the estimated proportion of energy intake for each household member.
- Usually overestimates salt intake (11).

National food disappearance data; food production data; food import/export data

Designed to capture production and consumption of agricultural commodities as well as import/export data.

- Do not reflect salt/sodium consumption.
- If salt is considered a broad general commodity,

usually cannot separate salt used for food from that used for other purposes e.g. animal feed, water softening, road salt for de-icing, etc.

Food sales data; scan data

Retail sales information usually collected by private sector market researchers.

- With regards to food, designed to collect detailed information on the sales of main retailers so brand information is likely to be available.
- Information about food consumption away from home (e.g. restaurants and cafeterias) is not included.
- Market or retailer specific i.e. there is no information on who is buying the products or how much individuals are consuming.
- Expensive to purchase.
- Can provide estimates of salt in cooking ingredients or processed foods when linked to nutrient information in appropriate food composition databases.
 - Geographic data and economic stratifications of households are likely available in markets where the surveys are conducted.
 - Can rely on periodicity.
 - Represents a large share of the sales.
 - Comparable methodologies applied across countries.
 - Utility increased when combined with data on food consumption patterns.

Attaching Questions or Modules to Existing Chronic Noncommunicable Disease Risk Factor Surveys

As an alternative to stand-alone food consumption surveys, it is worth considering whether questions or modules specific to salt intake can be attached to existing survey instruments that collect data on risk factors for chronic non-communicable diseases. Examples of such surveys are the WHO STEPwise approach available at <http://www.who.int/chp/steps/en/> and PanAm STEPS at <http://www.paho.org/english/ad/dpc/nc/panam-steps.htm>.

A set of standard questions is suggested in Population Level Sodium Determination in 24-hour Urine Samples, available at http://new.paho.org/hq/index.php?option=com_content&task=view&id=3072&Itemid=2376.

Section 3: Sodium Content of Foods

Overview

Food composition tables or databases provide detailed profiles of the nutritional composition of foods, usually for a particular country. They can contain information on a variety of components, among them: energy, macronutrients, minerals (such as sodium) and vitamins, and sometimes, individual fatty acids, amino acids and/or vitamin fractions (6). They are essential to determining the sodium content of foods consumed as identified in food consumption surveys.

Countries can use a variety of methods to construct national food composition tables, using direct or indirect data. Chemical analysis of major national foods and the calculation of values using yield and nutrient retention factors provide direct or primary data. Countries can also develop national food composition database by starting with data from another country with a similar food supply and then modifying and adding data to reflect national food culture. Some adjustments to data from other countries can be required to e.g. account for country-specific regulations on food fortification levels, the use of permissible sodium containing food additives and local food products. New or additional data can come from secondary sources, typically from industry, on food labels or in scientific literature.

- Given that data accuracy varies across sources, it is important to identify for each source in the food composition database, the country of origin of the data, whether it is primary or secondary data, if primary – the sample collection date, analytical methods used, number of samples and detailed food name identification, and if secondary – detailed food name identification and the original data sources. This will greatly assist the process of updating the database when new values become available.

Methods that Provide Direct or Primary Data

Descriptions and considerations

National/regional food composition databases

Reflects food products consumed in a country or in a region whose countries have a common food culture.

- Usually provide averages or generic composite values for certain food categories rather than brand-specific data.
- Require ongoing commitment to be kept up-to-date as food supplies are dynamic in that the

nutrient composition of food products changes e.g. the elimination of trans fats, reductions in sodium, discretionary addition of nutrients, new processes e.g. moisture enhancement of meats, and depending on the sales volumes of certain brands of a changed product e.g. white bread, the average sodium content for that particular food product may shift.

- Need to account for local traditional foods, indigenous peoples' foods and ethnic foods, foods provided by restaurants and street vendors.

Modifying/updating existing database from another country

- May save resources in that a large proportion of food products may already be accounted for in existing databases (with primary and secondary data).
- Need to supplement with some analysis of national foods to account for differences in e.g. national regulations on fortification levels and the addition of local traditional foods, indigenous peoples' foods and ethnic foods, foods provided by restaurants and street vendors.
- Can supplement with data from national secondary sources or scientific literature that reflects the national food culture.
- Need to recognize geographic variations – using values from other country databases can introduce unexpected errors.
- Within brand differences – same brand products available globally may have different formulations (e.g. added sodium) at sub-regional or country (local) levels, or have different inherent sodium content of base ingredients in various national markets, therefore it is important to note the country of origin for the product (where produced or processed).
- Requires ongoing commitment to be kept up-to-date.

Food Composition Tables and Databases from Selected Countries and Regions

Below are examples of national food composition databases and where they can be found. Other examples can be found through Langual at http://www.langual.org/langual_linkcategory.asp?CategoryID=4&Category=Food+Composition and http://www.fao.org/infoods/directory_en.stm.

ARGENTINA ARGENFOODS (Argentina Tabla de Composición de Alimentos)	http://www.unlu.edu.ar/~argenfoods/Tablas/Tabla.htm
BRAZIL TBCAUSP (Tabela Brasileira de Composição de Alimentos)	http://www.fcf.usp.br/tabela/
CANADA Canadian Nutrient File	http://webprod.hc-sc.gc.ca/cnf-fce/index-eng.jsp
CHILE Tabla de Composición Química de Alimentos Chilenos	http://mazingersisib.uchile.cl/repositorio/lb/ciencias_quimicas_y_farmaceuticas/schmidth03/index.html
CENTRAL AMERICA INCAP – Tabla de Composición de Alimentos de Centroamérica	http://www.tabladealimentos.net/tca/TablaAlimentos/inicio.html
COSTA RICA Tablas de Composición de Alimentos de Costa Rica (Macronutrientes y fibra dietética, Alimentos fortificados, Ácidos grasos)	http://devserver.paho.org/virtualcampus/costarica/drupal/
EUROPE EuroFIR (European Food Information Resource)	http://www.eurofir.org/eurofir_knowledge/european_databases
LATIN AMERICA LATINFOODS (Tabla de Composición de Alimentos de América Latina)	http://www.into.cl/latinfoods/
NEW ZEALAND FSANZ (Food Standards Australia New Zealand) – Australian Food Composition Tables (NUTTAB 2006)	http://www.foodstandards.gov.au/consumerinformation/nuttab2006/
UNITED KINGDOM UK Food Standards Agency - UK Nutrient Databank, Composition of Foods Integrated Dataset (CoF IDS) and The McCance and Widdowson's – The Composition of Foods (CoF) book series (6th Summary Edition, 2002)	http://www.food.gov.uk/science/dietarysurveys/dietsurveys/#h_1
UNITED STATES USDA (United States Department of Agriculture) – National Nutrient Database for Standard Reference	http://www.nal.usda.gov/fnic/foodcomp/search/

Using Indirect or Secondary Data Sources

Food composition databases can be constructed or modified using secondary sources of data, a process known as data compilation. In this case, a number of factors need to be addressed, key among them discussed in the table below.

A thorough discussion of quality in the compilation of nutrient data for a food composition database is provided in Chapter 10: Quality considerations in the

compilation of a food composition database. In Greenfield H and Southgate DAT. (2003). Food Composition Data: Production, Management and Use (pp.171-186). Rome: FAO, available at <http://books.google.co.cr/books?hl=es&lr=&id=KQRzKDr9bgcC&oi=fnd&pg=PR7&dq=Food+composition+data.+Production+management+and+use,+FAO&ots=nXTxTEoJ0c&sig=nOEjo0RB-sZ2nFrFcBqDx16FJ0g#v=onepage&q=Food%20composition%20data.%20Production%20management%20and%20use%2C%20FAO&f=false> Specific to LATINFOODS, Electronic Conferences Reports are available http://www.inta.cl/latinfoods/conferencia_electronica.html.

One on “Compilación de datos para bases de datos y tabla de composición química de alimentos” is available at <http://www.rlc.fao.org/foro/latfoods/infofinal.pdf>.

Another on “Evaluación de la calidad de los datos para bases de datos y tablas de composición química de alimentos” is available at http://www.inta.cl/latinfoods/inf_final_conferencia_esp.pdf.

Descriptions and considerations

Industry-provided data

Food label review for branded products

- Need to ascertain the source.
- Determine if the data are from laboratory analysis and if so the number of samples analyzed, sample preparation procedure and analytic method used, or whether values were taken/calculated from another data source e.g. calculation software, other databases, etc.
- Determine age/date of the data (may need to be tagged for collection or confirmed by the food company).
- Data may be limited e.g. only major brands, only certain cuts of meat.
- Label information is brand specific, need to account for all major brands.
- Can link label information on salt/sodium content to market share weighting (using the universal product code) to identify relative contribution of products to sodium intake.
- Should have some quantitative data to confirm the accuracy of label values in the country or the industry provided data (requires capacity to sample and analyze available data to confirm accuracy).
- Where certain products are typically imported, it is useful to know the origin(s) and whether imports from there are constant or not, as similar products can have variable formulations depending on the producer/country of origin.

- Where nutrition labelling is voluntary, nutrient data may not be consistently available.
- Where nutrition labelling is mandatory but does not require declaration of salt/sodium content, will need to find other sources of complementary data.
- Need to recognize issues with labels – rounding of values, application of compliance tolerances, conservative labelling practices, label information may not keep pace with reformulations.
- Popular products for which nutrition labelling is not required may need chemical analysis.

Restaurant chain nutrition information

- Usually web based.
- Generally less consistently available.
- Variability in information due to customization; usually only data for standard menu items are available.

Nutrition science literature for analyzed values

- Judgement is required to evaluate the relevance of the data to one's country, the sampling plan applied, the analytical and technical expertise of the laboratory and methods used before deciding on whether or not to use the published data.

Section 4: Salt Added at the Table and during Cooking

If food consumption data indicate that processed and packaged foods are the main contributors of salt in the diet, typical in higher income countries, determining qualitatively the frequency with which salt is added at the table and in cooking in households – the discretionary use of salt – provides sufficient information on which to base consumer awareness campaigns to curb the personal use/addition of salt.

Qualitative Methods

Descriptions and considerations

Supplementary questions in national, regional or sentinel site food consumption surveys or KAB surveys.

- For salt added at the table, 24-hour dietary recalls, FFQ surveys or KAB studies are supplemented with questions on “added salt” such as “what type of salt is usually added” and “how often”, with typical responses about frequency being “rarely”, “occasionally” or “very often”.
- To probe cooking practices in households, a sample question is “how often is ordinary or seasoned salt added in cooking or preparing

foods in the household” and should include “don’t know if salt is added in cooking” in case others in the household are cooking.

- Also select or probe if the respondent is the cook in the household and if yes, ask about the number of people for whom he/she prepares food.

Direct Quantitative Methods

In countries, regions or sub-populations undergoing nutrition transition – where households are known to rely on home-prepared meals at the same time that consumption of commercially prepared foods is increasing – it is important to quantify the intake of discretionary salt to differentiate it from other sources.

A number of quantitative methods are available to measure the amount of salt added at the table and in cooking (7). Typically, direct measurement is limited to a subsample of a population that is participating in a national or regional survey on food consumption or is resident in a sentinel area being studied. Other than the lithium tagging method, most of the methods described below require further research and validation and the resulting data represent rough estimates of discretionary salt use at the population level.

Descriptions and considerations

Lithium tagging of household salt

- A metabolic method.
- Normal household salt in sample households is for a specific period of time replaced by salt tagged with lithium. For each individual in the household, daily urinary excretion of lithium combined with that of sodium measures both discretionary and total salt intake (8,9,10).
- The gold standard providing the most accurate data on discretionary salt consumed.
- Can be cumbersome to the extent that is not feasible.
- Requires 24-hour urine collection from usually all members of the household (refer to the Protocol for Population Level Sodium Determination in 24-hour Urine Samples).
- Data have limited geographic and socio-economic strata comparability method requires review and validation for population level application.

Weighed household salt (disappearance method)

- Household salt is replaced by a defined quantity of salt that is measured after a specific timeframe to determine how much was used.
- Can significantly exaggerate consumption because of the large amounts of salt discarded in cooking water (11).

Using simulated meals

- Participants add salt to food models or pictures of meals which is then collected and measured.
- Simple to measure.
- Limited to what is assumed to be representative of typical meals.
- Limited to research studies or targeted individuals.

Observing cooking practices

- The amounts of salt added during cooking are observed.
- Semi-quantitative in that the observer estimates the amount of salt added by the cook.
- Difficult to generalize to large populations as household recipes can be highly variable.

Indirect “Subtraction” Method

Estimating absolute and proportional intake of discretionary salt using urinary salt excretion and primary or secondary food consumption data.

- Estimates intake of discretionary salt as:
- Discretionary salt intake = 24-hour salt excretion – salt intake attributed to commercially-available foods.
- Requires the estimation of sodium (salt) urinary excretion for a 24-hour period.
- Accuracy is highly dependent on the precision and certainty of measurements of the consumption of commercially prepared foods.
- Must account for the commercially prepared foods consumed outside the home.
- Despite the effort to account for non-food uses and losses, the values may still overestimate discretionary salt intake.

Secondary Sources

Both qualitative and quantitative determinations of the discretionary use of salt may be derived from or supplemented by scientific literature, with the cautions noted below.

Nutrition science literature

- Judgment is required to assess whether local/regional patterns of adding salt at the table and in cooking are similar to what is published in the literature.
- Assumptions on which to base accepting/adjusting the data may be necessary.

Section 5: Culturally or Regionally-specific High-sodium Foods

It is very important to account for what can be considered a sub-set of the general category of processed and

packaged foods that in some countries is subject to discretionary use in the household – the high-sodium prepared foods that are specific to the national food culture or to a region or sub-population within a country. Even though used in relatively small amounts at any one time, these products can contribute very high levels of salt to the diet because they are so frequently consumed. Examples are some pickled foods, salted fish, condiments, sauces (soy sauce, fish sauce, tomato sauce, specialty local sauces), marinades, curry pastes and soup mixes. Usually there are only a few such products in a given country. The sodium level per product is usually determined from manufacturer-provided data if laboratory analysis data are not available (12,13).

Descriptions and considerations

Supplementary questions in national, regional or sentinel site food consumption surveys, KAB surveys, household budget surveys, household food disappearance studies.

Specific local products known to be popular can be:

- queried in the 24-hour recall;
- listed in FFQs;
- added to household budget surveys;
- household food disappearance studies or
- included with questions about salt added at the table and in cooking;
- the survey instrument is customized to reflect national, regional or even more localized food cultures.

Nutrition science literature

Unless the food product featured in the literature is known to be the same or very similar in terms of salt/sodium content and patterns of use, and whose differences can be accounted for through a set of reasonable assumptions, published data on a very specific product from another country or region are not likely to accurately reflect intake in another country or region.

Section 6: Recommendations

Knowing the main sources of salt in the diet is essential to “making the case” for reducing dietary salt in that the information points out the directions for action. With key high-salt food products identified, frequency of their consumption measured and discretionary salt use understood, national public health agencies can frame the issue with baseline information, begin raising consumer awareness and at the same time engage with the food industry to encourage reformulations to lower salt/sodium content across the food products of concern. In cases where limits on the salt content of food products are regulated, monitoring the main sources of salt in the

diet and their salt levels informs the national authorities with the mandate to enforce regulations.

How the main sources of salt in the diet are determined depends on what would be most efficient, cost-effective and feasible in the context of a country's existing national nutrition policies, the capacities and resources that support them and what is already known about food and nutrient consumption. So as not to assume what countries may have in hand, the recommendations in this Section are presented in a sequence that begins with a fundamental question and progresses as information is accumulated. An overarching recommendation applying to any aspect of the sequence is for countries to use as much as possible existing processes and instruments that are known to be reliable and valid, and in the case of surveys, for them to have the required periodicity and representativeness, to which questions or modules specific to salt intake can be added at marginal cost. While the recommendations here are based on the methods reviewed in the previous sections, the preferred approach uses as much as possible validated methods, some with training materials and/or software. Nevertheless, it may be necessary to use approaches that require further validation, and where this is the case, investigators are encouraged to design their studies such that they can inform the refinement of the methods and to plan for dissemination of findings.

Determining Baseline Data on the Major Contributors to Salt Intake

Step 1: What is the overall use of discretionary salt?

In low and middle income countries and where there is evidence of nutrition transition, it is important to assess at the outset the overall discretionary use of salt in the household – salt added at the table and in cooking and the use of specific high sodium products – as this may be the most significant source of sodium in the diet. And since the quantities of salt and high sodium products added cannot be measured through 24-hour recalls, a combination of direct and indirect methods is needed.

- KAB surveys can be designed to give preliminary qualitative information on discretionary salt.
- Local food/nutrition experts can provide qualitative insights based on their experience in the area. These preliminary findings can be then further refined through e.g. focus groups.
- Household food disappearance studies can isolate the use of table salt and specific high sodium food products at the table and in cooking.
- If data on 24-hour sodium excretion are available as well as estimates of salt intake from commercially prepared foods, subtract one from the other

for an estimate of discretionary use. (*Note that if salt from commercially prepared foods is negligible, salt intake estimated by food disappearance together with changes in urinary sodium excretion may be sufficient to set intake baselines and subsequent monitoring*).

- To differentiate at the population level the personal use of table salt (added at the table and in cooking) from all other sources, the lithium-tagged salt method can be considered with results potentially contributing to further validation of this method.

Step 2: What are the food consumption patterns?

- Investigate whether established survey instruments with reliable periodicity and appropriate representativeness e.g. household income and expenditure surveys, food disappearance surveys or risk factor surveys, can be vehicles for questions or modules on food consumption that specifically investigate salt intake.
- If there is a possibility of administering 24-hour recalls or FFQS, conduct small sentinel site surveys e.g. one urban and one rural, to collect initial data on sources of dietary salt to confirm protocols and determine the expertise and resources (financial and personnel) needed to scale up to a systematic national survey.
- Conduct national surveys using 24-hour recalls or FFQs applying/adapting as much as possible the tools that are validated and supported.

Step 3: What is the proportional importance of discretionary sources versus commercially prepared foods?

- Combine data on discretionary sources with salt intake from commercially prepared foods to get the proportion of each.
- If data on 24-hour sodium excretion are available as well as estimates of salt intake from commercially prepared foods, subtract one from the other and calculate the relative proportion of discretionary sources.

Establishing and Maintaining Food Composition Databases and Tables

Regardless of the methods chosen to determine baseline data on the main sources of dietary salt, using national or international food composition databases is essential to ascertain salt/sodium intakes from food. National investments are therefore required to establish food composition tables from primary or secondary sources, and then to verify and maintain the quality

of data given their importance to nearly all activities in nutrition and food quality and safety (14).

Key questions to ask are:

- Is there a national food composition database with nutrient data including sodium that are reliable, up-to-date and available for local foods and processed and packaged foods.
- Is there a food composition database in another country that can be adapted to national food products.
- Is there an international or regional food composition database to which a country can contribute for the benefit of all countries using the database.
- If adjustments need to be made to a national or international database, what are the resources and expertise at hand, and in what timeframe is information needed.

The outcome being sought is determining the amount of salt consumed and identifying the main contributors of salt in the national diet in a manner that allows for setting targets, monitoring change in these sources and capturing new sources on an ongoing basis.

Deciding on Targets and Interventions

Based on knowledge about the main contributors to salt intake in a country, targets can be set for either the food groups that contribute the most to salt intake, for the full food chain, and/or for reductions in discretionary use, both table/cooking salt and high sodium food products. Targets can be set as averages, as maximums (upper limits) or sales weighted means per food category (see the examples below).

If targets are to apply to processed and pre-prepared food products, they should be set jointly with the food industry. A recommended first step is engaging with the sector of the food industry that accounts for high volume products that contribute a significant proportion to salt intake. Several countries have begun with the bread

sector, working with associations of small independent bakers and in some cases also supermarket chains that have in-store bakeries. Other countries have started with the producers of bread products and processed meats. Still others have approached several food producers to address salt levels in a full range of categories of packaged processed foods that are the main sources of dietary salt. Depending on resources and capacities available, subsequent or concurrent interventions can deal with food retailers, restaurants and street food vendors. Examples of targets and descriptions of how specific countries have negotiated with the food industry are shown below:

- For the United Kingdom, see <http://www.food.gov.uk/healthiereating/salt/saltreduction>
- For the [US] National Sodium Reduction Initiative, packaged food targets are at <http://www.nyc.gov/html/doh/downloads/pdf/cardio/cardio-salt-nsri-packaged.pdf> and restaurant food targets are at <http://www.nyc.gov/html/doh/downloads/pdf/cardio/cardio-salt-nsri-restaurant.pdf>.
- For the Canadian sodium reduction targets, see <http://www.hc-sc.gc.ca/fn-an/nutrition/sodium/sodium-reduction-targets-cibles-eng.php>.

Monitoring and Evaluation

Monitoring food consumption and the main sources of salt in the diet also requires collecting primary and/or secondary data as is the case with setting the baseline. It can be limited to the same food groups that determined the baseline to optimize resources, or just the targeted foods, or can include reassessing food consumption and updating food composition tables for new sources and levels of salt.

The ultimate impact analysis or evaluation of dietary salt reduction policies and interventions can include food product analysis, food consumption patterns, health status (mean blood pressure), and sodium intake measured through urine samples.

References

1. **Khohkar S, Pandar A, Cade J.** Eating habits: Food preparation and serving practices in ethnic populations: critical review and assessment of dietary survey requirements. Report to the Food Standards Agency, London, UK. FSA Project: N08015.
2. **Ferr-Luzzi A.** Keynote Paper: Individual food intake survey methods. Accessed September 18, 2010 at <http://www.fao.org/docrep/005/y4249e/y4249e0a.htm>.
3. **FAO.** Uses of Food Consumption and Anthropometric Surveys in the Caribbean – How to transform data into decision-making tools. Accessed September 18, 2010 at <ftp://ftp.fao.org/docrep/fao/008/y5825e/y5825e00.pdf>.
4. **Garriguet D.** Sodium consumption at all ages. Available at <http://www.statcan.gc.ca/pub/82-003-x/2006004/article/sodium/9608-eng.pdf>. Accessed October 1, 2010.
5. **Sarno F et al.** Estimated sodium intake by the Brazilian population, 2002-2003. *Rev. Saúde Pública.* 2009; 43:219-25. Available at http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0034-89102009000200002&lng=en&nrm=iso. Accessed September 18, 2010.
6. **Merchant AT, Dehghan M.** Food composition database development for between country comparisons. *Nutr J.* 2006; 5: 2. Available at <http://www.nutritionj.com/content/pdf/1475-2891-5-2.pdf>. Accessed September 18, 2010.
7. **Melse-Boonstra A, Rexwinkel H, Bulux J, Solomons NW, West CE.** Comparison of three methods for estimating daily individual discretionary salt intake: 24 hour recall, duplicate-portion method, and urinary lithium-labelled household salt excretion. *Eur J Clin Nutr.* 1999; 53:281-7.
8. **Leclercq C, Avalor V, Ranaldi L, Toti E, Ferro-Luzzi A.** Simplifying the lithium-marker technique used to assess the dietary intake of discretionary sodium in population studies. *Clin Sci (Lond).* 1990; 79:227-31.
9. **Sánchez-Castillo CP, James WP.** Accurate assessment of the quantitative significance of different sources of salt in the diet. *Arch Latinoam Nutr.* 1994 Sep;44(3):145-50.
10. **Venezia A, Barba G, Russo O, Capasso C, De Luca V, Farinero E, Cappuccio FP, Galletti F, Rossi G, Strazzullo P.** Dietary sodium intake in a sample of adult male population in southern Italy: results of the Olivetti Heart Study. *Eur J Clin Nutr.* 2010; 64:518-24.
11. **Sánchez-Castillo CP, James WP.** Defining cooking salt intake for patient counseling and policy making. *Arch Latinoam Nutr.* 1995; 45:259-64.
12. **Sharma S, Harris R, Cao X, Hennis AJ, Leske MC, Wu SY, et al.** Nutritional composition of the commonly consumed composite dishes for the Barbados National Cancer Study. *Int J Food Sci and Nutr.* 2007; 58:461-74.
13. **Sharma S, Yacavone MM, Cao X, Samuda PM, Cade J, Cruickshank K.** Nutritional composition of commonly consumed composite dishes for Afro-Caribbeans (mainly Jamaicans) in the United Kingdom. *Int J Food Sci and Nutr.* 2009; 60 Supp 7:140 – 50.
14. **Bangkok Declaration** at the 8th International Food Data Conference, Bangkok, Thailand, 3rd October 2009. Accessed September 18, 2010 at www.fao.org/infoods/Bangkok%20Declaration.final.doc.
15. **Charlton KE, Steyn K, Levitt NS, Jonathan D, Zulu JV, Nel JH.** Development and validation of a short questionnaire to assess sodium intake. *Public H Nutr.* 2007;11:83-94.
16. **Levy RB, Claro RM, Bandoni DH, Mondini L.** Per capita versus adult-equivalent estimates of calorie availability in household budget surveys. *Cad. Saúde Pública, Rio de Janeiro.* 2010;26.

Appendix 1: Examples from the Region

Barbados

The Barbados National Commission for Chronic Non-Communicable Disease is spearheading a country-wide nutrition intervention and improvement programme aiming to reduce chronic diseases (in particular, heart disease, stroke and diabetes). The prevalence of hypertension, the principal risk factor for cardiovascular disease, exceeds 50% in the 40 to 80 age group, making the Barbadian high salt diet a target for intervention.

In 2010, the Barbados Ministry of Health sent out a request for proposals to develop a Salt Intake Study with two components to:

- Determine a baseline of actual sodium intake (through 24-hour urine sampling).
- Identify the main sources of dietary sodium in the population.

By 2005, a quantitative food frequency questionnaire (QFFQ) had been developed and was in use in the Barbados National Cancer Study through a collaboration between Stony Brook University in New York, the University of the West Indies and the Cancer Research Center of Hawaii. The list of foods in this QFFQ instrument had been derived from food intake data collected in 2000 through the Barbados Food Consumption and Anthropometric Survey (BFCAS). For the BFCAS, 1600 respondents 18 years of age and over completed a single 24-hour dietary recall, and 50 (of 63) randomly selected residents gave additional updated information on foods that they consumed. All foods, beverages and supplements reported by at least two BFCAS participants were included in the draft QFFQ. This instrument was piloted in a random sample of individuals (n=50) selected from the nationwide electoral register. Participants completed a 24 hour dietary recall followed by an interviewer-administered QFFQ. Foods with minimal contribution to nutrient intake were excluded, and the final QFFQ contained 148 food and drink items.

There was evidence of some secular changes in the Barbadian diet between 2005 and 2010, and the current QFFQ, will therefore have to be amended for use in the Salt Intake Study. To this end ongoing work to date has entailed:

- Phase 1 – a pre-pilot in April 2010:
 - to update and amend the existing QFFQ by the inclusion of additional food items and to create a more comprehensive QFFQ to allow the evaluation of salt consumption;
- Phase 2 – a pilot of the amended QFFQ in May 2010;
- Phase 3 – administration of the updated QFFQ to a nationally representative sample of the population.

It is anticipated that these new dietary data will provide information on nutrient intake and cooking practices to be targeted for a nutritional intervention aimed at reducing salt intake.

The first step of Phase 1 began with the selection of a random sample of participants from the electoral register who were mailed a letter of invitation to participate in the study, followed by telephone or personal contact by study staff to make appointments for recall interviews. Nutritionists were trained to conduct a series of three non-consecutive 24 hour dietary recalls with each participant. These recalls also included questions probing for additions to the diet such as salt, sugar, seasonings and condiments. Information on portion size (using food models, standard units or household utensils, later converted to standardized weights), specific types of foods and drink including brand names, cooking methods (boiled, fried, baked, “lime and salt”) and food sources were also recorded.

Twenty-five men and 27 women (aged 25 to 90 years; mean ages were 46 and 55, respectively) participated in Step 1 (83% response rate). Data represented 148 days of dietary recall. Preliminary results showed that the mean daily sodium intake among men aged 27-50 years and 51-73 years was 4,313 mg and 2,653 mg, respectively, while women 25-50 years and 51-90 years consumed 2,897 mg and 2,126 mg of sodium per day, respectively. The leading dietary sources of sodium were rice and peas (5.4%), fried flying fish (4.4%), and macaroni pie (3.2%). Bread, fish, rice, poultry, and sweets were major food group sources of sodium, contributing a combined 50% to total sodium consumption. Mean sodium intake levels exceeded Adequate Intake recommendations across all gender-age groups. In Step 2, a focus group with Barbadian nurses, a dietitian and researchers facilitated the draft of the QFFQ with the addition of new food items reported more than once by participants in Step 1, with particular attention paid to items rich in sodium e.g. condiments and preserved foods. Of note, questions about the practice of adding salt at the table and others asking about participants’ willingness to reduce dietary sodium were included. The new draft QFFQ was initially piloted in May 2010 with 25 respondents who added other foods they frequently consumed. Final updates were completed by July 2010.

The Salt Intake Study has now been approved by the funding agency and the start of data collection is anticipated in mid-2011.

Prepared in collaboration with Professor Sangita Sharma, Endowed Chair in Aboriginal Health, Professor in Aboriginal and Global Health, University of Alberta, Canada and Anselm Hennis, Professor of Medicine and Epidemiology, the University of the West Indies, Cave Hill Campus, Barbados.

Brazil

The National Household Budget Survey (HBS – Pesquisa de Orçamentos Familiares) administered between July 2002 and June 2003 by the Brazilian Institute of Geography and Statistics (IBGE, Instituto Brasileiro de Geografia e Estatística) provided the data from which the main sources of sodium in the Brazilian diet were derived (4). The HBS is designed to obtain information on the earnings and expenditures of families in order to adjust the structure of the national consumer price index.

The 2002/03 version of the HBS surveyed a probabilistic sample representative of all Brazilian households as well as each of the country's five macro-regions (according to the urban/rural status of the household) or 26 states. Sampling was based on a complex (two-stage) strategy involving 443 previously defined socio-geographic strata of census tracts (aiming to assure high representativeness with a minimum sample size). Subsequently, census tracts were selected from within each stratum and households were selected from within each tract. Finally, in order to make data collection uniform across the four trimesters of the year, the interviews carried out within each stratum were spread out across the 12 months of the survey. A total of 48,470 households participated in the survey.

With regards to diet, the HBS collects detailed information on all food and drink purchases made by households during a period of seven consecutive days, including the name and specification of each item, its price and where it was acquired (i.e. grocery store, bakery). The 48,470 households contributed a total of 969,989 records. Data were collected electronically with the help of portable computers and specially developed software.

Since the seven-day reference period is not sufficient to characterize the food-purchase pattern of each household, the survey unit was adapted to be groups of households (rather than individual households) corresponding to the 443 HBS strata.

The household units in each stratum are homogeneous in terms of territorial domain and family socioeconomic status and they were surveyed uniformly throughout the trimesters of the year.

Initial data analysis excluded any non-edible fraction from each of the purchases using correction factors recommended by IBGE. The edible fraction of each food item was then converted into energy (kcal) and sodium (grams) using version 1 of the Brazilian Food Composition Table (TACO – Tabela Brasileira de Composição dos Alimentos), or where an item was not in TACO, using the United States official food composition table version 15. (An extension to the statistical package Stata – AQUINUT – was developed for this purpose) (5). In the specific case of foods preserved in salt such as salted/dried

beef and salted fish, an additional adjustment allowed the sodium concentration of these items to refer to the desalted product. Thus for each survey unit (household stratum), the daily per capita availability of energy and sodium was calculated.

Given that meals eaten outside the home and the fraction of food purchases not consumed are unknown, the household sodium availability to actual intake was estimated by considering the sodium content per 2,000 kcal of total energy consumed – the Brazilian recommendation for daily per capita energy intake. Food and beverages were then classified into four groups: 1) salt and salt-based condiments; 2) processed foods with added salt; 3) in natural foods, or processed foods without added salt; and 4) ready-made meals. The proportion that each food group contributes to total household sodium availability was described for the country as a whole and according to quintiles of per capita income distribution.

The results indicate that the amount of sodium available for consumption in Brazilian households was 4.7 g/person/day, more than two times the maximum internationally recommended intake level of 2 g/person/day. Excessive consumption was identified in all regions, in both rural and urban settings, and across all income strata. Although most of the available sodium in all income strata originates from discretionary salt (added at the table and in cooking) and salt-based condiments, the proportion derived from processed foods has a strong and growing importance as household purchasing power increases in Brazil. The complete set of results was published in the journal “Revista de Saúde Pública” (6).

Canada

In 2004, Statistics Canada carried out the Canadian Community Health Survey (CCHS) 2.2 – Nutrition using a 24-hour food recall instrument to which a statistically representative sample (based on census data) of over 35,000 people responded from all provinces (the survey excluded members of the armed forces, residents of the three northern territories, people living on native reserves, in institutions and some remote locations). A subset of respondents (>10,000) completed a repeat recall three to 10 days later (to assess day-to-day variation, which was used for determining usual intakes). The response rate was 76.5% and 72.85% respectively. The survey provided information on the then current intakes of dietary sodium by the Canadian population, which is the baseline against which the effectiveness of strategies to reduce salt consumption would be measured.

Respondents were asked to list all foods and beverages consumed during the 24 hours (from midnight to

midnight) before the day of their interview. Interviewers used a Canadianized version of the USDA Automated Multi-pass Method with five steps to help respondents remember what they had had to eat and drink:

- a quick list (respondents reported all items in whatever order they wished);
- questions about specific food categories and frequently forgotten foods;
- questions about the time and type of meal;
- questions seeking more detailed, precise descriptions of foods/beverages and quantities consumed;
- a final review.

An additional salt questionnaire was administered. People who replied “None” when asked “What type of salt do you usually add to your food at the table?” were classified as “never adding salt at the table”. Otherwise, respondents were asked how often they added salt to their food, with the answer options of rarely, occasionally, or very often. Respondents were also asked about certain health conditions that were expected to last or had already lasted six months or more that had been diagnosed by a health professional. Those who answered “yes” to “Do you have high blood pressure?” were defined as having hypertension (7,8).

The nutrient values, including sodium levels for foods that respondents consumed were obtained from Health Canada’s Canadian Nutrient File (version 2001b) (9). The average daily intakes of sodium, as well as percentages of the population who exceed the Upper Level (UL) were reported for each DRI age and sex group for age 1 up (7). Proportions of the population who reported adding salt to food at the table or in cooking as well as the 10 main sources of sodium were provided. Additional analyses were done by province and by frequency of high blood pressure (adults aged 31 and older).

The CCHS survey were used for a further detailed analysis of food sources of sodium for four age groups derived by regrouping the Dietary Reference Intakes (DRI) age–sex groups according to expected similarities in sodium intake patterns (8). The child DRI age–sex groups were divided by age into two groups: young people aged 1 to 8 years and those aged 9 to 18 years. The adult DRI age–sex groups were divided by sex into two groups: males aged 19 years and older, and females aged 19 years and older.

The food groups used in this study were based on an established classification system developed in the Food Directorate of Health Canada and used in previous nutrition surveys. Some of the food groups were altered for sodium modelling purposes e.g. by separating them into canned, pickled, smoked, or dried foods, usually higher in sodium than the rest of the group.

The analyses focused on estimating the sources of so-

dium by food group, with a number of parameters being reported for each food group:

- food intake per person, in grams per day, calculated as the total grams of the foods reported within the group divided by the total population;
- the sodium amount in foods, in milligrams per 100 g, calculated from the total sodium consumed, divided by the total grams of the foods consumed within the group;
- the percent of total sodium intake, calculated from the total sodium consumed within the group divided by the total sodium consumed by the population;
- the daily sodium intake per person in milligrams per day, calculated as the total sodium intake within the group divided by the total population;
- the percent of the population consuming the food, calculated from the number of consumers having reported eating at least one of the foods within the group divided by the total population; and
- the daily sodium per consumer in milligrams per day, calculated as the total sodium intake within the group divided by the number of consumers having reported eating at least one of the foods within the group.

In addition, the average daily energy provided by each food group was calculated by multiplying the energy density in kcal per gram by the per capita food intake for each food group. The amount of sodium provided by the quantity of food supplying 100 kcal of energy was determined by dividing the daily per capita sodium intake from each food group by the per capita energy intake for that group.

The analyses conducted in this study (8) were based on first-day recalls only. Usually a repeat recall is used to adjust for intra-individual variability to estimate the usual intake distribution. However, for the food groups considered in the sodium modeling, the frequency of their consumption did not meet the requirements for estimating a usual intake distribution from the CCHS 2.2 sample.

- 1 **Hennis A, Wu SY, Nemesure B, Leske MC; Barbados Eye Studies Group.** Hypertension prevalence, control and survivorship in an Afro-Caribbean population. *J Hypertens.* 2002 Dec;20(12):2363-9.
- 2 **Sharma S, Cao X, Harris R, Hennis AJ, Leske MC, Wu SY;** Barbados National Cancer Study Group Dietary intake and development of a quantitative food-frequency questionnaire for the Barbados National Cancer Study. *Public Health Nutr.* 2007 May;10(5):464-70.
- 3 **Sharma S, Harris R, Cao X, Hennis AJ, Leske MC, Wu SY:** Nutritional composition of the commonly consumed composite dishes for the Barbados National Cancer Study. *Int J Food Sci Nutr* 58:461-474, 2007.

- 4 **Instituto Brasileiro de Geografia e Estatística.** Coordenação de Índices de Preços. Pesquisa de Orçamentos Familiares 2002-2003: análise da disponibilidade domiciliar e estado nutricional no Brasil. IBGE; 2004.
- 5 **Núcleo de Pesquisas Epidemiológicas em Nutrição e Saúde.** Conversor de aquisições de alimento em energia e nutrientes (AQUINUT): versão 1.0 [internet] [cited 2007 Oct 12]. Available from: <http://www.fsp.usp.br/nupens>
- 6 **Sarno F, Claro RM, Levy RB, Bandoni DH, Ferreira SRG, Monteiro CA.** Estimated sodium intake by the Brazilian population, 2002-2003. *Rev. Saúde Pública.* 2009; 43:219-25. Available at http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0034-89102009000200002&lng=en&nrm=iso. Accessed September 18, 2010.
- 7 **Garriguet D.** Sodium consumption at all ages. *Health Reports.* 2007;18:47-52.
- 8 **Fischer PWF, Vigneault M, Huang R, Arvaniti K, Roach P.** Sodium food sources in the Canadian diet. *Appl. Physiol. Nutr. Metab.* 2009;34:884-92.
- 9 **Health Canada.** 2001. The Canadian Nutrient File. Available at www.healthcanada.gc.ca/cnf. Accessed on January 11, 2011.

Acknowledgements

Mary L'Abbé

Department of Nutritional Sciences, University of Toronto, Canada

Adriana Blanco Metzler

Health and Nutrition Unit, Costa Rican Institute for Teaching and Research on Nutrition and Health

Norm Campbell

Libin Cardiovascular Institute of Alberta, University of Calgary, Canada

Francesco Cappuccio

WHO Collaborating Centre for Nutrition, Warwick University, England

Omar Dary

Academy of Educational Development, Washington DC, USA.

Daniel Ferrante

Health Promotion and Chronic Disease Control, Ministry of Health, Argentina

Christina Howitt

Chronic Disease Research Centre, Barbados

Charmaine Kuran

Nutrition Evaluation Division, Food Directorate, Health Canada

Branka Legetic

PAHO Secretariat

Barbara Legowski

PAHO Secretariat

Rafael Moreira Claro

School of Public Health, University of São Paulo, Brazil

Renata Levy

School of Medicine, University of São Paulo, Brazil

Ana Beatriz Vasconcellos

Food and Nutrition Coordination, Ministry of Health, Brazil

Eduardo Nilson

Food and Nutrition Coordination, Ministry of Health, Brazil

Rosangela Pereira

University of Rio de Janeiro, Brazil

Rosely Sichieri

University of Rio de Janeiro, Brazil

**Protocol for the Survey on Knowledge,
Attitudes and Behavior toward Dietary
Salt and Health**

Introduction

In September 2009 PAHO established an expert group to explore the current epidemiological situation regarding cardiovascular disease and its link to excessive salt consumption in the Americas and review existing policies, interventions and programs aimed at reducing dietary salt and issue evidence-based recommendations for salt reduction in the region.

Clear food labeling with regard to salt and/or sodium together with consumer awareness can help reduce the salt intake and is therefore recommended.

Justification

The 37th session of the CODEX Committee on Food Labeling did not reach consensus whether salt or sodium should be used on food labels and established an Electronic Working Group. Section C of the comments of the EWG members (Annex to CCFL minutes of the 38th session, Canada, May 2010) show that in Latin America no research has been undertaken with respect to public/consumer understanding of the terms salt and sodium, little campaigning to raise consumer awareness on the consumption of high amounts of salt has been undertaken, while all countries show to have guidelines and recommendations to this respect.

Main Objective

For this reason the PAHO expert group proposed to conduct a survey to be realized in 5 countries by national consumer associations, a project to be led by Consumers International and financed by PAHO. The overall aim of this quantitative survey is to establish a base line on consumer knowledge, behavior and labeling preference with respect to salt and sodium. This can be used to provide input to the next CODEX committee meeting on salt vs. sodium labeling of food and to an in-depth qualitative follow-up survey on the same topic.

Hypothesis

Using 20 (multiple options and open) questions, we intend to establish:

- the attitudes of consumers with regard to salt and health;
- the knowledge of consumers with respect to his/her use of salt or sodium in food;
- the average knowledge of consumers of the relation between high salt consumption and possible health problems;
- whether the consumer suffers/has suffered from

a salt-related disease (and has knowledge of this condition);

- whether the consumer does something to control his/her salt intake;
- the preference of consumers for labeling food (salt, sodium, both, warning labels, percentages, absolute quantity);
- relations of the above-mentioned with age, sex, educational level, attitude.

A direct relationship is expected between educational level or the fact that consumers suffer from a salt-related disease and knowledge of the possible health problems high salt consumption can provoke. A direct and inverse relationship between age of healthy consumers and knowledge of the possible health problems high salt consumption can provoke is expected.

Limited knowledge of the possible health problems high sodium consumption can provoke is expected in the majority of the respondents. Labeling preference for salt is expected for consumers with lower levels of education.

This survey should confirm these expectations or prove them incorrect.

Methodology and Data Analysis

The consumer organizations will conduct the survey with the following remarks:

- establish first the **effectiveness** of the questions, i.e. make sure they are understandable for the people in your country and avoid technical language where possible (e.g., do not use scientific names of diseases), once this has been done print sufficient forms;
- find out if in your country a consent of some ethical review board is needed to conduct the survey; if yes, please make sure you obtain one (and change questions if necessary);
- ask a minimum **400** different respondents in your country in two batches (so in the second batch, emphasis can be given to missing demographic groups);
- provide an even distribution of men and women (e.g., if two people conduct the interviews, one can only interview women while the other only interviews men);
- provide an even distribution in age as much as possible; please ask **only** adults (> 18 years);
- if possible, divide the sample in two representative groups; one group consisting of people living in cities, the other of people living in rural areas (please make sure these groups can be identified).

- avoid that the interviewee can read the questionnaire as much as possible (e.g., use a notebook computer or a notepad);
- avoid leading the interviewee, i.e., stick to the questions;
- answer the questions in the order provided, i.e. do **not** turn back to correct earlier answers;
- when using paper forms:
 - identify the questionnaires by using initials of the interviewer and a number (e.g., AB-001);
 - circle the options clearly on the questionnaire; in case of corrections mark the first answer given by X and circle the correct one; except where indicated **only one option** is allowed;
 - write open answers clearly legible on the questionnaire.
- when using a notebook :
 - indicate correct answer with color or erase other options; except where indicated **only one option** is allowed;
 - type open answers on the form;
 - do not forget to save the form; the filename can be the identification of the interviewer, e.g., AB-001.
- Q1–Q3 + Q6 +Q8 + Q10 + Q12–Q14 + Q16–Q19: read the question aloud and the options if needed;
- Q4: read question aloud but **not** the options; indicate option(s) mentioned (more than one option allowed) and write down in case “Other” is chosen;
- Q5: read question aloud and the options; each should get an answer;
- Q7: read question aloud and write down the answer;
- Q9: ask only if Q8 = yes; read question aloud and write down the answer including unit (e.g. grams), if Q8 ≠ yes, skip this question;
- Q11: ask only if Q10 = yes; read question aloud and write down the answer; if Q10 ≠ yes, skip this question;
- Q15: ask only if Q14 = salt/sodium/salt and sodium; read question aloud and write down the answer, if Q14 ≠ yes, skip this question;
- Q20: read question aloud and the options; only one is allowed;
- Q21: in case the interviewee has any comments, please note those here;
- after the interview, make sure all questions have been answered, thank the interviewee and hand out a PAHO Patient/Consumer factsheet on salt to the interviewee;
- process each form using the attached excel spreadsheet*;
 - in case there are not sufficient worksheets (40)

within the spreadsheet, please copy the entire spreadsheet (always keep one clean copy of the spreadsheet);

- please name each sheet using the initials and numbers to identify the interview;
- please mark options with “x”;
- send electronic file(s) to Consumers International Santiago.

All answers will be processed as percentages of sub-groups or the total of respondents, except:

- age (will be used to establish age groups);
- Q7 (what is done to control salt intake; groups of similar answers will be established);
- Q9 (recommend daily salt intake; answers will be shown as a range);
- Q11 (difference between salt and sodium; groups of similar answers will be established);
- Q15 (reason for labeling preference; groups of similar answers will be established);
- Q21 (groups of similar comments will be established).

References

<http://www.surveysystem.com/sdesign.htm>

*To save time, when using a notebook computer, the interviewer can type the answers directly into the excel sheets instead of the word forms.

Appendix 1:

Questionnaire on Knowledge, Attitudes, and Behavior toward Dietary Salt and Health

N°:	Date:	Interviewer:
-----	-------	--------------

Age	
-----	--

Sex	female	male
-----	--------	------

Children under 16 and living at home

yes	no	no answer
-----	----	-----------

Highest education level

none	primary	secondary	higher	no answer
------	---------	-----------	--------	-----------

Attitude statements

I try to eat a healthy diet	agree	disagree	don't know	no answer
Eating a diet high in salt can cause serious health problems	agree	disagree	don't know	no answer
I try to minimize the amount of fat I eat	agree	disagree	don't know	no answer
My health is generally good	agree	disagree	don't know	no answer
There is too much pressure to eat healthily these days	agree	disagree	don't know	no answer
I try to minimize the amount of salt I consume	agree	disagree	don't know	no answer
I know in general how much salt food contains	agree	disagree	don't know	no answer
There is sufficient nutritional information on labels of food and drinks	agree	disagree	don't know	no answer

1. How often do you add salt to food at the table?

never	rarely	sometimes	often	always	don't know
-------	--------	-----------	-------	--------	------------

2. In the food you eat at home salt is added in cooking

never	rarely	sometimes	often	always	don't know
-------	--------	-----------	-------	--------	------------

3. How much salt do you think you consume?

too much	right amount	too little	don't know	no answer
----------	--------------	------------	------------	-----------

4. What sort of health problem do you think can be caused by a high salt diet?

(do *not* read aloud + tick all that apply)

a) high blood pressure	b) osteoporosis	c) stomach cancer	d) kidney stones
e) heart attack/heart failure	f) stroke	g) asthma	h) other (specify):
i) none	j) don't know	k) no answer	

5. Do you suffer from or have you suffered from: ?

High blood pressure	yes	no	don't know	no answer
Heart attack	yes	no	don't know	no answer
Stroke	yes	no	don't know	no answer
Kidney stones	yes	no	don't know	no answer
Asthma	yes	no	don't know	no answer
Osteoporosis	yes	no	don't know	no answer
Stomach cancer	yes	no	don't know	no answer

6. Limiting the amount of salt/sodium I eat is important to me.

not at all	somewhat	very	don't know	no answer
------------	----------	------	------------	-----------

7. What, if anything, do you do to control your salt or sodium intake?

8. Do you know if there is a recommended amount for salt/sodium to be eaten per person per day?

yes	no	don't know	no answer
-----	----	------------	-----------

9. If yes, please indicate the amount?

10. Do you know the difference between salt and sodium?

yes	no	don't know	no answer
-----	----	------------	-----------

11. If yes, please indicate the difference?

12. Do you pay attention to indications on packages like "no added salt", "low in salt", "light", "free of trans fat"?

always	often	sometimes	rarely	never	don't know	no answer
--------	-------	-----------	--------	-------	------------	-----------

13. How often do you read nutrition labels on food packages?

always	often	sometimes	rarely	never	don't know	no answer
--------	-------	-----------	--------	-------	------------	-----------

14. What do you prefer on nutrition labels on food packages?

salt	sodium	salt and sodium	don't know	no answer
------	--------	-----------------	------------	-----------

15. Why?

16. Would you like labelling of food indicating high/medium/low levels of salt or sodium?

yes	no	don't know	no answer
-----	----	------------	-----------

17. Would you like to see a clear warning label on the package if foods are high in salt?

yes	no	don't know	no answer
-----	----	------------	-----------

18. Would you like labelling of food indicating the amount of salt or sodium in grams or milligrams?

yes	no	don't know	no answer
-----	----	------------	-----------

19. Would you like labelling of food indicating salt or sodium as a percentage of the amount recommended to be eaten per person per day?

yes	no	don't know	no answer
-----	----	------------	-----------

20. Do you prefer labelling of food indicating salt or sodium per portion or the total amount per package?

per portion	per 100 gr	total per package	don't know	no answer
-------------	------------	-------------------	------------	-----------

21. Comments:



**Optimizing Salt
and Iodine Intake**

**White Paper on Improving
Public Health by Optimizing
Salt and Iodine Intakes**

Background

The International Council for the Control of Iodine Deficiency Disorders (ICCIDD) estimates that two billion people worldwide live in areas at risk of iodine deficiency. It states that among the methods to prevent iodine deficiency disorder (IDD), using salt as a vehicle to deliver supplementary iodine to the diet is most simple, practical and effective (1). Together with WHO and UNICEF, it recommended in 1996 the iodine level at the point of production of 20-40 ppm, assuming an average per capita salt intake of 5-10 g per day (2).

In 2006, the report from a WHO Forum and Technical Meeting responding to the WHO Global Strategy on Diet, Physical Activity and Health recommended that average population level salt intake from all food sources be < 5g/day per person, in order to reduce the sodium intake to less than 2 g/d per person (3). Salt added to food is a major factor increasing the blood pressure in normotensive and hypertensive people, whether adults or children [4,5,6]. Increased blood pressure is the leading risk factor for death worldwide and the second leading risk for disability by causing heart disease, stroke and kidney failure (7,8).

In 2007 a WHO Expert Consultation on "Salt as a Vehicle for Fortification", taking into account both the status of IDD and the evidence of harmful effects of high salt diets, emphasized the need for and benefits of coordination between salt iodization and dietary salt reduction programs (9). It concluded that

"Policies for salt iodization and reduction of salt to < 5 g/day are compatible, cost effective and of great public health benefit. At the country level, close collaboration between salt iodization and salt reduction programs as a coalition is urgently required so that their aims are congruent." [9, p. 15]

Most recently, with dietary salt reduction mobilizing in the Americas, the PAHO/WHO Regional Expert Group guiding the initiative indicated in its 2009 Policy Statement (10) that

Salt intake can be reduced without compromising micronutrient fortification efforts. National governments are to review national salt fortification policies and recommendations to be in concordance with the internationally recommended target of < 5 g salt /day per person by 2020 or national targets if lower.

And the Iodine Network, in a resolution dated February 20, 2009 (11) stated

BE IT FURTHER RESOLVED THAT the World Health Organization is urged to share with the Network Board any data, when and if it becomes available, about the achievement of salt reduction in various countries and its projected trends of salt consumption to inform Board discussion and national salt iodization programs to enable ongoing efforts to calibrate iodine fortification levels to ensure appropriate population iodine intakes.

Currently underway on the advice of the WHO Nutrition Guidance Expert Advisory Group (NUGAG) is a review of the evidence on how varying levels of population salt intake can impact the effectiveness of salt iodization programs, intending that salt reduction and salt iodization strategies work efficiently and effectively together. The results will feed into future revised WHO/UNICEF/ICCIDD salt iodization program guidelines, to become the starting point for newly coordinated efforts between iodine nutrition and dietary salt reduction. Needed as interventions are readied for implementation are current accurate baselines of actual iodine and salt intakes (assessed as sodium in the urine) and of the main sources of dietary salt and iodine (i.e. table salt and processed and pre-prepared food sources), information that is lacking in most countries. This would be followed by rigorous simultaneous measurement of urinary iodine and sodium, and food consumption patterns to monitor the progress of and feed critical information back to both programs.

Synchronization of salt iodization and dietary salt reduction programs brings together several stakeholders at international and national levels: the agencies working to optimize iodine supplementation and those focused on cardiovascular disease prevention; national governments; and various sectors of the salt and food industries. When the knowledge and experience of the stakeholders involved in the two programs are coordinated, with the stakeholders playing their respective roles within a framework for action directed at a common goal of mutual benefit, cost savings can be realized for healthcare systems.

A group of technical experts and stakeholders in both IDD and dietary salt reduction programs, convened by PAHO, has developed this White Paper to facilitate a broad collaboration between the programs, having agreed to a common goal and a Framework for Collaborative Action. It is directed to stakeholders for the two programs active within countries as well as those operating at the international level.

The Common Goal

The achievement of optimal intakes of sodium and iodine.

A Framework for Collaborative Action

- 1. Common and coordinated messaging** at global, regional and national levels to
 - a. policy and decision makers;
 - b. the salt and food industries;
 - c. stakeholders among the health professions;
 - d. the public and consumers.
 - 2. Common advocacy platforms** to
 - a. integrate the development/adjustment of iodine fortification of salt and dietary salt reduction policies and programs
 - i. at national or sub-national levels taking into account localized food supplies and dietary practices;
 - ii. noting the need for collaboration among the diverse sectors and groups within the sectors involved in both initiatives;
 - b. implement effective and regular quality assurance and monitoring programs for iodization of salt and iodine nutrition;
 - c. synchronize national efforts to monitor iodine as dietary salt is reduced and where iodine deficiency is a concern, advance policies for the voluntary or mandatory use of iodized salt or iodine-containing premixes in commercially produced food at levels appropriate to population iodine needs;
 - d. harmonize wherever possible cross country approvals processes to admit new food products with low salt content and an adequate amount of iodine;
 - e. emphasize the importance of optimal iodine intake;
 - f. emphasize the importance to health and the cost savings to health care systems of reduced dietary salt intake.
 - 3. Concurrent surveillance** of salt and iodine intake where feasible to inform salt iodization and dietary salt reduction programs including but not limited to
 - a. methods that optimally assess and monitor salt and iodine intake including potassium where a public health concern;
 - b. comprehensive food surveys to distinguish the main sources of salt and iodine in the diet (including potassium where a public health concern) with questions to assess:
 - i. the discretionary use of iodized salt at the table and in household cooking;
 - ii. salt intake through the consumption of processed foods, restaurant meals and street food and
 - iii. the proportion of iodine in the diet contributed by each source;
 - c. methods that account for vulnerable and diverse populations;
 - d. establishing, promoting and supporting laboratory proficiency for iodine and sodium analysis, (and potassium where a public health concern);
 - e. knowledge, attitudes and behavior surveys on salt consumption;
 - f. monitoring the plans and patterns of the processed food industry with regards to
 - i. provision of sodium (and/or salt) data on food labels;
 - ii. the feasibility of including iodine on labels;
 - iii. the markets where new salt-containing products are being or will be supplied/imported, especially in countries undergoing nutrition transition, to anticipate changes in salt intake levels and whether the products use iodized salt or are otherwise a source of iodine.
- 4. Coordinated evaluations** of national salt iodization and dietary salt reduction programs
 - a. applying a common set of principles including transparency and minimized conflicts of interest;
 - b. committed to information sharing;
 - c. independent of food and salt industries;
 - d. demonstrating the link between action and disease outcomes.
- 5. Strategic joint research** to fill knowledge gaps relevant to both salt iodization and reduction of dietary salt that emphasizes but is not limited to pilot and case studies in countries of differing economic and cultural make-up on
 - a. how to most effectively optimize sodium and iodine intake;
 - b. the most effective and feasible collaborative surveillance methods to determine sodium and iodine intake and the sources of salt and iodine in the diet.
- 6. Shared forums with relevant sectors of the food industry** to deal with iodine and sodium additives and promote
 - a. the voluntary or mandatory use of iodized salt or iodine-containing premixes in commercially produced food;
 - b. improved capacity and technology of the salt industry to ensure consistent and high standards of iodization of salt of small and medium sized salt producers.

- c. calibration of iodization levels in salt based on the different salt intake levels with
 - i. the food processing industry and the restaurant and catering sectors consistently using iodized salt;
 - ii. the food processing industry and the restaurant and catering sectors reducing the salt content of processed and pre-prepared foods;
 - iii. joint technical assistance and knowledge sharing between sectors;
 - iv. compatible positions on issues held in common e.g. international trade agreements and regulatory or voluntary frameworks governing the sectors such that both salt iodization and dietary.

salt reduction programs can achieve their goals in the established timeframes.

- 7 Coordinated mapping of existing and needed resources** and mobilization of resources towards but not limited to
 - a. Concurrent surveillance, policy development, advocacy and consumer education.

Disclaimer: The findings and conclusions in this meeting summary are those of the authors and do not necessarily represent the official position of their organizations or of the Pan American Health Organization.

References

- 1 **ICCIDD. The Global Picture.** Accessed April 5, 2011 at <http://www.iccidd.org/pages/protecting-children/fortifying-salt.php>.
- 2 **World Health Organization.** 1996. Recommended iodine levels in salt and guidelines for monitoring their adequacy and effectiveness. Based on a joint WHO/UNICEF/ICCIDD consultation, World Health Organization, 8-9 July 1996, Geneva, Switzerland. Accessed April 4, 2011 at http://whqlibdoc.who.int/hq/1996/WHO_NUT_96.13.pdf.
- 3 **World Health Organization.** 2007. Reducing salt intake in populations: report of a WHO forum and technical meeting, 5-7 October 2006, Paris, France. Accessed April 4, 2011 at http://www.who.int/dietphysicalactivity/Salt_Report_VC_april07.pdf.
- 4 **He FJ, Marrero NM, MacGregor GA.** Salt and blood pressure in children and adolescents. *J Hum Hypertens.* 2008;22:4-11.
- 5 **He FJ, MacGregor GA.** A comprehensive review on salt and health and current experience of worldwide salt reduction programmes. *J Hum Hypertens.* 2009;23:363-84.
- 6 **He FJ, MacGregor GA.** Importance of salt in determining blood pressure in children: Meta-analysis of controlled trials. *Hypertension.* 2006;48:861-69.
- 7 **World Health Organization.** The World Health Report 2002: Reducing Risks, Promoting Healthy Life. 8 Hsu C, McCulloch CE, Darbinian J, Go AS, Iribarren C. Elevated blood pressure and risk of end-stage renal disease in subjects without baseline kidney disease. *Arch Intern Med* 2005;165:923-28.
- 9 **World Health Organization.** 2008. Salt as a Vehicle for Fortification: report of a WHO expert consultation, 21-22 March 2007, Luxembourg. p 15. Accessed April 4, 2011 at http://whqlibdoc.who.int/publications/2008/9789241596787_eng.pdf.
- 10 **Pan American Health Organization.** Policy Statement on Dietary Salt Reduction. Accessed April 4, 2011 at http://new.paho.org/hq/index.php?option=com_content&task=view&id=2022&Itemid=1766.
- 11 **Personal communication, Lucie Bohac,** 2011, from unpublished Minutes of the Meeting of the Board of the Iodine Network on February 20, 2009 in Jaipur, India.

**Improving Public Health in the
Americas by Optimizing Sodium
and Iodine Intakes**

Key Messages

In Latin American countries, iodized salt has the potential to secure the sustained elimination of iodine deficiency at the same time that dietary salt consumption is reduced to prevent cardiovascular diseases. Synchronization of programs for salt iodization and dietary salt reduction to achieve a common goal – the optimal intake of sodium and iodine in the Americas – will be cost effective and of great public health benefit.

Lowering dietary salt intake to < 5 g/day is compatible with salt iodization programs based on flexible iodization levels for salt.

Needed at the outset of program coordination are up-to-date and accurate baselines of actual salt (sodium) and iodine intakes and the main sources of dietary salt and iodine (i.e. household iodized salt added at the table and in cooking or other sources), information that is currently lacking in most countries in the Americas.

Background

An estimated two billion people worldwide live in areas of iodine deficiency (1). The International Council for the Control of Iodine Deficiency Disorders (ICCID) states that among the methods to prevent iodine deficiency disorder (IDD), using salt as a vehicle to deliver supplementary iodine to the diet is most simple, practical and effective (2). WHO and UNICEF together with ICCIDD recommended in 1996 iodine concentrations in salt that assume an average per capita salt intake of 10g per day (3).

In 2006, the report from a WHO Forum and Technical Meeting responding to the WHO Global Strategy on Diet, Physical Activity and Health recommended that average population level salt intake from all food sources be < 5g per person per day (4). Salt added to food is a major factor increasing the blood pressure in normotensive and hypertensive people, whether adults or children (5,6,7). Increased blood pressure is the leading risk factor for death worldwide and the second leading risk for disability by causing heart disease, stroke and kidney failure (8,9).

In 2007 a WHO Expert Consultation on “Salt as a Vehicle for Fortification”, taking into account both the status of IDD and the evidence of harmful effects of high salt diets, emphasized the need for and benefits of collaboration between salt iodization and dietary salt reduction programs (10). Among its recommendations.

Policies for salt iodization and reduction of salt to < 5 g/day are compatible, cost effective and of great public health benefit. At the country level, close collaboration between salt iodization and salt reduction programs as a coalition is urgently required so that their aims are congruent.

The level of iodine fortification needs to be adjusted by national authorities responsible for the implementation and monitoring of universal salt iodization in light of their own data regarding dietary salt intake. The average national level of salt consumption must provide key guidance for the concentration of iodine in salt.

Most recently, with dietary salt reduction mobilizing in the Americas, the PAHO/WHO Regional Expert Group guiding the initiative indicated in its 2009 Policy Statement (11) that Salt intake can be reduced without compromising micronutrient fortification efforts. National governments are to review national salt fortification policies and recommendations to be in concordance with the recommended salt intake [national targets or in their absence, the internationally recommended target of < 5 g/day/person by 2020]. And the Iodine Network, in a resolution dated February 20, 2009 stated BE IT FURTHER RESOLVED THAT the World Health Organization is urged to share with the Network Board any data, when and if it becomes available, about the achievement of salt reduction in various countries and its projected trends of salt consumption to inform Board discussion and national salt iodization programs to enable ongoing efforts to calibrate iodine fortification levels to ensure appropriate population iodine intakes.

Following up on the 2007 WHO Expert Consultation and the recommendations from 2009 of the PAHO/WHO Group for dietary salt reduction and the Iodine Network, PAHO convened a small group of technical experts and some stakeholders in both IDD prevention and dietary salt reduction programs on 3-4 January 2011 in Washington DC. They agreed that they hold a goal in common – the optimal intake of sodium and iodine in the Americas. The group then drafted objectives, expected results and a background document for a subsequent meeting involving an expanded group of experts and stakeholders in both programs to facilitate a broader collaboration and to define the next steps needed to advance the synchronization of dietary salt reduction and salt iodization programs in the region.

The larger group of stakeholders met in Washington DC on 31 March and 1 April 2011. The background material became the basis for a Position Document and Framework for Collaborative Action. Their meeting objectives and outcomes were as given below.

Objectives:

1. Define the common ground for simultaneous salt iodization and salt reduction initiatives and discuss a draft position document.
2. Agree to a Framework for Collaboration/Coordination.
3. Define next steps to operationalize the Framework.

Outcomes:

1. A Position Document that reflects areas of collaboration/coordination.
2. A Framework for Collaborative Action.
3. Discussion on next steps in IDD (salt iodization) and dietary salt reduction for the Americas region.

On day one of the meeting, presentations brought the participants up to date on both topics of dietary salt reduction and salt iodization. They included: the evidence for why salt reduction should be implemented at the population level; recent advances in IDD prevention; plans for the ICCIDD for Latin America and participation in the International Resources Laboratories for Iodine Network (IRLI); review of the 2007 recommendations in “Salt as a Vehicle for Fortification”; an update on the activities of the Iodine Task Force; and the WHO (global) activities on salt intake reduction. The day concluded with a guided discussion toward defining the common ground for collaboration between salt iodization and dietary salt reduction programs to be stated in the Position Document expected as a meeting outcome.

On day two, participants were divided into three groups to prepare responses to a draft version of a *Position Document to Improve Public Health by Optimizing Sodium and Iodine Intakes* that included a Framework for Collaborative Action. They also worked out several next steps to operationalize the components of the Framework, oriented at governments, consumers/the public and the salt and food industries.

NB For the purpose of this document, the word **salt** is used to refer to **sodium** and **sodium chloride**.

Presentation Summaries

The evidence for why salt reduction should be implemented at the population level – Norm Campbell

- In 2009 WHO reported high blood pressure (systolic >115 mmHg) as the leading single risk factor for death (12). Its impact on vascular diseases is

profound, attributed to 60-70% of strokes, 50% of heart failures, 25% of heart attacks, 20% of kidney failures and is associated with dementia due to cerebrovascular damage.

- Blood pressure is well known to rise with age in modern industrialized societies. The US Framingham Heart Study reported in 2002 that men and women 55 to 65 years of age who were non-hypertensive had an estimated lifetime risk of 90% of becoming hypertensive (13). In societies where people eat unprocessed foods, are physically active and lean, they do not develop hypertension (14).
- A vast and conclusive body of evidence, ranging from animal studies, migration studies, epidemiological studies (cross sectional and cohort, examining hypertension and vascular disease), clinical trials, and meta-analyses points out the relationship between salt intake, hypertension and vascular disease. A Cochrane review in 2006 concluded that even a modest reduction in salt intake for a duration of four or more weeks has a significant effect on blood pressure in individuals who are normotensive as well as in those with elevated blood pressure. The meta-analysis is consistent with other findings where the lower the salt intake, the lower the blood pressure (15).
- If dietary salt were reduced to recommended levels, hypertension prevalence is estimated to decrease by 30%. Even small reductions in blood pressure can lower mortality rates for stroke and coronary heart disease (16). If dietary salt were reduced by even 15% worldwide over 10 years, an estimated 8.5 millions deaths could be averted. In the US, if salt intake fell to 3g/day/person, in one year between \$10 and \$30 billion could be saved, there could be up to 260,000 fewer cardiovascular disease events and up to 90,000 fewer deaths. In low and middle-income countries, reducing dietary salt is slightly more cost effective than smoking reduction.
- A high salt diet is also a probable cause of gastric cancer, and has possible associations with osteoporosis, calcium containing renal stones and increased severity of asthma. Because salty foods cause thirst they are likely an important contributor to obesity among children and adolescents through association with increased consumption of high-calorie soft drinks (6,17).
- People are generally unaware of their own level of salt consumption. With few exceptions, average consumption is over 5.8g/day after age 5 and for many, intake is over 10g/day. In developed econo-

mies about 80% of the salt consumed is added during food processing while in less developed economies, most of the salt consumed is discretionary, added at the table and in cooking. In countries undergoing nutrition transition, there is a shift in the main sources of dietary salt, from discretionary use to the “hidden” salt in processed foods as they become increasingly available to consumers (18).

- Salt intakes around the world are much higher now than the physiologic levels that supported human evolution. The current levels are linked to major causes of death and disability worldwide. Reducing dietary salt is estimated to be one of the most effective and cost effective public health interventions (19).

Recent advances in the prevention of IDD - Ruben Grajeda

- Iodine is an essential component of the hormones produced by the thyroid gland. Thyroid hormones regulate many key biochemical reactions, especially protein synthesis and enzymatic activity. Major target organs are the developing brain, muscle, heart, pituitary, and kidney.
- Severe iodine deficiency in pregnancy can cause hypothyroidism, poor outcomes of pregnancy (spontaneous abortion and stillbirth), cretinism and in some countries is a major cause of irreversible mental retardation.
- Mild to moderate iodine deficiency in utero and in childhood results in less severe learning disability, poor physical growth and diffuse goiter.
- In adults, mild to moderate iodine deficiency appears to be associated with higher rates of more aggressive sub-types of thyroid cancer and increases the risk for non-toxic and toxic nodular goiter and associated hyperthyroidism.
- Worldwide two billion people have insufficient iodine intake from their usual diet. In the Americas 98 million people are estimated to have insufficient iodine intake. In 2006, the Americas had the lowest global prevalence of insufficient iodine intake at 10%. Between 2003 and 2006, the rate in the Americas was stable while in other WHO regions it fell (20,21).
- Strategies/options to prevent IDD include fortification of salt, bread, water and milk, and iodine supplementation. Salt iodization is the most cost effective intervention to prevent IDD. Recommended intakes are standardized by age and population group (22).

Age or population group	IOM		Age or population group	WHO RNI
	EAR	AI or RDA		
Infants 0-12 months		110-130	Children 0-5 yr	90
Children 1-18 yr	65	90	Children 6-12 yr	120
Children 9-13 yr	73	120		
Adults ≥ 14 yr	95	150	Adults ≥ 12 yr	150
Pregnancy	160	220	Pregnancy	250
Lactation	200	290	Lactation	250

IOM, Ref. 8; WHO, Ref. 54

UI (ug/liter)	Iodine intake	Iodine nutrition
School-aged children		
<20	Insufficient	Severe iodine deficiency
20-49	Insufficient	Moderate iodine deficiency
50-99	Insufficient	Mild iodine deficiency
100-199	Adequate	Optimum
200-299	More than adequate	Risk of iodine-induced hyperthyroidism in susceptible groups
>300	Excessive	Risk of adverse health (iodine-induced hyperthyroidism, autoimmune thyroid disease)
Pregnant women		
<150	Insufficient	
150-249	Adequate	
250-499	More than adequate	
≥500 ^a	Excessive	
Lactating women ^b		
<100	Insufficient	
≥ 100	Adequate	
Children less than 2 yr of age		
< 100	Insufficient	
≥ 100	Adequate	

There is no information about iodine nutrition for pregnant and lactating women in the WHO assessment table, and the upper limits of the median UI for lactating women and children less than 2 yr of age were not specified.

a The term excessive means in excess of the amount needed to prevent and control iodine deficiency.

b In lactating women, the numbers for median UI are lower than the iodine requirements because of the iodine excreted in breast milk.

- There are several methods to assess iodine status: thyroid size (a population's history of iodine nutrition and its present status); urinary iodine concentration (indicator of recent iodine intake); level of thyroid stimulating hormone (useful biomarker in neonates, pregnant and lactating women); thyroglobulin (useful marker in children and adolescents); and thyroid hormone concentration (30,31).
- Excess iodine intake induces hypothyroidism, autoimmune thyroiditis with chronic hypothyroidism and hyperthyroidism after long-standing iodine deficiency (producing goiters).
- In the US, even though iodine status is adequate, pregnant women are at high risk of iodine deficiency just as are women in other developed economies (23).
- Salt iodization programs should be monitored and evaluated to assure adequate iodine intake.

Plans of the ICCIDD for Latin American Countries and Participation of the IRLI Network – Eduardo Pretell

- In Latin American countries, with improvements in the production and monitoring of iodized salt, salt as a vehicle for iodine has the potential to secure the sustained elimination of IDD at the same time that dietary salt consumption is reduced to prevent cardiovascular diseases.
- Most countries in Central and South America and the Caribbean along with Mexico have official IDD control programs with the exception of Belize, Argentina and Guyana (uncertain in Haiti). Where programs exist, coverage with the household use of iodized table salt is 100% except for Guatemala (60%), the Dominican Republic (78%) and Bolivia (94%).
- Rudimentary technology to produce iodized salt remains an obstacle to the provision of good quality iodized salt in many countries in Latin America. In about half of the countries, small/medium scale producers provide all the iodized salt. In Guatemala for example, only 43.5% of the iodized salt at production level contained 30-60 ppm of iodine and at retail, 64% of iodized salt had >15 ppm. In Mexico, where small/medium scale producers account for 17% of iodized salt, only 54% of their products were found to have >15 ppm whereas 96% of the iodized salt supplied by large producers (83% of total iodized salt production) had >15 ppm.
- Across Latin American countries, the presence of


salt with ≥ 15 ppm iodine at the retail/household level can vary from less than 5% in Haiti (2006) to 100% in Uruguay (2006).

- The resource laboratory for Latin America that is part of the International Resource Laboratories (IRLI) Network is in Lima Peru – the Laboratory of Endocrinology at the High Altitude Research Institute at Cayetano Heredia Peruvian University. The IRLI laboratories' main activities are quality assurance of laboratories for urinary iodine, technical support to improve their capacity for diagnosis and monitoring, and the processing of urine samples (surveys, research) as requested by countries.
- In 2006 the Peru resource laboratory was at that time able to provide technical support to laboratories in Argentina, Bolivia, Brazil, Chile, Costa Rica, Cuba, El Salvador, Mexico, Panama, Paraguay and Uruguay. Six professionals from Argentina, Brazil, Cuba and Mexico were trained and supported to set up laboratories.
- There were 25 registered laboratories in Latin American countries in 2006 but only 15 were operative, listed below. An inter-laboratories assay run from 2003 to 2006 showed 12 out of 21 participating laboratories to have made significant progress.

Laboratories processing iodine in urine

- Centro Nacional de Investigaciones Nutricionales. Salta, Argentina.
- Instituto Nacional de Laboratorios de Salud. La Paz, Bolivia.
- Centro de Investigación y Desarrollo de Tecnología de Alimentos. Santa Cruz, Bolivia.
- Instituto de Nutrición y Tecnología de los Alimentos, Univ. de Chile. Santiago, Chile.
- Instituto Nacional de Salud. Bogotá, Colombia.
- Inst. Costarricense Nutr y Enseñanza Nutr y Salud. MOH. San José, Costa Rica.
- Instituto de Nutrición e Higiene de los Alimentos, MOH. Havana, Cuba.
- Laboratorio de Yodurias. MOH. Quito, Ecuador.
- Laboratorio de Bioquímica. INCAP. Guatemala, Guatemala.
- Lab Central de Referencia de Estudios en Salud Pública. Panama.
- Instituto Nacional de Alimentación y Nutrición. Asunción, Paraguay.

Fig. 2: Urinary Iodine



- Laboratorio de Micronutrientes. Fac Ciencias y Filosofía, UPCH. Lima, Peru.
- Centro Nacional de Alimentación y Nutrición, MOH. Lima, Peru.
- Fac. Química. Univ. Rep. Oriental del Uruguay. Montevideo, Uruguay.
- Dep. Bioquímica, Universidad de los Andes. Mérida, Venezuela.
- Regular monitoring of iodine nutrition is not carried out in all countries. In half of the countries with data on urinary iodine levels since 2005, the median µg/l ranges from approximately 190 µg/l to 440 µg/l

(Paraguay, Honduras, Peru, Ecuador, Chile, Cuba, Panama, Bolivia, Venezuela). In countries with data before 2005, the urinary iodine levels were found to range from about 40 µg/l (Dominican Republic in 1993) to about 420 µg/l (Columbia in 2002).

- At this time, ICCIDD is mainly pursuing the improvement of the quality of iodized salt, the implementation of effective monitoring and reporting systems, and sustained communication and education. A regional meeting is planned for 1-2 August 2011 in Lima Peru with the following agenda:
 - discussion of the IDD control programs in each country;

- how to improve the capacity and responsibility of the salt industry;
- importance of optimal iodine nutrition in early life;
- implementation of WHO recommendations;
 - securing the elimination of IDD by USI strategy;
 - reduction of salt consumption (sodium) to prevent cardiovascular disease;
- current and future role of Agencies and NGOs.
- Reaching the above objectives will facilitate the implementation of programs for dietary salt reduction.
- Starting pilot studies to reduce dietary salt without risking optimal iodine intake could be considered in one or two countries where effective IDD programs are implemented. This will require the collaboration/integration/coordination of various stakeholders in both salt iodization and salt reduction programs including ICCIDD given its network of volunteer experts/focal points in almost all Latin American countries.
- There were 19 recommendations in the 2007 document “Salt as a vehicle for fortification” (9). The current initiative to realize a collaboration between salt iodization and dietary salt reduction programs provides an opportunity to analyze why the recommendations have not been implemented and at the same time it poses an important challenge – to critically examine what has been achieved with salt iodization programs and the bases on which they have been judged to be safe and successful.
- Analyzing one recommendation – “the iodine concentration in salt should be determined considering both the level of salt consumption and median urinary iodine (UI) of the population” – introduces a number of issues:
 - While UI is found to be sufficient in various regions in a country, the concentration of iodized salt (table salt) and its availability (indicators of IDD program effectiveness) by region do not correspond, in some cases, suggesting inadequate intake. Conversely, high quality iodized table salt is readily available while UI is found to be insufficient, particularly in vulnerable groups (pregnant and lactating women).

Reviewing the Key Recommendations in “Salt as a Vehicle for Fortification” – Omar Dary

- Safe, efficacious and sustainable programs for salt iodization and dietary salt reduction require the involvement of several sectors and many players within each of them in a logical sequence where a clear separation of roles is important as is transparency of what each sector and its players are contributing. This is particularly relevant with regards to the salt and food industries whose participation is essential and at the same time needs to be specifically defined.
- Currently the use of iodized salt in food processing is not consistent across countries in the Americas. A potential approach to make universal the use of iodized salt would have the following sequence. Nutritionists and researchers would first complete the scientific and epidemiological assessments of the appropriate sodium and iodine ratios needed in a country, and governments would then prepare the national policies and strategies with corresponding standards/regulations. The “premix” manufacturers (in the case of salt iodization) could then respond with production of micronutrient premixes that the food industries incorporate into their production processes. The salt and food industries also have a role, working collaboratively with governments, in social marketing and public education on what constitutes healthy intake of sodium and iodine.
- The dietary sources of salt vary. While household salt is iodized, salt used in food manufacturing may not be; the use of iodized salt in processed foods is not currently measured or monitored. Where processed foods made with un-iodized salt are the main source of salt in the diet (while household table salt intake is relatively low), there is risk of iodine deficiency. Hence salt reduction and salt iodization policies must vary from country to country based on food consumption patterns.
- The extent to which natural sources of iodine are present in the diet and the use of iodine supplements (or not) need to be considered.
- There is still much to learn about UI levels e.g. how to adjust for urine volumes. Collaboration between salt iodization and salt reduction programs has potential to accelerate a better understanding of UI as an indicator of the effectiveness of salt iodization programs.
- The Iodine Task Force (ITF) operates under the IDD Network umbrella. A Steering Group with WHO, UNICEF and ICCIDD helps to assure that the work is aligned with and feeds into the systematic review

Update on the Iodine Task Force – Lucie Bohac

on salt iodization and also feeds into a revision of field-appropriate program guidelines.

- A Management Group (A. Timmer, J. Gorstein, F. van der Haar, L. Bohac) is responsible for day-to-day management, defines structures and processes, coordinates background research, supports five technical working groups, and is responsible for communication and logistic support.
- The five working groups comprise experts in the various areas from within and outside the Network member organizations. Their responsibilities are:
 - WG 1 – establishment of salt iodization standards to achieve optimal iodine intake.
 - WG 2 – collection and interpretation of data on iodine status.
 - WG 3 – collection and interpretation of iodine content in household and food industry salt.
 - WG 4 – other iodine interventions.
 - WG 5 – goal and program indicators.
- The Task Force was conceived in April 2010 and was underway by September 2010. Each WG is led by a nominated focal point and consists of five to seven subject matter specialists. Each has set its expected outputs on a time trajectory. The groups will converge on a joint draft final report due at the end of May 2011.
- Linkages to salt intake reduction initiatives depend on the elements of national strategies, whether there are:
 - standards and norms affecting the supply of household salt;
 - legislation concerning the salt supply to food manufacturing industry;
 - legislation affecting the combination of the salt supply channels (households, food industry and livestock);
 - programs based on diverse and/or multiple food supply channels:

- spontaneous fluctuations in salt or iodine intake e.g. special events/circumstances requiring adjustment such as high iodine intake from natural sources and natural sodium intake fluctuations.

- There are several opportunities for collaboration between the salt iodization and salt reduction programs, e.g. using each other's expertise and outcomes. Going forward, it is essential to preserve the benefits of collaboration by using realistic projections of national achievements from each policy. USI strategies (standards) have to adjust according to achievements made from national salt intake reduction policies. Programs need to work collaboratively in the monitoring of progress.

Population salt reduction strategies: activities and plans at WHO Headquarters – Godfrey Xuereb

- Of the six objectives in the 2008-2013 Action Plan for the Global Strategy for the Prevention and Control of Non-communicable Diseases, directly relevant to dietary salt reduction is “reducing and preventing risk factors”. Under each objective are sets of actions for Member States, a WHO Secretariat and international partners.
- The WHO population salt reduction strategy has three platforms. Platform 1 – Create enabling environments – was organized jointly with the UK FSA in the summer of 2010. There was an information exchange forum with the private sector and NGOs on population-based salt reduction strategies followed by a technical meeting.
- Platform 2 was organized jointly with the Government of Canada (Health Canada) in the fall of 2010. It was also an information exchange forum with the private sector and NGOs followed by a technical meeting, both on strategies to monitor and evaluate population sodium consumption and sources of sodium in the diet.
- Platform 3 intends to bring together dietary salt reduction and salt fortification with iodine, to facilitate the coordination of the two strategies.
- In the meantime, a Nutrition Guidance Expert Advisory Group (NUGAG) is considering evidence that suggests lower sodium intake (1.2g/day) may confer additional health benefits (compared to the current guideline of <2g sodium or <5g salt/day), and is undertaking systematic reviews of literature to answer two priority questions:
 - What is the effect of reducing sodium intake to 1.2 g/day versus 2 g/day on blood pressure and

incidence of stroke, coronary heart disease, and cardiovascular disease in hypertensive and non-hypertensive adults?

- What is the effect of reducing sodium intake to the equivalent of 2g/day (based on caloric intake) versus > 2g/day on blood pressure in children?
- Also under consideration, at the request of CO-DEX, is potassium consumption for the general population, suggested by evidence to have health benefits, for which WHO does not have a current guideline/optimal level. Systematic reviews of literature currently underway are seeking to answer the question:
 - What is the effect of consuming > 90 mmol/day of potassium versus < 90 mmol/day on blood pressure and incidence of stroke, coronary heart disease and cardiovascular disease in hypertensive and non-hypertensive adults?
- NUGAG is also updating a Cochrane systematic review from 2002 to answer – is salt iodization safe and effective for reducing IDD? Subgroups to be examined are: by age (<59 months, 5-12 years, women of reproductive age, pregnant women); population salt consumption (<5, 5-9.9, 10-14.9, ≥15g/day); iodine consumption based on urinary iodine excretion; concentration of iodine in salt (<20, 20-40, >40ppm); and availability of iodized salt (household only, processed foods only, USI, unknown).
- NUGAG will have draft recommendations based on the findings from all reviews at a November 2011 meeting.

Meeting Outcomes

The Way Forward

- Meeting participants responded to the background document prepared by the group that met in January 2011. It became the Position Document, see Appendix 1.
- Meeting participants prepared responses to five questions to inform the way forward with governments, the salt and food industries and consumers:
 - What arguments should be used to promote a common view as well as national ownership (for dietary salt reduction and improving iodine status)?
 - What information is needed to be convincing, and how to get it and present it?
 - What messages and attitudes should be avoided because they may hinder plans and intentions?

What is still needed to overcome internal discrepancies?

- Who should make contact with the sector, when and how?
- What is the sequence of activities, where and at what level (regional, sub-regional, national)?

Governments

- The essential starting point to influence governments is evidence-based reviews developed through international collaborations of researchers mobilized and supported by WHO, PAHO, ICCIDD and UNICEF. With the lead agencies committed to collaboration and scientific reports ready, clear and simple messages, oriented for regional collaboration and avoiding directive approaches, can be distilled from the science such that if change is expected, solutions are obvious and allow for tailoring to suit national contexts to minimize political and public anxieties about re-balancing sodium and iodine intakes. Then on a country level, researchers and academics can be mobilized to prepare local evidence to build the case for a national initiative. They, together with representatives of the international agencies and local partners in various sectors in the country e.g. associations of health professionals, civil society leaders, NGOs, consumer associations, can as a coalition present the case for coordinated salt reduction and salt iodization (or alternatives for the latter) to national policy and decision makers.
- Essential for a coordinated response by the salt and food industries to achieve the optimal intakes of sodium and iodine is a sequence of actions to level the playing field for both the salt producers and food manufacturers. It begins with an enhancement of existing national regulatory frameworks that currently govern the prevention of IDD through salt iodization, that being regulation to require universal salt iodization (USI) and the universal requirement for food manufacturers to use it where not currently in effect. This would be followed by an examination of food import and export policies to ensure their congruence with the mandatory requirement for iodized salt in food products.
- As salt and iodine intakes are rebalanced, governments need to monitor both the intake levels of sodium and iodine along with the concentrations of iodine in salt and of salt in food products. Evaluations of progress need to be transparent and apply methods and indicators that are valid and reliable.
- Important to knowledge transfer is documentation of pilot projects and case studies to feature lessons

learned and demonstrate the successful recalibrations of iodine to salt ratios and harmonization of salt reduction and salt iodization programs.

Salt and food industries

- With the fundamental requirement for USI made mandatory, where the iodine to salt density ratio is based on total average salt intake of less than 5g/day/person, the opportunity arises for food manufacturers to adopt a standardized salt/sodium content target for their food products, e.g. mg Na/kcal value of the food, to deliver optimal iodine and sodium levels together. Where voluntary reformulations to reduce salt content by the food industry are protracted, and given the critical public health importance of recalibrating the iodine to salt ratio to reach the optimal intakes of both, governments can consider regulating a salt/sodium density to advance the progress of both programs.
- Essential to constructive engagement of the salt and food industries (the latter comprising food manufacturers and food service establishments) is an emphasis on positive outcomes – the reassurance that public agencies (governments, NGOs, civil society) are prepared to increase public awareness, in particular among vulnerable groups, of the benefits of combined optimal intakes of sodium and iodine and increase consumer demand for products that are in line with intake targets. Healthier foods should not have a negative impact on sales (if marketing and pricing policies are supportive) and the scientific evidence confirms that a range of iodine concentrations in salt makes no taste difference to consumers and people can adapt to lower salt content in relatively short time periods.
- Corporate leadership among the salt and food industries, especially with champions or umbrella associations, to coordinate the reduction of salt content of food products while retaining appropriate iodine fortification levels can accrue efficiencies to both industries. Governments need to be ready to congratulate the successes and where necessary provide technical and economic support to e.g. small salt producers to improve their supply of a consistent quality of iodized salt.

Consumers

- Avoiding mixed messages is crucial. Public education to raise awareness and where necessary change behavior is essential, to disseminate the information on the positive health effects of optimal intakes of both sodium and iodine, why dietary salt needs to

be reduced while iodine intake needs scrutiny especially among specific populations such as pregnant and lactating women and school age children, and where salt is added at the table and in cooking, that it should be iodized.

- Where processed foods are or are becoming the main sources of salt in the diet, consumers need to become aware that they can demand control of their salt intake to keep it optimal for their health. Most currently available processed and pre-prepared foods have excessive amounts of salt/sodium that is added before the product is sold.
- The engagement of experts in consumer behavior, to design and evaluate information campaigns, is important to ensure as much as possible that messages are clear and understood, are context specific i.e. take into account whether the main sources of salt in the diet are discretionary salt use or processed and pre-prepared foods, and that vulnerable populations are being reached effectively.
- It is important for healthcare professionals to be well informed to reinforce the public education campaigns with consistent messages and advice.

Conclusions and Next Steps

Education and sensitization within agencies

- Agencies involved in salt iodization and dietary salt reduction programs should ensure that staff is educated on the intersection of the programs and sensitized to the potential for unintended conflicting advocacy and education messages.

Engage the international stakeholders

- Once the currently involved stakeholders approve the Position Document, PAHO will approach UNICEF for its endorsement. Other stakeholders to be approached are e.g. the InterAmerican Heart Foundation and Latin American Societies for Nephrology and Hypertension.
- WHO Headquarters Platform III meeting on iodine fortification will likely be attached to the upcoming NUGAG meeting in Korea on 28 November – 2 December 2011.

Discussion and further research on key technical issues

- A number of key technical issues require further thought and discussion and possibly research, in-

volving technical sub-groups of experts, one example being the selection of target groups in the case of concurrent sodium and iodine intake surveillance.

Engage the salt and food industries

- As soon as all key stakeholders approve the Position Document, the salt and food industries can be engaged as broadly as possible, including small and medium enterprises, to elaborate their roles in implementation.

Prepare for the pilot studies

- PAHO and the main stakeholders and governments in the region will establish the criteria on which to base the selection of countries in Central and South America eg the indicators of active and functional programs for dietary salt reduction and salt iodization, where the collaboration and harmonization of the programs can be piloted.
- PAHO and the main stakeholders will approach their respective focal points in the selected countries plus local academics to assemble teams of researchers and technical experts to initiate grant proposal preparation.

Mobilize resources

- PAHO will explore the receptivity of (NIH) NHLBI and the Child Health and Endocrinology Institute as well as the Swiss Federal Institute of Technology.

Timeline

- The Position Document will be completed by mid/end April.
- All key stakeholder approval to be completed by mid May 2011.
- The full meeting report to be ready by the end of June 2011 (in time to inform the Iodine Task Force (ITF) at its next meeting).
- A technical group in mid summer convened by PAHO and ITF, place to be determined, (using the outcomes of the five ITF working groups and the PAHO 24-hour urine sampling protocol to measure sodium and iodine intake) to identify the elements of a joint iodine and sodium surveillance protocol that are ready now (guidance and tools) and what still needs to be researched, to ultimately draft a comprehensive protocol for joint surveillance for the pilot studies.
- PAHO and MI will prepare a joint intervention for the upcoming June 2011 annual meeting of the Institute of Food Technologists.
- The criteria for selecting countries for the pilot studies to be ready for the August 2011 ICCIDD Latin America workshop in Peru.
- The research framework for the pilot studies to be presented to the Dietary Salt Reduction Expert Group in September 2011.
- The pilot study project to be presented at the WHO Platform III meeting in November/December 2011.

References

- 1 **de Benoist B, McLean E, Andersson M, Rogers L.** Iodine Deficiency in 2007: Global progress since 2003. *Food Nutr Bull.* 2008;29:195-202.
- 2 **ICCIDD. The Global Picture.** Accessed April 5, 2011 at <http://www.iccidd.org/pages/protecting-children/fortifying-salt.php>.
- 3 **World Health Organization. 1996.** Recommended iodine levels in salt and guidelines for monitoring their adequacy and effectiveness. Based on a joint WHO/UNICEF/ICCIDD consultation, World Health Organization, 8-9 July 1996, Geneva, Switzerland. Accessed April 4, 2011 at http://whqlibdoc.who.int/hq/1996/WHO_NUT_96.13.pdf.
- 4 **World Health Organization. 2007.** Reducing salt intake in populations: report of a WHO forum and technical meeting, 5-7 October 2006, Paris, France. Accessed April 4, 2011 at http://www.who.int/diet-physicalactivity/Salt_Report_VC_april07.pdf.
- 5 **He FJ, Marrero NM, MacGregor GA.** Salt and blood pressure in children and adolescents. *J Hum Hypertens.* 2008;22:4-11.
- 6 **He FJ, MacGregor GA.** A comprehensive review on salt and health and current experience of worldwide salt reduction programmes. *J Hum Hypertens.* 2009;23: 363-84.
- 7 **Feng J, MacGregor GA.** Importance of salt in determining blood pressure in children: Meta-analysis of controlled trials. *Hypertension.* 2006;48:861-69.
- 8 **World Health Organization.** The World Health Report 2002: Reducing Risks, Promoting Healthy Life.

- 9 **Hsu C, McCulloch CE, Darbinian J, Go AS, Iribarren C.** Elevated blood pressure and risk of end-stage renal disease in subjects without baseline kidney disease. *Arch Intern Med.* 2005;165:923-28.
- 10 **World Health Organization. 2008.** Salt as a Vehicle for Fortification: report of a WHO expert consultation, 21-22 March 2007, Luxembourg. Accessed April 4, 2011 at http://whqlibdoc.who.int/publications/2008/9789241596787_eng.pdf.
- 11 **Pan American Health Organization.** Policy Statement on Dietary Salt Reduction. Accessed April 4, 2011 at http://new.paho.org/hq/index.php?option=com_content&task=view&id=2022&Itemid=1766.
- 12 **World Health Organization. 2009.** Global Health Risks. Mortality and burden of disease attributable to selected major risks.
- 13 **Vasan RS, Beiser A, Seshadri S, Larson MG, Kannel WB, D'Agostino RB, Levy D.** Residual lifetime risk for developing hypertension in middle-aged women and men – The Framingham Heart Study. *JAMA.* 2002;287:1003-10.
- 14 **Meneton P, Jeunemaitre X, de Wardener HE, MacGregor GA.** Links between Dietary Salt Intake, Renal Salt Handling, Blood Pressure and Cardiovascular Disease. *Physiol Rev.* 2005;85:679-715.
- 15 **He FJ, MacGregor GA.** Effect of longer-term modest salt reduction on blood pressure. *Cochrane Database of Systematic Reviews* 2004, Issue 1. Accessed April 6, 2011 at http://www.worldactiononsalt.com/evidence/docs/cochraneupdate_2006_salt_metaanalysis.pdf.
- 16 **Whelton PK, He J, Appel LJ, Cutler JA, Havas S, Kotchen TA, Roccella EJ, Stout R, Vallbona C, Winston MC, Karimbakas J;** National High Blood Pressure Education Program Coordinating Committee. Primary prevention of hypertension: clinical and public health advisory from the National High Blood Pressure Education Program. *JAMA.* 2002;288:1882-88.
- 17 **He FJ, Marrero NM, MacGregor GA.** Salt intake is related to soft drink consumption in children and adolescents: a link to obesity? *Hypertension* 2008 Mar;51(3): 629-34.
- 18 **Brown IJ, Tzoulaki I, Candeias V, Elliott P.** Salt intakes around the world: implications for public health. *Int J Epidemiol.* 2009;38:791-813.
- 19 **Asaria P, Chisholm D, Mathers C, Ezzati M, Beaglehole R.** Chronic disease prevention: health effects and financial costs of strategies to reduce salt intake and control tobacco use. *Lancet.* 2007;370:2044-53.
- 20 **de Benoist B, McLean E, Andersson M, Rogers L.** Iodine deficiency in 2007: global progress since 2003. *Food Nutr Bull.* 2008;29:195-202.
- 21 **Pretell E, Grajeda R.** Iodine nutrition in Latin America. *IDD Newsletter.* 2009;31:1-5.
- 22 **Zimmermann MB.** Iodine deficiency. *Endocr Rev.* 2009;30:376-408. Epub 2009 May 21.
- 23 **Caldwell KL, Makhmudov A, Ely E, Jones RL, Wang RY.** Iodine Status of the U.S. Population, National Health and Nutrition Examination Survey, 2005-2006 and 2007-2008. *Thyroid.* Epub 2011 Feb 16.

Appendix 1: Meeting Participants and Contributors to the Position Document

<p>Lucie M. Bohac Coordinator</p>	<p>Iodine Network 180 Elgin St, Suite 1000, Ottawa, ON, K2P 2K3 CANADA tel: +1 613 782-6812 / fax: +1 613 782-6838 www.iodinenetwork.net</p>	<p>lbohac@micronutrient.org</p>
<p>Kimberly Harding Program Officer (Research)</p>	<p>Micronutrient Initiative 180 Elgin St, Suite 1000, Ottawa, ON, K2P 2K3 CANADA Tel: +1 613 782 6821 / Fax: +1 613 782 6838 www.micronutrient.org</p>	<p>kharding@MICRONUTRIENT.ORG</p>
<p>Malia Boggs Technical Advisor Child and nutrition program</p>	<p>USAID Office of Health, Infectious Diseases and Nutrition USAID/GH/HIDN/NUT Room 3.07-041, Third Floor, RRB, Washington, D.C. 20523-3700 / Phone: 202-712-1294</p>	<p>mboggs@usaid.gov</p>
<p>Norm Campbell PAHO /WHO Expert group, Chair</p>	<p>Departments of Medicine, Community Health Sciences and Physiology and Pharmacology, TRW, The University of Calgary,3280 Hospital Drive NW, Calgary Alberta, T2N 4Z6 / Tel: 403-210 3955</p>	<p>ncampbel@ucalgary.ca</p>
<p>Prof. Dr.med. Michael B. Zimmermann</p>	<p>Institute of Food, Nutrition and Health Swiss Federal institute of Technology (ETH) Zürich LFV E19, Schmelzbergstrasse 7, CH-8092 Zürich, Switzerland / Tel. +41 44 632 8657</p>	<p>michael.zimmermann@ilw.agrl.ethz.ch</p>
<p>Franco Cappuccio Professor</p>	<p>Cephalon Chair – Cardiovascular Medicine and Epidemiology, Director-WHO Collaborating Centre for Nutrition University of Warwick, Warwick Medical School Clinical Sciences Research Institute CSB bldg. UHCW Campus, Clifford Bridge Road Coventry CV2 2dx, UK</p>	<p>F.P.Cappuccio@warwick.ac.uk</p>
<p>Omar Dary Expert in micronutrients PAHO /WHO Expert group, leads sub group on fortification issues</p>	<p>Academy for Educational Development. U.S. OFFICES AED Headquarters 1825 Connecticut Ave., NW, Washington DC / Tel: 202-884-8436.</p>	<p>odary@aed.org</p>
<p>Mary L'Abbé Professor and Chair</p>	<p>Department of Nutritional Sciences Faculty of Medicine, University of Toronto Fitzgerald Bldg. 150 College Street Toronto, ON, Canada M5S 3E2 Tel: (416) 978-7235</p>	<p>Mary.Labbe@utoronto.ca</p>
<p>Barbara Legowski Consultant, PAHO Secretariat salt initiative</p>	<p>17 Loch Isle Rd Ottawa, ONK 2H8G5, Canada / Tel(613) 726 2625</p>	<p>legowski@rogers.com</p>
<p>Eduardo Pretell ICCIDD Coordinator for Americas</p>	<p>Avenida Paseo de la República 3691 Oficina 401-A, San Isidro, Lima 27, Perú</p>	<p>dreapretell@gmail.com</p>
<p>Arun Chockalingam Director, Global Health</p>	<p>National Heart, Lung, and Blood Institute Office of Global Health 31 Center Drive Suite 5A06D, Bethesda, Md. 20892 Tel: 301-496-3620</p>	<p>chockalingama@mail.nih.gov</p>

James Hospedales Senior adviser, NC coordinator	Pan American Health Organization Area of Health Surveillance and Disease management 525 23rd street, NW / Phone: 202 974 3695	hospedaj@paho.org
Lynnette Neufeld	Micronutrient Initiative 180 Elgin St, Suite 1000, Ottawa, ON, K2P 2K3 CANADA Tel: +1 613 782 6821 / Fax: +1 613 782 6838 www.micronutrient.org	lneufeld@MICRONUTRIENT.org
Godfrey Xuereb Team Leader (Population-based Prevention)	WHO- HQ Surveillance and Population-based Prevention Unit Department of Chronic Diseases and Health Promotion Tel. direct: +41 22 791 2617 / Fax direct: +41 22 791 1581	xuerebg@who.int
Darwin Labarthe Director, Division for Heart Disease and Stroke Prevention, NCCDPHP	Centers for Disease Control and Prevention Division for Heart Disease and Stroke Prevention National Center for Chronic Disease and Health Promotion Atlanta 770-488-8053, phone / 770-488-8334, FAX	dil3@cdc.gov
Kevin Sullivan Epidemiologist/Associate Professor	Centers for Disease Control and Prevention Division of Nutrition, Physical Activity and Obesity National Center for Chronic Disease Prevention and Health Promotion, Atlanta, GA And Department of Epidemiology, Rollins School of Public Health at Emory University, Atlanta, GA 30322 404-727-8714, phone / 404-727-8737, fax	cdckms@emory.edu
Branka Legetic PAHO Secretariat salt initiative	Regional adviser, noncommunicable disease prevention and control Pan American Health Organization Area of Health Surveillance and Disease management 525 23rd street, NW / Phone: 202 974 3892	legeticb@paho.org
Christine Swanson Senior Nutrition Scientist	National Institute of Health Office of Dietary Supplements 6100 Executive Blvd. Room 7517 / Tel: 301-435-2930	Swansonc@od.nih.gov
Ruben Grajeda Regional adviser, Micronutrient program	Pan American Health Organization Family and Community Health area 525 23rd street, NW / Phone:	grajedar@paho.org
Kathleen L. Caldwell Chief, Inorganic Toxicology Laboratories	National Center for Environmental Health Center for Disease Control 4770 Buford Hwy. Mail stop F-18 / Tel: 770-488-7990	Klc7@cdc.gov
Mary E. Cogswell RN, Dr PH Senior Scientist	Centers for Disease Control and Prevention Epidemiology and Surveillance Branch Division for Heart Disease and Stroke Prevention National Center for Chronic Disease and Health Promotion 770-488-8053, phone / 770-488-8334, FAX	mec0@cdc.gov
Catherine (Cay) Loria PhD, MS, MA, FAHA, Nutritional Epidemiologist	National Heart, Lung, and Blood Institute Division of Cardiovascular Sciences 6701 Rockledge Dr., Suite 10018 Bethesda, MD 20892-7936 FED EX ZIP: 20817 301-435-0702 / 301-480-5158 (fax)	loriac@mail.nih.gov



**Targets Setting and
Voluntary Industry
Engagement**

**A Guide for Setting Targets
and Timelines to Reduce the
Salt Content of Food**

Key Messages

Typical modern diets provide excessive amounts of salt¹. While people can control the salt added at the table and in cooking, it is the food industry that determines how much salt is added during food processing. Engaging the food industry in setting and following a schedule of targets and timelines to reduce the use of salt requires first a secure national commitment to dietary salt reduction.

Countries that intend to set targets for reduced salt content in foods should examine the approaches and achievements of already active countries – evidence of what reformulations are feasible and that progress can be made. In the Americas, as of January 2013, there are targets and timelines for reduced salt in foods in five countries: Argentina, Brazil, Canada, Chile and in the United States the National Salt Reduction Initiative (NSRI).

Canada and the NSRI are accounting for all food categories that contribute salt to the diet. Argentina and Brazil are taking a step-wise approach, beginning with priority food categories. Chile began with voluntary agreements to reduce the salt in bread, followed by a law requiring warning labels on packaged products in which salt and other nutrients exceed upper limits.

This document gives links to the comprehensive strategies in these five countries and presents the targets and timelines for the food categories common to all or most of them as of January 2013: all five countries have targets for bread and bakery products; four are dealing with biscuits and cookies and snacks; three have targets for cakes, meats, dairy products, mayonnaise, soups and pasta.

Ideally, targets should be set both an average and maximum target per category or per sub-category. The two targets affect the range of products in a category, this way protecting more of the population, no matter which products from within the category people consume.

It is vital that progress relative to national targets and timelines, whether voluntary or regulated, be monitored and made public. Asking companies to regularly report the nutrient profiles of their foods in a standardized electronic format to a central database can facilitate monitoring.

Countries should also assess from time to time their progress against what other countries in the Region and elsewhere are achieving and should monitor advances in food technology that may offer new reformulation options.

1 While sodium is the technically and scientifically correct term for the nutrient of public health concern, the term salt is used in this document. Whether a country refers to salt or sodium in its national discourse on the nutrient is a matter of national discretion.

Equivalencies: 5 gm salt (NaCl) = 2000 mg sodium (Na) = 87 mmol sodium = 87 mEq sodium

Introduction

Typical modern diets provide excessive amounts of salt, from early childhood through adulthood. In high income countries processed foods account for most dietary salt intake, and in lower and middle income countries their consumption is rapidly increasing as nutrition transition evolves. While people can control the salt added at the table and in cooking, it is the food industry that determines how much salt is added during food processing. A core component of any national program to reduce dietary salt consumption is government working with the food industry to lower the amount of salt added during food processing. Governments are justified in intervening directly with the food industry because salt is such a common food additive and salt intake levels are consistently high. The food companies that participate in reducing the salt content of their products demonstrate corporate social responsibility and stand to benefit from a positive public image. As more companies participate in such health promoting activity, the potential grows for a shift in the whole food market towards improving the nutrition profile of processed foods.

While a number of food companies have ongoing initiatives to reduce the salt content in their products, and while their product development schedules should be taken into consideration when setting national targets and timelines for reducing the use of salt, experience to date has shown that when governments and the food industry coordinate their efforts in a national strategy to reduce the overconsumption of salt, there have been larger and more timely reformulations of food products.

Purpose of the Guide

Audience

This is a step-wise Guide intended to assist governments and public health authorities with initiating and maintaining interactions with food companies to set and then monitor targets and timelines for gradually lowering the salt content of processed foods. It has been developed by representatives of public health authorities in countries in the PAHO region that have committed to reducing dietary salt intake and are involved with the food industry to stimulate product reformulations.

Primary aims

- give advice and recommendations on how to initiate and maintain the involvement of food industries and other relevant agencies and entities in a transparent and open process of developing and monitoring a schedule of targets and timelines intended to reduce the amounts of salt added to specific food categories.

- disseminate the targets and timelines for food categories in common as reported as of January 2103 for five countries (Appendix 1) – Argentina, Brazil, Canada, Chile and the US National Salt Reduction Initiative (NSRI) – to inform countries that intend to reduce the overconsumption of salt of what others are seeking to achieve. Appendix 1 has a compilation of:
 1. the food categories identified for salt reduction, common to all or most of the five countries;
 2. for each food category above, the targets and timelines set to date in each country.
- An on-line version of Appendix 1 is to be updated as progress is made.

Secondary aims

- Foster a harmonization of targets for common food categories as more countries engage the food industry and the lead countries demonstrate successes.
- Provide links to the national salt reduction program details in the five Pan American countries with targets and timelines as of January 2103, presented in Appendix 2.

Step 1 – Secure the National Strategy to Reduce Dietary Salt

Establish the national importance of dietary salt reduction

Salt intake exceeding physiologically adequate levels has a causal and linear relationship with greater-than-optimum levels of blood pressure – the science is incontrovertible. Useful references and resources with evidence and justifications for a national dietary salt reduction initiative include:

- the Political Declaration of the High-level Meeting of the [UN] General Assembly on the Prevention and Control of Non-communicable Diseases (2011) at http://www.un.org/ga/search/view_doc.asp?symbol=A/66/L.1;
- the PAHO initiative on Cardiovascular Disease Prevention through Dietary Salt Reduction at http://new.paho.org/hq/index.php?option=com_content&view=article&id=2015&Itemid=4024&lang=en
 - core scientific and policy references at http://new.paho.org/hq/index.php?option=com_content&view=article&id=3043&Itemid=2375
- *WHO Guideline: Sodium intake for adults and children* at http://www.who.int/nutrition/publications/guidelines/sodium_intake_printversion.pdf

WHO guidelines for sodium and potassium intake at http://www.who.int/nutrition/publications/guidelines/sodium_intake/en/index.html and http://www.who.int/nutrition/publications/guidelines/potassium_intake/en/index.html

- US Institutes of Medicine report *Strategies to Reduce Sodium Intake in the United States* (2010) at <http://www.iom.edu/Reports/2010/Strategies-to-Reduce-Sodium-Intake-in-the-United-States.aspx>

Prepare the arguments, supported by epidemiologic data, as to the national importance of dietary salt reduction. Collect data on mortality and morbidity due to risk factors related to high salt diets, and if available, also the economic burden related to them. Consider also determining the health economic benefits of reduced salt intake. Examples of methodologies are in:

- Wang G, Labarthe D. The cost-effectiveness of interventions designed to reduce sodium intake. *J Hypertens.* 2011; 29:1693–99.
- Cobiac LJ, Vos T, Veerman JL. Cost-effectiveness of interventions to reduce dietary salt intake. *Heart.* 2010; doi:10.1136/hrt.2010.199240.

Ensure high level political commitment and operational support for the national initiative and publicize it. As timelines for reformulations typically span a number of years, the government initiative must be able to sustain ongoing discussions and negotiations along with performance monitoring for the period.

Develop a comprehensive strategy with a clear goal; indicate whether it will take a voluntary or regulatory approach

Referring to international experiences and achievements, define a comprehensive strategy for dietary salt reduction. The strategy can begin by setting a national goal for salt intake, as a number of countries in the Americas have done, or in its absence, adopting:

- the target recommended by the PAHO Expert Group – less than 5g salt intake/day/person by 2020 or
- the WHO global target – a 30% relative reduction in mean population intake of salt, aiming to achieve the recommended level of less than 5g salt per adult per day.

Governments can follow several different approaches for setting targets for reduced salt in foods. They can focus on the whole food supply (e.g. the food processing industry, restaurants, food caterers, the informal

food sector), on specific sectors of the food supply and all their respective food categories (e.g. processed food manufacturers), or on specific components of the food supply or key food categories (e.g. breads, processed meats). Regardless of which sector of the food supply or which food categories are being addressed, the approach must include a monitoring framework that assesses industry performance against commitments made to targets and brings feedback to the next stages of reduction. See Section 6 Monitoring Performance. Whether voluntary or regulated, successful initiatives have strong government oversight. To be avoided is relying solely on the food industry to self-monitor. See

- US Institutes of Medicine report *Strategies to Reduce Sodium Intake in the United States* (2010) at <http://www.iom.edu/Reports/2010/Strategies-to-Reduce-Sodium-Intake-in-the-United-States.aspx>
- Moodie R, Stuckler D, Monteiro C, Sheron N, Neal B, Thamarangsi T, Lincoln P, Casswell S. Profits and pandemics: prevention of harmful effects of tobacco, alcohol, and ultra-processed food and drink industries. *The Lancet*. 2013; 381:670-79.

The voluntary approach undertaken to date by several jurisdictions is based on negotiations with food industries to establish reduction targets, often complemented with a voluntary commitment mechanism. Most have committed to gradual reductions of salt content per food category. Success depends on:

- genuine commitment by the food industry to reach targets;
- the strong position of the government to set and hold to the lowest feasible targets possible, and
- government commitment to monitor progress.

A regulatory approach typically sets an upper limits of salt in foods and depends on enforcement. It has been most commonly used for widely consumed products such as breads. An advantage of the voluntary approach over the regulatory one is that it can be initiated relatively rapidly in most countries that have basic national data on salt intake and on the main food sources of salt, and it retains some flexibility for adjusting reduction targets over time. On the other hand, a regulatory approach has been assessed as more effective in reducing salt consumption and for industry, levels the playing field across food processing sectors.

It is also possible for voluntary approaches to be followed by legal measures. If this is the intention from the outset, the food industry is initially prompted to respond to voluntary targets and is in a ready position once regulations are passed. This may also be an outcome in cases where voluntary targets are not being

fully achieved by industries or when small but cumulatively significant segments of the food market have not responded to voluntary targets.

Chile began with voluntary agreements with industry to gradually reduce the salt content in food, in particular in bread made in small bakeries and by supermarket chains. Most recently it has also enacted a Law on Nutrient Composition of Foods and Advertising (N°20,606, 2012) that requires a warning label on food packages where the salt and other nutrients exceed upper limits. The intention is to encourage the food industry to improve the overall nutritional quality of food products. See Appendix 2 for links to the Chile approach.

Leverage and coordinate with supporting initiatives, agencies and resources

Position dietary salt reduction within the larger national public health agenda

Look for a convergence of interests, where concerted dietary salt reduction can strengthen other national policies, instruments and processes already in place that refer directly to salt e.g. national nutrition guidelines or the prevention of non-communicable diseases e.g. cardiovascular and renal diseases, hypertension.

Wherever possible, link to other diet related risk factor programs e.g. obesity, school health, trans fatty acid elimination, where the health risks of high salt can be profiled.

Coordinate with the national salt iodization program

It is crucial to check and account for impacts of reduced dietary salt on the prevention of iodine deficiency disorders where salt delivers supplementary iodine (I). In some cases, household table salt is the sole source of the supplement whereas in other cases, both table salt and the salt used in food processing are iodized. See http://new.paho.org/hq/index.php?option=com_content&view=article&id=2015&Itemid=1757 for the *White Paper on Improving Public Health by Optimizing Salt and Iodine Intakes, 2011* and the *Final Report on Improving Public Health in the Americas by Optimizing Sodium and Iodine Intakes – A Meeting Summary*.

Secure food labelling

Clear easy-to-read food labels can be a central tool for directing consumers to healthier choices and at the same time are very often a key source of data on the salt content of packaged foods – required for the baselines on which the case is made to reduce salt use and central to monitoring performance against targets. Ideally official food composition tables are the sources of nutrient information about national foods. If tables are not up to date, label data become a key source of

nutrient contents. For label data to be reliable to serve both consumers and nutrition policies, a system to confirm the accuracy of labels needs to be in place including feedback and enforcement of requirements.

The regulatory framework governing food labels needs to respond effectively and efficiently to the pace of food composition changes. Examining the framework may open opportunities to consider other tools or incentives to lower salt use e.g. additional nutritional information, health claims, health seals, warning labels to alert consumers (e.g. foods with high salt content). These additional sources of information can be a useful complement to nutrition labels.

Consult food technology experts

The food technology sector and food research centres are a particularly important resource for governments to learn about opportunities and constraints regarding food product reformulations e.g. typical reformulation schedules, technology or food safety issues related to the role of salt in specific food products that have an impact on reformulating, schedules for new product development, and how these relate to targets and timelines.

Examine/promote healthy food procurement policies

Determine where reformulated products can represent a new market opportunity for food companies. Public entities may have food procurement policies for their cafeterias for which they are responsible and may be in a position to adjust their policies to take into account reduced levels of salt in the food products being purchased or offered for sale. Key examples are the procurement policies for schools, hospitals, long term care facilities, daycares and prisons.

Involve other government ministries and agencies

A cross-government approach must be considered as there may be ministries other than health that have roles and responsibilities affecting the food supply. For example, the agriculture ministry may control meat and dairy processing and where there are separate food regulatory agencies, they will be needed to monitor food labels, test food products or have a role in harmonizing salt content across food categories in cases where food products are the responsibility of multiple ministries.

In cases where countries are federations with authorities and responsibilities for food processing distributed across regional, state or local governments, the food industries will engage through these levels. There may also be resources at regional or state levels that can support national implementation or be coordinated to support local efforts e.g. local laboratories for analyzing salt content.

Countries in common or neighbouring markets across which food products circulate can explore the potential convergences of interests on issues such as food labelling or salt reduction efforts.

Use international references

International references such as the *Codex Alimentarius* have standards for labelling of salt that are parallel to the efforts to reformulate food. The *Guidelines on Nutrition Labelling* (CAC/GL 2-1985) recommend that all pre-packaged foods be labelled with nutrition information including the amount of sodium. The *Guidelines for Use of Nutrition and Health Claims* (CAC/GL 23-1997) set out conditions for claims related to sodium.

Examine trade agreements

Where relevant, understand the trade patterns and agreements regarding the processed foods imported from within a common market or from elsewhere, and the extent to which your country has control over these products e.g. nutrition labelling requirements. Consider international or trade agreement levers that can protect national public health in cases of imported products.

Examine also the salt reduction approaches taken in other countries with which there is some relevant relationship e.g. neighbouring country, country with similar food culture, similar economic status (see Appendix 2).

Control advertising to children

It is especially important for all stakeholders in a healthy food supply to be consistent in supporting the public to make healthy food choices. This includes how and what foods are advertised and especially the nature of advertising to children as they are a vulnerable population. As some products intended for children have high salt content, the World Health Organization recommendations for marketing of foods and non-alcoholic beverages to children are relevant to dietary salt reduction. See <http://www.who.int/dietphysicalactivity/marketing-food-to-children/en/index.html>

Monitor the commitments of transnational companies

Appendix 2 includes links to public commitments regarding salt content made by transnational and large national food companies in the Americas as of January 2013.

An international initiative that is engaging a number of transnational food companies in the region is the Salt Smart Consortium that operates through the Pan American Forum for Action on NCDs. Country-based or sub-regional branches of the multinationals may be predisposed to participate in a national initiative. See <http://new.paho.org/panamericanforum/?p=626>.

Consider warning labels with or in the absence of targets

Where systems of warning labels are mandatory, review that salt is featured. Warning labels are a means of complementing reformulation commitments, especially those set as maximum targets. Public health authorities set an upper limit of salt per standardized unit or standardized serving of food product, that if exceeded, requires that consumers be warned. Refer to Appendix 2 for links on warning label legislation in Chile and Appendix 3 for the pros and cons of warning labels.

Step 2 – Prepare Data

There are a number of sets of data necessary to initiate engagement with the food industry. A government must prepare some, and depending on its capacity and resources, can prepare the rest or request them from the food industry or from other private commercial sources. In cases where the food industry supplies data on the salt content of its food products, government must provided with the data in a way such that it can be verified as accurate, prior to negotiations and during monitoring of progress.

As the objective for data preparation is a set of priority food products with proposed targets and timelines for salt content reductions, the criteria for selection of the categories and targets must be transparent to all stakeholders and grounded in evidence that the proposed reductions in salt content are technically feasible and will achieve measurable reductions in salt intake.

Select the food categories and determine baseline salt content

National food consumption data are very important as they identify the foods and eating habits that account for salt intake levels. The consumption data coupled with data on the salt content of products in the food supply (from national or international databases) point out the relative contributions of various foods to total salt intake, hence the products on which reformulations should be focussed and the sectors of the food industry to approach can be established.

Important food consumption data are:

- foods that people eat and the amounts and frequency of consumption;
- salt content of the most commonly consumed foods;
- the amount of salt added at the table and in cooking;
- intake of high salt foods that are culturally or regionally specific within the country.

Select food categories based on national data on the

sources of salt in the diet and on knowledge of the feasibility of salt reductions.

- The proportion of total salt intake attributed to a food category can be derived by collecting food intake data (estimated through various instruments e.g. 24 hour food recalls or household budget surveys) which is then matched with appropriate food composition data that include the salt content for the products identified. Food composition data can come from either nationally specific or regional databases containing foods with some regional specificity e.g. LATINFOODS (Tabla de Composición de Alimentos de América Latina at <http://www.into.cl/latinfoods/>).
- It is critical that any sources used to determine the salt content of foods be current. (Alternatives to food composition data that include salt content are food labels, food analysis or values provided by the food industry).
- National food intake data can also be derived from FAO food balance sheets.
- A possible data source is private commercial enterprises that collect and sell product market share data. While they can supply precise and comprehensive information as to all the brands in each food category and their relative share of market, necessary to calculate sales weighted averages for the salt content per food category, it is proprietary information and costly. As well, depending on the country, data may be limited in terms of market coverage and the level of detail needed.
- Give attention to processed foods even in cases where discretionary salt use is high as nutrition transition towards processed foods is accelerating especially in countries with rising economies.

The PAHO document *A Review of Methods to Determine the Main Sources of Salt in the Diet*, available at http://new.paho.org/hq/index.php?option=com_docman&task=doc_download&gid=11559&Itemid= is a comprehensive guide to identify key food categories and salt content of foods. The FAO/INFOODS food composition databases is one international source for food categories and provides salt content data as well, available at <http://www.fao.org/infoods/infoods/tables-and-databases/faoinfoods-databases/en/> Knowing the salt levels in locally available food products also allows comparing the levels in same or similar foods in other parts of the world, identifying in particular the global brand products for which lower salt versions are available elsewhere. This is a lever for encouraging multinational companies to replicate product improvements in your country. Refer to the work on global brand products by the Global Food Monitoring Group at <http://www.georgeinstitute.org/global-health-landscape/food-policy/food-monitoring-group>

and <http://www.georgeinstitute.org/global-health-landscape/food-policy/food-monitoring-group/resources> High income countries in the region have chosen to set targets and timelines for the full range of processed foods that contain added salt. See Appendix 2 for links to the comprehensive sets of food categories with targets and timelines in Canada and the US NSRI. In these cases processed foods account for an estimated 75% of total salt consumed and public health authorities have been able to allocate resources sufficient to address all of these products.

Other countries in the region are taking a step-wise approach, beginning with salt reductions in priority food categories, selecting those that:

- account for a large proportion of total salt intake;
 - a food category with a high volume of consumption e.g. bread or a product within a category that holds a large percentage of market share, even though not high in salt, may account for a large proportion of total salt intake, while other foods, with relatively low volumes of consumption, may also contribute large amounts of salt to the diet because of their very high salt content (e.g. cured meats, soya sauce).
- have been shown to be highly consumed by vulnerable groups especially children;
- have high baseline salt levels;
- affect the supply chain beyond what is sold to individual consumers e.g. wholesale pre-mixes such as used by bakeries for bread and other baked goods; soup bases supplied to restaurants;
- have food technology solutions for reducing salt content.

Selection of priority food categories can also be influenced by what other countries have already selected, particularly countries in a common market or countries from which products are imported. A key example is bread – all countries in the region with targets and timelines are addressing bread, ranging from artisanal bakery bread (in Argentina, Brazil and Chile), to private label brand supermarket bread (Chile), to packaged breads in Canada and the US NSRI.

Other considerations in category selection are whether the sector associated with a food category has capacity to reformulate and whether there is an existing relationship with a sector, established through previous public health interventions, where there is willingness to participate.

Propose a schedule of targets and timelines for discussion

Ahead of discussions with the food industry, public health authorities should draft targets and timelines for

priority food categories. The criteria selected to guide the process must be clear and transparent to the food industries as well as civil society.

Examine first the baseline and ranges of salt content and targets per food category that have been agreed to to date e.g. Appendix 1 for the Pan American region. This can provide evidence of the technical feasibility of reformulations, that technical and food safety issues have been dealt with. Acknowledging these factors during negotiations will better ensure that reductions agreed to are sustainable and meaningful. If for example baseline salt content for a particular food category or product in your country is the same or similar to that found in another country, particularly a neighbouring country or trading partner, consider proposing a schedule of reduction targets and timelines similar to those already negotiated.

Ideally target values should be modelled to demonstrate that they lead to the achievement of the national salt intake goal or are otherwise based on criteria that ensure that proposed reductions will have a meaningful impact on salt intake at a population level. This is to be balanced by the targets and timelines being feasible.

Set targets as averages/ means, or maxima; ideally as both

Consider first if targets are to be set as an average/ mean or as a maximum level of salt per food category or product. Ideally both should be used. Averages address the most popular foods within a category, more so if sales weighted averages (SWA) are used. Whether simple or sales weighted, averages allow flexibility in the levels of salt in different products within a category and are helpful in considering the natural variation in salt levels of some foods (as cheeses and cured meats). On the other hand, averages are difficult for consumers to understand and for national authorities to compare and monitor across individual products.

Maximum or upper levels of salt are straightforward and transparent, easier to administer, monitor and use to compare salt in products. They ensure meaningful salt reductions in products with the higher amounts of salt in the category. They do not however indicate whether the salt content range in foods is changing, making it difficult to predict the impact of targets on average intakes. In the long run maxima may even suggest that as long as salt content is at or below the maximum, no further reductions are needed.

Ideally, set both an average and maximum target per category. Companies will work towards the average and in addition will be guided by the maximum level. The two targets affect the range of products in a category, this way protecting more of the population, no

matter which products from within the category people consume. If a category has very wide variations in salt content, consider, if resources permit, creating sub-categories to recognize the differences with separate targets. **Refer to Appendix 3 for a thorough discussion of pros and cons of averages, sales weighted averages, maxima, setting percentage reductions and requiring warning labels.**

If determining average salt content is not feasible, set maximum targets with timelines for food categories or products.

For averages/means

- check existing average/mean targets for salt content in other countries (Appendix 1);
- determine the average/mean salt content per standard unit of measure for a food category and the salt content distribution;
- if necessary remove outliers in a consistent fashion.

For maxima

- check existing salt content maxima in other countries (Appendix 1);
- set maxima as between the 50th and 75th percentile of the salt content distribution;
- adjust downward as salt content is reduced based on the range of salt content in different products (i.e. serial reductions with maxima set at the “rolling” 75th percentile).

Consider sales weighted averages/means (SWA/SWM) where available

- check existing SWA/SWM for salt content of food products for Canada and the US NSRI (Appendix 1);
- SWA/SWM are calculated using the salt levels of products within a category weighted by their volume of market share in kilograms;
- requires purchasing expensive market share data from the private sector; however these data cannot be published or released publicly;
- for monitoring whether targets are being met, requires ongoing purchase of the market data.

Note that there may be food categories for which salt substitutes might be necessary in order to meet targets or where lowering salt levels beyond a certain target creates a food safety risk. Seek the objective and independent advice of food technology experts on such issues and if needed, consider supporting or encouraging research on the roles of salt in these cases.

Alternatively, consider percentage reductions from a baseline salt content in the products in a food category. The WHO global target of a 30% relative reduction in

mean population intake of salt is an overall guide. This still requires knowing the sources of salt in the diet and the impact of various reductions in salt by source. If for example 50% of salt intake is attributed to discretionary use at home and 50% comes from processed foods, then if all reduction is to be achieved by lowering the salt content of only processed foods, these foods would have to have salt content lowered by 60%.

Standardize the presentation of salt content

To facilitate the reduction efforts by industries and for consumers to know how much salt they get from a food product, it is important to standardize how salt content is presented.

- It is recommended that targets be expressed as salt per 100g or 100ml of product.
- If salt content is expressed per serving size or portion, standard reference amounts should be indicated.
- For products that require reconstitution before consumption (e.g. soup cubes), indicate whether the target refers to the product “as sold” or “as consumed” per 100g or 100ml (once it has been prepared per the manufacturer’s instructions).

Propose timelines

The timeline should be such that the national salt intake goal or the internationally recommended goal can be reached within six to 10 years. Consider the initial targets to span four years with interim targets at e.g. one or two year intervals to underpin monitoring and timely feedback. Discussions may be reopened within the timeframe and targets adjusted as advances are made or technical barriers encountered. If taking a regulatory approach, the timeline will reflect compliance dates. Ensure that proposed timelines are grounded in experiences proven successful elsewhere and/or are based on evidence of achievability supported by the food technology sector.

Step 3 – Identify the Key Stakeholders

Effective dietary salt reduction at the population level requires a multi-sector approach. The greatest impact will be achieved when governments, the food industry, civil society and non-governmental organizations (NGOs) coordinate their efforts. It is essential that all participants in the position of negotiating on behalf of or supporting the public health authorities declare no conflicts of interest.

Critical is that all agencies involved in dietary salt re-

duction efforts publicly state their support for the approach of setting targets and timelines and where relevant, deliver consistent messages and education to the food industry and the public.

Outline the process and principles of engagement with the food industry

There must be a clear understanding and agreement that the selection of food categories, targets and timelines are to make a meaningful impact on salt intake at the population level.

Consider setting terms for technical cooperation. It must be understood that commitments include acceptance of a transparent monitoring framework, that agreements will be documented and official. Consider a process of formal signing of agreements that include roles and responsibilities, targets and timelines and consequences of not meeting targets.

There also needs to be agreement that industry commitments and performance are to be made public, the details of which can be negotiated. Broadcasting the agreements, new targets and industry progress through for example press releases or other means are opportunities to publically reinforce information and education messages and also reinforce the salt reduction policies within the government and to the food industry. This can also be positively linked to building the market for low/no salt products, to the industry's advantage.

The timeframe within which agreements on targets and timelines are expected must be clarified.

Select stakeholders in the food industry

Data on the main sources of salt will direct the selection of food industries to engage in product reformulations. Begin with representatives of food categories that account for the highest market coverage and highest potential effect on population intake of salt. Ultimately all food categories should be addressed and the responsible companies invited to participate in reformulation.

All food companies with products in a food category should be involved to ensure that they are equitably responsible for salt content reductions and to avoid market distortions and differential consumer preferences – issues that are viewed as threats by the food industry.

Umbrella food industry organizations or single large companies can be approached. If for example food markets are highly concentrated, negotiating with individual companies may be fast and effective but caution needs to be exercised to avoid government associa-

tion with any particular company or brand to appear as an endorsement. In more fragmented markets, dealing with a food industry association, if it represents the majority of the sector (based on either number of companies or market share), will reach more companies simultaneously although it may take more time to agree on targets. If there is less than optimal coverage of a sector by an association, consider approaching sub-associations or undertake outreach that can connect outliers to the initiative.

Associations of importers may be especially important to engage in countries where a significant proportion of the food supply is imported.

Take advantage of existing contacts and relationships with food industry and of any relevant initiatives that are currently involving or have involved the food industry e.g. trans fatty acid elimination, fortification of bread.

Engage both manufacturers and retailers per food category where relevant.

Include other government ministries and agencies.

Refer to Section 3 regarding leveraging and coordinating with other government entities. Where other government departments or agencies outside the health sector are involved in the production or regulation of food preparation, it may be advantageous to involve them in discussions with the food industry in order to best understand issues or constraints specific to the sector. Partnerships with these departments or agencies can also be useful in addressing research needs or identifying regulatory hurdles.

Engage NGOs

NGOs can both legitimize and support the national goal and the target setting processes. They can play a key role in balancing the public discourse regarding what consumers can do to reduce salt intake and what the food industry is able to do. They can also extend the reach of consumer awareness and education campaigns through their local and national networks and can offer champions to reinforce them.

Consumer organizations can be particularly helpful if not integral in monitoring whether food companies meet the commitments they make and can strengthen the government position to maintain the momentum to set and reach targets.

Societies of health professionals can also assist with raising awareness, disseminating information, advocating action on behalf of patients and encouraging industry commitments.

Step 4 – Plan the Meetings

The design and formats of meeting between governments and food industries will vary from country to country. Much depends on whether there are pre-existing relationships with the food industry, their nature and the extent of communications. Regardless of the differences between countries, a number of points should be taken into consideration when launching the negotiations on targets and timelines.

Agree on a way of working, clarify expectations

- State a clear purpose for all meetings.
 - Circulate an agenda in advance of each meeting with objectives specified and invite food companies to send in advance specific questions they want addressed during the meeting.
 - At initial meetings, discuss what activities are already underway.
 - If applicable, invite leaders in industry to present current contributions or progress already made.
 - Meet separately with companies whose products are a concern related to high salt content and/or slow progress in reformulation. One-on-one meetings generally provide more detailed information about progress and future plans as well as successes and challenges. The information from individual companies may be useful in understanding the issues for a particular food category.
 - Consider sector-specific meetings, with groups of companies that produce similar products, with the objective of sharing problem solving and addressing common questions related to salt reduction initiatives. Such meetings can result in partnerships of industry groups to jointly work on technical solutions and reporting mechanisms.
 - Trade associations can be useful in presenting generic barriers and technical issues and solutions but due to the proprietary nature of some information, may not be able to elaborate on specific issues.
 - In preparation for meetings, be aware of the range of salt content of company products as well as the range of salt content of the product category and additionally in global brand products in the same food categories. For the latter, see the Global Food Monitoring Group at <http://www.georgeinstitute.org/global-health-landscape/food-policy/food-monitoring-group> and <http://www.georgeinstitute.org/global-health-landscape/food-policy/food-monitoring-group/resources>.
 - Foster as open and honest sharing of information as possible without violating trade secrets.
- Keep brief notes of all key points covered in meetings for future meetings and document challenges and solutions. Circulate to all participants.
 - Where companies have changes in staff or resources, or there are company mergers or acquisitions, monitor progress and follow-up as needed with meetings.
 - Welcome meetings at company offices as there may be opportunities to tour food processing plants that can provide greater understanding of challenges and solutions.
 - Discussions with food companies are most likely to address individual food categories because the function of sodium salts, food safety issues, the processing technologies and consumer thresholds for acceptability may be specific to a category. Expect discussions with food industry representatives to take varying amounts of time according to the food category.
 - There must be the clear understanding that the draft targets and timelines are to be finalized in a timely manner.

Require reformulation schedules

Ask for company-specific reformulation schedules that include the products to be reformulated, by how much and by when. Ensure that there is a focus on products with a larger market share.

Step 5 – Monitor Performance

Industry accountability regarding targets and timelines in either voluntary or regulated contexts must be clarified in terms of a monitoring framework, with the framework included in the agreements made when industry engagements are initiated. It should stipulate what will be monitored, how, at what frequency, how performance results will be disseminated and the consequences in cases where targets are not met.

Propose and agree on a monitoring framework

In general terms, a monitoring framework can include:

- Process outcomes: progress with reformulations including food technology findings; advance reports to inform on initial results; dissemination of experiences/lessons learned; information on consumer preferences.
- Intermediate outcomes: salt content of food products/categories from e.g. laboratory analyses or labels (NB small reductions may not appear on labels as a range of variations in salt content may be tolerated before a label change is required).
- Long term outcomes: salt consumption, blood pressure, etc.

Establish and maintain public attention to the targets

Launch the dietary salt initiative in a public forum where companies will state their commitments and indicate their plans. Plan subsequent events where company progress and achievement of targets can be applauded with media coverage. Document the successes and challenges

Consider the different sources of data for monitoring voluntary approaches or to ensure compliance with regulation

Valid and reliable data on the nutrition composition of foods are central to both public and private interests. In particular consumers need data especially on food labels to be accurate for their decision making.

Data required for monitoring commitments to voluntary targets or to ensure compliance with and enforcement of regulations can come from various sources. Setting baselines and ongoing monitoring of performance against targets requires that data be available on an ongoing basis such that collection can be most readily sustained. This applies equally to countries that require warning labels.

Label data can be a key source for nutrient content information where keeping them updated is mandated and enforced. National food regulatory agencies are a potential resource.

Consider requiring companies to regularly (annually or biennially) report the nutrient profiles of their foods per 100g or 100ml of product (salt and other nutrients as food composition) in a standardized electronic format to a central database.

Other sources of food composition data are food analyses conducted by government agencies, consumer organizations or research institutions.

Engage consumer organizations, scientific societies, health related NGOs

Consider inviting and supporting other organizations to monitor progress against the targets. Supply if possible the data required for monitoring or support the collection of data.

Continue to meet with food industry representatives

Meet regularly with sector representatives to continue to understand the technical issues and barriers being faced as well as learn about the successes in reducing

salt use. If within a category or sector certain individual companies are slower to reduce salt, it may be worthwhile to meet with them separately to review the importance of salt reductions, as well as gain an understanding of their particular issues.

Dealing with Technical Issues, Controversies, Barriers

Food companies are especially concerned about certain issues when discussing salt reduction e.g. that all companies in a sector are involved and committed; there may be need for new technologies; consumer acceptability of reformulated foods; the reductions already made or underway; and food safety. Public authorities that are to be involved with the food industry must make themselves as aware as possible of these issues in advance of meeting with stakeholders and the proposed approach including the targets and timelines should reflect an understanding of the issues.

It is important to stress that for most food categories it is possible to initiate salt reduction through simple salt removal e.g. a series of reductions of less than 10% are generally undetected by consumers. In later stages of reformulation, further lowering of salt can pose greater challenges as there may be need for salt substitutes and/or changes to food processing methods and equipment. For these reasons stepwise approaches have been adopted, to allow reductions to parallel the taste adaptation of consumers and also permit changes in production processes. Discussions with food technologists in advance of meetings with industry, to understand reformulation opportunities and challenges per food category, are recommended.

Use the science and the evidence

- Indicate that the science base on the health advantages of reduced salt intake is solid and frequently updated by internationally recognized scientific and public health organization e.g. American Heart Association (http://my.americanheart.org/professional/General/Cutting-Sodium-to-Prevent-CVD-and-Stroke_UCM_424966_Article.jsp); the Canadian Hypertension Education Program (<http://www.hypertension.ca/chep-recommendations>).
- Do not immediately respond to challenges raised by a single company; first carefully consider information from other companies and other countries, along with that from food technology specialists.
- Should the issue of micronutrient fortification arise, particularly the use of iodized salt to prevent iodine deficiency disorders, see <http://new.paho.org/hq/>

[index.php?option=com_content&view=article&id=2015&Itemid=1757](http://www.georgeinstitute.org/global-health-landscape/food-policy/food-monitoring-group/resources) for the *White Paper on Improving Public Health by Optimizing Salt and Iodine Intakes, 2011* and the *Final Report on Improving Public Health in the Americas by Optimizing Sodium and Iodine Intakes – A Meeting Summary*.

Refer to existing reformulations and targets

- If industry reformulation costs are cited as a barrier, refer to the established targets and timelines in Appendix 1. The existing experiences have proven that there is room for meaningful reductions, particularly in the first stages of reformulation for many food categories where the adjusted levels of salt remain within the acceptable range for taste.
- Reformulation is an ongoing activity for the food processing industry; a substantial proportion of the reformulation costs are part of ongoing business expenses. Companies can accommodate gradually lower salt content targets into product reformulation cycles to coincide with marketing strategies and label changes.
- Be prepared for companies being reluctant to reduce salt content of all brands especially those with global recognition. For the latter, refer to the Global Food Monitoring Group at [http://www.georgeinstitute.org/global-health-landscape/food-policy/food-monitor-](http://www.georgeinstitute.org/global-health-landscape/food-policy/food-monitoring-group)

[ing-group](http://www.georgeinstitute.org/global-health-landscape/food-policy/food-monitoring-group/) and <http://www.georgeinstitute.org/global-health-landscape/food-policy/food-monitoring-group/resources> for data on the salt content in best in class and best in world global brand products and global variations in the same product.

Encourage technology transfer

- Encourage industry, especially small and medium enterprises (SME), to discuss and share technical solutions for reformulations.
- If feasible contribute funds for technical research. There may be government departments or agencies other than health, universities or research centres that can fund or be partner in relevant research.
- Consider supporting food technology development among national SME e.g. through small grants in order for all companies in a sector to be actively reformulating their products.
- Seek public commitments from the food industry for technology transfer and otherwise encourage technology transfer by other means available.

Invoke corporate social responsibility

- Give examples of industry leaders and be willing to highlight the success stories.
- Set mechanisms whereby industry leaders can report on progress.

Appendix 1

Table 1 - Targets and Timelines for Food Categories in Common in Argentina, Brazil, Canada, Chile and the National Salt Reduction Initiative in the United States (As of January 2013).

		Baselines sodium - mg/100g			Sodium Targets mg/100g										
					Average/mean			SWA/SWM			Upper limit				
Food Category	Country and products	average/mean (year)	SWA 2009/2010	upper limit (year)	2012	2014	2016	2012	2014	2016	2012	2014	2016		
Bread and bakery products	Argentina														
	Artisanal bread	920 (2011)		1100 (2011)	805	690									
	Bakery products	1300 (2011)		1600 (2011)	1200	1000									
	Brazil														
	Industrially produced bread and buns														
	Loaf bread	431		796								645	522		
	Buns	524		656								531	430		
	Artisanal bakery (French) bread	648										616	586		
	Canada														
	Pantry bread, rolls, bagels, croissants, flatbread		469						430	380	330				520
	Hearth bread		531						520	490	470				600
	Chile														
	Artisanal bakery bread	780 (2010)					450								
	Private label supermarket bread						450								
	NSRI														
Savory breads and rolls		485						440	360						
Biscuits and Cookies	Argentina														
	Crackers without bran	601 (2011)		907 (2011)	570.95										
	Crackers with bran	781 (2011)		890 (2011)	741.95										
	Sweet without filling	367 (2011)		539 (2011)	348.65										
	Sweet with filling	234 (2011)		451 (2011)	222.3										
Snack cookies	1190 (2011)		3000 (2011)	1130.5											

		Baselines sodium - mg/100g			Sodium Targets mg/100g									
					Average/mean			SWA/SWM			Upper limit			
Food Category	Country and products	average/mean (year)	SWA 2009/2010	upper limit (year)	2012	2014	2016	2012	2014	2016	2012	2014	2016	
Biscuits and Cookies	Brazil													
	Salted biscuits	700		1220							923	699		
	Sweet Biscuits	360		490							419	359		
	Filled cookies	254		600							398	265		
	Canada													
	Cookies		363					320	280	240				390
	Crackers		859					770	690	600				930
	NSRI													
Filled and unfilled cookies, sandwich cookies and tea biscuits		367					310	260						
Crackers		918					780	640						
Cakes	Brazil													
	Cakes without filling	336		463							392	332		
	Filled cakes	250		330							282	242		
	Roulade	205		240							221	204		
	Mixes for aerated cakes	336		568							476	398	334	
	Mixes for creamy cakes	268		412							349	295	250	
	Canada													
	Baked desserts (cakes, doughnuts, muffins, pastries etc)		349					310	270	230				400
NSRI														
Cakes, snack cakes, muffins and toaster pastries		359					310	250						
Meats	Argentina													
	Sausages, ham, morcilla	1218 (2011)		1370 (2011)	1120.56									
	Salami	1680 (2011)		2000 (2011)	1596									
	Chorizos	1750 (2011)		1950 (2011)	1662.5									
	Hamburgers	766 (2011)		1100 (2011)	651.1									
	Chicken	680 (2011)		980 (2011)	625.6									

		Baselines sodium - mg/100g			Sodium Targets mg/100g									
					Average/mean			SWA/SWM			Upper limit			
Food Category	Country and products	average/mean (year)	SWA 2009/2010	upper limit (year)	2012	2014	2016	2012	2014	2016	2012	2014	2016	
Meats	Canada													
		Uncooked bacon-belly		619				610	590	580				610
		Fully cooked breakfast strips and bacon substitutes		961				940	930	910				960
		Uncooked fresh sausage		789				750	700	660				690
		Cooked sausage		990				940	880	830				870
		Cooked deli meats		1028				970	910	850				890
		Dry cured, fermented deli meats		1592				1510	1420	1330				1400
		Canned chicken or turkey		554				520	480	450				470
		Canned meat		865				840	820	790				830
		NSRI												
		Cold cuts		1085				980	810					
		Pepperoni and dry salami		1834				1740	1560					
		Cooked sausage		898				810	720					
		Uncooked sausage		838				750	670					
		Hot dogs		1059				950	850					
		Bacon		1792				1610	1470					
		Uncooked whole muscle meat and poultry		NA								450	400	
		Canned meat and sausage		987				940	840					
		Canned chicken and turkey		403				380	340					
Dairy	Argentina													
		Cream cheese	583 (2011)		650 (2011)	553.85	524.7							
		Danbo cheese	600 (2011)		700 (2011)	570	540							
		Cuartirolo cheese	583 (2011)		700 (2011)	553.85	524.7							
		Tybo cheese	633 (2011)		705 (2011)	601.35	569.7							
		Porsalut cheese	600 (2011)		720 (2011)	570	540							
		Mozzarella	666 (2011)		725 (2011)	632.7	599.4							

		Baselines sodium - mg/100g			Sodium Targets mg/100g									
		average/ mean (year)	SWA 2009/ 2010	upper limit (year)	Average/mean			SWA/SWM			Upper limit			
Food Category	Country and products				2012	2014	2016	2012	2014	2016	2012	2014	2016	
Dairy	Canada													
		Cottage cheese	375					350	330	280			410	
		Cream cheese, cream cheese products, soft unripened goat cheese	472					440	410	350			600	
		Brie, camembert, cheddar, swiss, monterey, jack, brick, colby, gouda, mozzarella	720					710	700	670			770	
		Feta and feta-style	1323					1270	1210	1100			1530	
		Hard cheese grated and ungrated												
		Processed cheese and other cheese products	1610					1520	1420	1240			1670	
		NSRI												
		Grated hard cheese	1530					1450	1300					
		Cheddar, colby, jack, mozzarella, muenster, provolone, swiss	668					630	600					
		Cream cheese	408					390	350					
		Cottage cheese	347					330	290					
		Processed cheese	1393					1250	1040					
	Snacks	Argentina												
		Snacks	1500 (2011)		2000 (2011)	1275								
		Brazil												
		Extruded corn snacks	832		1288							1090	852	747
		Potato chips	548		720							650	586	529
		Canada												
		Chips, popcorn, extruded corn snacks	676					580	490	400				800
		NSRI												
		Flavoured chips	711					570	430					
		Unflavoured chips	524					470	420					
	Puffed corn snacks	969					820	680						

		Baselines sodium - mg/100g			Sodium Targets mg/100g									
					Average/mean			SWA/SWM			Upper limit			
Food Category	Country and products	average/ mean (year)	SWA 2009/ 2010	upper limit (year)	2012	2014	2016	2012	2014	2016	2012	2014	2016	
Mayon- naise	Brazil	1269		1567							1283	1051		
	Canada		760					680	610	530			840	
	NSRI		713					640	570					
Soups	Argentina													
	Soups in cubes	432 (2011)		449 (2011)	410.4	388.8								
	Instant soups	255 (2011)		286 (2011)	242.25	229.5								
	Ready to consume	331 (2011)		358 (2011)	314.45	297.9								
	Cream soups	337 (2011)		355 (2011)	320.15	303.3								
	Canada													
	Bouillon and broth		296						280	260	240			360
	Condensed wet		302						280	260	240			360
	Ready to serve		280						280	260	240			360
	Fresh and instant oriental noodles		309						280	260	240			360
	Dry		388						280	260	240			360
	NSRI													
	Canned		326						280	230				
	Broth and stock		352						320	260				
	Dry soup		820						700	570				
	Pasta	Brazil												
		Instant pasta	2036		4718							1920.7		
Canada														
Shelf stable pasta, noodles and rice or other grains with sauce or seasoned			368						330	300	270			440
NSRI														
Shelf stable dry seasoned pasta and stuffing mix		700 mg/ cup						630 mg/ cup	560 mg/ cup					

Appendix 2

Links to National Targets and Timelines including Industry Commitments made Public

Argentina

Menos Sal + Vida at <http://www.msal.gov.ar/ent/index.php/informacion-para-ciudadanos/menos-sal--vida>

Company commitments at

http://www.msal.gov.ar/ent/images/stories/ciudadanos/pdf/2012-06_listado-empresas-adheridas-Menos-Sal-Mas-Vida.pdf

Brazil

Nilson EAF, Jaime PC, de Oliveira Resende D. “Iniciativas desenvolvidas no Brasil para a redução do teor de sódio em alimentos processados” [Initiatives developed in Brazil to reduce sodium content of processed foods] at http://new.paho.org/journal/index.php?option=com_content&task=view&id=116&Itemid=215

Company commitments at

<http://portalsaude.saude.gov.br/portalsaude/noticia/6839/893/acordo-para-reducao-de-sodio-inclui-novos-alimentos.html>

Canada

Complete Health Canada Data Table at http://www.hc-sc.gc.ca/fn-an/legislation/guide-ld/2012-sodium-reduction-indust_data_table-eng.php

Chile

Estrategia de Reducción de SAL/SODIO en los Alimentos at

http://www.redsalud.gov.cl/portal/url/page/minsalcl/g_proteccion/g_alimentos/reduccion_sodio.html

Programa de reducción de sal/sodio en pan at

<http://www.minsal.gob.cl/portal/url/item/9a0feb846ce5475de04001011e01795f.pdf>

Law No. 20,606 on the Nutrient Composition of Food and its Advertising [la Ley de Composición de Alimentos y su Publicidad] at http://www.europarl.europa.eu/meetdocs/2009_2014/documents/d-cl/dv/ley_20606_comp_alim_/ley_20606_comp_alim_en.pdf

Estudio “Propuesta de Criterios y Recomendación de Límites Máximos de Nutrientes Críticos para la Implementación de la Ley de Composición de Alimentos y su Publicidad” at

<http://www.minsal.cl/portal/url/item/d68cf20e14279b92e0400101650119e3.pdf>

Propuesta a Consulta sobre Modificación del Reglamento Sanitario de los Alimentos (Consultation on Proposed Amendments to the Sanitary Regulation of Food] at

<http://www.minsal.gob.cl/portal/url/item/d2682a13c0994729e04001016501108f.pdf>

National Salt Reduction Initiative in the United States

Complete NSRI Packaged Food Categories and Targets at <http://www.nyc.gov/html/doh/downloads/pdf/cardio/packaged-food-targets.pdf>

Complete NSRI Restaurant Food Categories and Targets at

<http://www.nyc.gov/html/doh/downloads/pdf/cardio/cardio-salt-nsri-restaurant.pdf>

Company commitments at <http://www.nyc.gov/html/doh/html/diseases/salt.shtml>

Appendix 3: Pros and Cons for Types of Targets that Can Be Set

(Key source: European Salt Action Network document “Setting Targets and Industry Engagement”)

Averages	
Pros	Cons
<ul style="list-style-type: none"> Allows some flexibility in the levels of salt in different products within a category, some lower and some higher allowing for different flavours. Usefully applied to foods where there is some natural variation in salt levels e.g. cheeses, cured meats. Useful for companies to use an average target per particular category to monitor its own compliance. 	<ul style="list-style-type: none"> More difficult for consumers and public health agency responsible for dietary salt reduction to compare levels in individual products to monitor compliance May be perceived as unfair by companies that have already lowered the salt content of their products.
<ul style="list-style-type: none"> A simple average is the average salt level in foods within a category regardless of consumption levels; relatively straight forward to derive. 	<ul style="list-style-type: none"> Industry may reduce the salt in the saltiest low selling products in a category thereby reduce the simple average salt level of the category but have no significant impact on actual salt intake.
<ul style="list-style-type: none"> Sales weighted average (SWA) is the “gold standard” – is the average salt level in a food category weighted by the relative sales volumes of products within the category ie indicates product consumption at market level more accurately than the simple average. Monitoring SWA of different food categories combined with food intake survey data allow the most accurate estimates of population intake of salt. 	<ul style="list-style-type: none"> To industry, the SWA may reveal sensitive market-share information. Market share data are prohibitively expensive; must be re-purchased to monitor SWA as changes are made; cannot be published or distributed; an unlikely option for LMIC. Much more difficult to monitor than simple averages in an independent transparent way. Does not address products high in salt. The SWA can be brought down with a focus on a few high volume products and not address targeted products at e.g. children. Monitoring on the basis of SWA would not allow a common approach for the whole food supply as some sectors e.g. large restaurant chains, food services industry do not provide sales volume data for menu items.
<ul style="list-style-type: none"> Encourage more action by companies as more products likely need to be reformulated. 	
<ul style="list-style-type: none"> Based on UK modeling, an average target in the majority of cases requires far less reduction in salt content in a food category than a maximum to achieve a similar overall reduction in salt intake. 	

Maxima	
Pros	Cons
<ul style="list-style-type: none"> • Sets a clear ceiling for all products in a category. • Set at around the average for a food category at point in time and reduce as average shifts down. 	<ul style="list-style-type: none"> • Suggests that if a food product is at or below the maximum, no further action is needed; no incentive for products to have salt levels less than the maximum. • Companies whose products have salt levels initially below the maxima have no incentive to reduce salt content. • Undermines adoption of best in class/best in world that are based on SWA.
<ul style="list-style-type: none"> • Easy for consumers and food companies to compare products. 	
<ul style="list-style-type: none"> • Moves salt content of products above the maxima down to the maxima. 	<ul style="list-style-type: none"> • If high salt content products are not high volume regarding consumption, may have little or no impact on actual salt intake levels. • Companies whose products are below the maximum do not participate.

Setting percentage reductions	
Pros	Cons
<ul style="list-style-type: none"> • May be easier for the food industry. • Combine with maxima (averages for food categories at point in time). 	<ul style="list-style-type: none"> • Still need data on before and after levels of salt content for monitoring purposes.

“Low” and “high” salt labeling of foods	
Pros	Cons
<ul style="list-style-type: none"> • Requires setting of maximum levels. • Conversely need to determine what is “low salt”. 	<ul style="list-style-type: none"> • Must be compulsory (regulatory mechanism). • Requires resetting of maximum levels as salt content falls.

Acknowledgements

Mary L'Abbé,

Department of Nutritional Sciences, University of Toronto, Canada

Pedro Acuna,

Departamento Alimentos y Nutrición, Ministerio de Salud, Chile

Álvaro Flores Andrade,

Departamento Alimentos y Nutrición, Ministerio de Salud, Chile

Simon Barquera,

Instituto Nacional de Salud Pública, Mexico

Norm Campbell,

Libin Cardiovascular Institute of Alberta, University of Calgary, Canada

Critian Cofre,

Departamento Alimentos y Nutrición, Ministerio de Salud, Chile

Maria Cristina Escobar,

Departamento Enfermedades No Transmisibles, Ministerio de Salud, Chile

Daniel Ferrante,

Dirección de Promoción de la Salud y Control de Enfermedades No Transmisibles, Ministerio de Salud, Argentina

Charmaine Kuran,

Food Directorate, Health Canada

Hubert Linders,

Fundraising Officer and Project Coordinator, Regional Office for Latin America and the Caribbean, Chile

Eduardo Nilson,

Coordenação de Alimentação e Nutrição, Ministério da Saúde, Brazil

PAHO Secretariat:

Branka Legetic,


Area of Health Surveillance and Disease Management

Barbara Legowski,

Technical Advisor

Christine M Campbell,

Intern



9 789275 117699

