

# Oral Health Status of Older Adults in the Americas





# Oral Health Status of Older Adults in the Americas

Washington, D.C., 2023



**Decade of Healthy Aging in the Americas**  
**situation and challenges**

## Oral Health Status of Older Adults in the Americas

ISBN: 978-92-75-12663-9 (PDF)

ISBN: 978-92-75-12664-6 (Print version)

© Pan American Health Organization, 2023

Some rights reserved. This work is available under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 IGO license (CC BY-NC-SA 3.0 IGO).



Under the terms of this license, this work may be copied, redistributed, and adapted for non-commercial purposes, provided the new work is issued using the same or equivalent Creative Commons license and it is appropriately cited, as indicated below. In any use of this work, there should be no suggestion that the Pan American Health Organization (PAHO) endorses any specific organization, product, or service. Use of the PAHO logo is not permitted.

**Adaptations:** If this work is adapted, the following disclaimer should be added along with the suggested citation: “This is an adaptation of an original work by the Pan American Health Organization (PAHO). Views and opinions expressed in the adaptation are the sole responsibility of the author(s) of the adaptation and are not endorsed by PAHO.”

**Translations:** If this work is translated, the following disclaimer should be added along with the suggested citation: “This translation was not created by the Pan American Health Organization (PAHO). PAHO is not responsible for the content or accuracy of this translation.”

**Suggested citation:** Pan American Health Organization. Oral Health Status of Older Adults in the Americas. Washington, D.C.: PAHO; 2023. Available from: <https://doi.org/10.37774/9789275126639>.

**Cataloguing-in-Publication (CIP) data:** Available at <http://iris.paho.org>

**Sales, rights, and licensing:** To purchase PAHO publications, email: [sales@paho.org](mailto:sales@paho.org). To submit requests for commercial use and queries on rights and licensing, visit: <https://www.paho.org/en/publications/permissions-and-licensing>.

**Third-party materials:** If material that is attributed to a third party, such as tables, figures, or images, is reused from this work, it is the user’s responsibility to determine whether permission is needed for that reuse and to obtain permission from the copyright holder. The risk of claims resulting from infringement of any third-party-owned material or component from this work rests solely with the user.

**General disclaimers:** The designations employed and the presentation of the material in this publication do not imply the expression of any opinion whatsoever on the part of PAHO concerning the legal status of any country, territory, city, or area or of its authorities, or concerning the delimitation of its frontiers or boundaries. Dotted and dashed lines on maps represent approximate border lines for which there may not yet be full agreement.

The mention of specific companies or of certain manufacturers’ products does not imply that they are endorsed or recommended by PAHO in preference to others of a similar nature that are not mentioned. Errors and omissions excepted, the names of proprietary products are distinguished by initial capital letters.

All reasonable precautions have been taken by PAHO to verify the information contained in this publication. However, the published material is being distributed without warranty of any kind, either expressed or implied. The responsibility for the interpretation and use of the material lies with the reader. In no event shall PAHO be liable for damages arising from its use.

FPL/HL/2023

# Contents

<b>Abbreviations and acronyms</b> .....	<b>iv</b>
<b>Acknowledgments</b> .....	<b>v</b>
<b>Aging and oral health</b> .....	<b>1</b>
<b>Oral health status</b> .....	<b>4</b>
Country data .....	4
Coronal dental caries .....	8
Root caries .....	9
Periodontal diseases .....	9
Tooth loss .....	12
Edentulism .....	13
<b>Utilization of dental services</b> .....	<b>14</b>
<b>Discussion</b> .....	<b>17</b>
<b>Gaps in knowledge and methods</b> .....	<b>18</b>
<b>Conclusions</b> .....	<b>19</b>
<b>References</b> .....	<b>21</b>

## List of Tables

Table 1. Dental caries, root caries, periodontal status, number of teeth, and edentulism among older adults in selected countries of the Americas .....	4
---	---

## Abbreviations and acronyms

CPI	community periodontal index
CPITN	community periodontal index of treatment needs
DALY	disability-adjusted life year
DMFT	decayed, missing, and filled teeth
PAHO	Pan American Health Organization
WHO	World Health Organization
YLD	years lived with disability

## Acknowledgments

This report was developed by the Healthy Life Course Unit of the Department of Family, Health Promotion and Life Course of the Pan American Health Organization (PAHO). It was drafted by Eugenio Beltran, with important collaboration from Carolina Hommes, Regional Advisor for the Healthy Life Course Integrated Approach.

This publication is part of a series titled “The Decade of Healthy Aging in the Americas: Situation and Challenges”, and is the result of an interagency effort, coordinated and edited by Patricia Morsch, Enrique Vega, and Pablo Villalobos, under the supervision of Luis Andrés de Francisco Serpa, from PAHO.

The purpose of the series is to provide continuous updates on the different areas of action of the Decade of Healthy Aging (2021–2030) in the Americas, as well as on other related aspects.

The collaboration of the experts from PAHO, the United Nations and the Inter-American System, and the academic world who participated in the initiative and formulated essential feedback and recommendations for the project to see the light is appreciated.

## Aging and oral health

At any point in an individual's lifetime, the oral health status represents the cumulative effect of risk indicators, social determinants, personal and familial beliefs, knowledge, health literacy, disease risk factors, and preventive and care behaviors. Thus, oral diseases measured at any point in life represent a snapshot of the individual's life course. Life course has been defined as the study of the long-term effects of risk factors and social exposures from gestation through life, including biological, behavioral, and psychosocial pathways (1). Furthermore, oral diseases, mainly dental caries, periodontal diseases, and their sequelae (tooth loss and edentulism), share common risk factors with other chronic diseases and are preventable (2).

Despite the availability of efficacious and efficient preventive interventions, the risk of oral disease remains throughout the lifetime. Critical life stages with increased risk include early infancy, school age, entering college or university, entering (and leaving) the labor force, and older adult years (3). This publication offers a glance at the oral health status of older adults among countries in the Region of the Americas.

The starting point for this review was an analysis of the literature in PubMed, Medline, and Google Scholar after 2005 using keywords for oral diseases, sequelae, and the names of the countries in the Region. The following sections present the findings for adults aged 60 or 65 years and older on disease occurrence and utilization of dental care services. The evidence table includes the most relevant information on national or regional oral health data from selected studies. Supporting studies not included in the evidence table are included in the discussion of the major findings.

The World Health Organization (WHO) defines healthy aging as “the process of developing and maintaining the functional ability that enables well-being in older age” (4, 5), where functional ability means having the capacity to be and do what people value (5). Functional ability involves the environment, the intrinsic capacity, and the relationship between them. Different health factors, including oral health, might influence specific domains within intrinsic capacity, including mobility, vitality, and psychological, cognitive, and sensory capacities.

Aging also brings changes to the oral tissues and their function (6) such as changes in the quality and amount of saliva (hyposalivation). Dryness of the mouth (xerostomia) is a clinical presentation of Sjögren's



syndrome and post-radiation treatment for cancer; it is also a side effect of therapeutics for systemic conditions in older adults. Both hyposalivation and xerostomia are considered risk factors for dental caries, conditions that can be alleviated using artificial saliva.

With age, the supporting tissues around the tooth (gingiva, periodontal ligament, and alveolar bone) migrate toward the tooth's apex. This condition is known as gingival recession and exposes the tooth root to the intraoral environment. The tooth root is covered by cementum, over which biofilm can increase its susceptibility to dental caries. Regular dental visits and fluorides can keep exposed roots free of dental caries. In these examples, team coordination with other healthcare professionals is essential in delivering effective interventions (7).

Aging is also associated with functional limitations that may affect the ability of individuals to hold a toothbrush and clean their teeth, especially in the interdental areas. These limitations put the individual at a higher risk of dental caries and periodontal diseases. The changes may be incremental with age and may affect the functional abilities of older adults but are also modifiable or controllable. One example is the availability of electric toothbrushes for those with functional inability to hold the toothbrush due to an underlying impairment (8).

Aging is also associated with social determinants that create barriers and limitations in seeking and receiving appropriate dental care. Lack of proper dental care for dental caries and periodontal diseases leads to pain and infection. When dental services are inadequate, both diseases progress toward tooth loss and sometimes to edentulism (missing all teeth). Tooth loss impacts daily living and quality of life, impairing chewing, nutrition, and social interaction with others (3, 9) and demands prostheses and dentures (10). Tooth loss may also affect choice of diet due to an inability to cut and chew. Therefore, the effects of social determinants earlier in life, which increase the risk of disease and affect access to appropriate care, may lead to functional disability to cut and chew and impact nutrition later in life.

Another characteristic affecting the oral health of older adults is their domestic/living circumstances. Many functional older adults live in their own homes (independent living, community dwelling). In contrast, others live in long-term care settings (nursing homes, institutions) due to physical or mental limitations. The oral health of older adults in long-term care facilities is affected by the severity of disease and sequelae prior to moving into the facility and the ability of the long-term care to provide or organize appropriate dental care for the residents.

Furthermore, most residents may have only limited dental coverage, and services need to be paid out of pocket. Therefore, older adults in long-term care have poorer oral health (11), lower utilization of dental services, higher prevalence of tooth loss and untreated tooth decay, increased need for urgent care and prosthetic appliances, and a higher rate of edentulism than independent-living older adults (12–14). A 2021 review suggested that people in long-term care often experience a rapid decline in their oral health after moving into the facility, even if their dentition was stable upon admission (15).

A related and concerning issue among the health of older adults is frailty. A consensus paper defined frailty as a “syndrome with multiple causes and contributors that is characterized by diminished strength, endurance, and reduced physiologic function that increases an individual’s vulnerability for developing increased dependency or death” (16). Frail individuals may not live independently and require assistance to perform everyday activities, including oral hygiene. Oral frailty has been defined by including characteristics such as the number of teeth, chewing inability, loss of motor skills, low tongue pressure, and difficulties in eating and swallowing (17). Many of the studies described later in this chapter exclude frail adults because they were not able to respond to questionnaires or follow instructions during an examination. Thus, information on the oral health of frail older adults is limited. Some studies have focused on dementia and oral health. For example, a systematic review (18) reported higher rates of root caries and gingival bleeding in older adults with dementia, which may be associated with cognitive and behavioral impairment. The reviewers noted that cognitive decline, loss of memory, learning disabilities, attention deficits, deterioration of motor skills, reduction in salivary flow, and difficulties with eating and swallowing were associated with an increase in oral diseases. In a three year cohort study in Mexico, the incidence of frailty among home-dwelling older adults was associated with loss of teeth and severe periodontitis (19).

A further issue is the delivery of dental care. Dental treatment is expensive and many adults lose their dental benefits upon retirement. If they have received complex and costly treatment before retirement, the cost of maintaining and replacing therapies needs to be paid from personal funds—thus, the call for the inclusion of dental care within older adults’ medical benefits (20). A recent systematic review reported that culturally and linguistically diverse older migrants have greater oral health needs and poor oral status and face barriers in accessing dental care in their host countries (21).

These biological, personal, familial, psychosocial, and societal factors exert burden and determine inequalities in oral health within and between countries. They are critical in older populations because of their cumulative effect over the life course. However, many of these effects may be prevented or controlled with appropriate interventions supported by upstream policies and regulations and cost-effective restorative interventions.

## Oral health status

### Country data

Some national and regional oral health surveys in the Americas have included data specifically focused on older adults (Table 1), but data are lacking for most countries. The data provided are comparable because countries have used the decayed, missing, and filled teeth (DMFT) index recommended by the WHO *Oral Health Surveys Basic Methods* (22, 23) and the national health and nutrition examination surveys protocols (24, 25). Table 1 shows the countries for which data are available: Brazil, Canada, Chile, United States of America, Mexico, Puerto Rico, and Uruguay.

**Table 1.** Dental caries, root caries, periodontal status, number of teeth, and edentulism among older adults in selected countries of the Americas

Author, year of publication	Country (data collection)	Age (years)	Sample	No.	Coronal caries prevalence	Caries severity	Root caries	Periodontal status <sup>a</sup>	No. of teeth	Edentulism
Álvarez et al., 2015	Uruguay (2010–2011)	35–44 & 65–74	National survey	769	P <sub>65–74</sub> = 99.6%	DT <sub>65–74</sub> = 0.7 MT <sub>65–74</sub> = 20.9 DMFT <sub>65–74</sub> = 24.1				
Beck et al., 2014	United States (2008–2011)	18–74	Cross-sectional study of Latinos	15 848		DS <sub>65–74</sub> = 0.61 DMS <sub>65–74</sub> = 16.3 MT <sub>65–74</sub> = 9.0	12% (all age groups)			4.1% (all age groups)
Bidinotto et al., 2018	Brazil (2004–2008)	60+	Prospective cohort; four year outcomes	T0 = 388 T4 = 273			ID = 4.9 per 100 root-year. 47% had new lesions			33% became edentulous during follow-up

<sup>a</sup> Includes only periodontal pocket (probing depth) data because it is the most comparable measurement in different methods. Measures are restricted to population estimates, not quadrants, teeth, or periodontal sites. Ppdx denotes the sample (population) prevalence of periodontal depth at age xx. In addition, the values indicating the probing depth used as cut-off point are within parenthesis. For example Ppdx(4+) means the prevalence at age xx years with at least one periodontal pocket of 4 mm or more. For studies reporting the community periodontal index (CPI), we are reporting the proportion giving 4–5 mm or 6+ mm as the worst score in all six sextants.

<b>Centers for Disease Control and Prevention, 2019</b>	United States (2021.)	65+	National survey	2011–2016: 3514 1999–2004: 3539	2011–2016: DFT >0 P <sub>65+</sub> = 96% DT >0 P <sub>65+</sub> = 16% 1999–2004: DFT >0% P <sub>65+</sub> = 93% DT >0 P <sub>65+</sub> = 18%	2011–2016: DT <sub>65+</sub> = 0.3 MT <sub>65+</sub> = 6.6 FT <sub>65+</sub> = 9.8 DFT <sub>65+</sub> = 10 DMFT <sub>65+</sub> = 17 1999–2004: DT <sub>65+</sub> = 0.4 MT <sub>65+</sub> = 8.8 FT <sub>65+</sub> = 8.7 DFT <sub>65+</sub> = 9.1 DMFT <sub>65+</sub> = 18		Mean = 21 (range 16–22)	2011–2016: P <sub>65+</sub> = 17% P <sub>65–74</sub> = 13% P <sub>74+</sub> = 23% 1999–2004: P <sub>65+</sub> = 27% P <sub>65–74</sub> = 24% P <sub>74+</sub> = 31%
<b>Dye et al., 2015</b>	United States (2011–2012)	50–64 65–74 75+	National survey		DFT >0: P <sub>50–64</sub> = 97% P <sub>65–74</sub> = 96% P <sub>75+</sub> = 97% DT >0 P <sub>50–64</sub> = 26% P <sub>65–74</sub> = 19% P <sub>75+</sub> = 19%				P <sub>65–74</sub> = 13% P <sub>75+</sub> = 26% P <sub>NHW</sub> = 17% P <sub>NHB</sub> = 29% P <sub>H</sub> = 15% P <sub>NHA</sub> = 24%
<b>Eke et al., 2016 and 2018<sup>b</sup></b>	United States (2009–2012) (2009–2014)	≥65	National survey	1983, with 1511 dentate				2009–2012: P <sub>pd65–74</sub> (4+) = 51% P <sub>pd65–74</sub> (6+) = 13% P <sub>pd75+(4+)</sub> = 45% P <sub>pd75+(6+)</sub> = 10% 2009–2014: <sup>c</sup> P <sub>pd65+(4+)</sub> = 41% P <sub>pd65+(6+)</sub> = 9%	P <sub>65–74</sub> = 14% P <sub>75+</sub> = 24%
<b>Griffin et al., 2019</b>	United States (1999–2004 and 2011–2016)	65+	Comparing two national surveys and state-based self-reported data (BSS)	1999–2004: 3539 2011–2016: 3514	DT >0 P <sub>65–74</sub> = 15.4% P <sub>≥75</sub> = 16.5%	MT <sub>65–74</sub> = 5.6 MT <sub>≥75</sub> = 7.9 DT <sub>65–74</sub> = 0.3 DT <sub>≥75</sub> = 0.3			P <sub>65–74</sub> = 13% P <sub>≥75</sub> = 23% Variation among states from 25% (Minnesota) to 43% (Arkansas), median 32% <sup>d</sup>

<sup>b</sup> All reports on nationally representative data from the United States using NHANES measure pocket depth and attachment loss on all or a selected number of periodontal sites. Different case definitions have been used and are not included in this review. Two reports of four and six years are included.

<sup>c</sup> The 2018 report including all six years of data did not stratify estimates by age group among older people.

<sup>d</sup> Included only adults aged 65 years and over living in long-term care, assisted living facilities, or home limited.

<b>Lorenzo S et al., 2015</b>	Uruguay (2010–2011)	65–74	First Oral Health Survey (national)	411				P <sub>bleed</sub> = 35% <sup>e</sup> P <sub>3–5mm</sub> = 28.5% P <sub>moderate</sub> = 34% <sup>f</sup> P <sub>severe</sub> = 17%		
<b>Mariño et al., 2015</b>	Chile (2009–2010)	60+	National survey of living older adults. Secondary analysis	4213						20% (self-reported)
<b>Mariño &amp; Giacaman, 2014</b>	Chile (2011.)	65–74	Regional Oral Health Survey, Maule	438		DMFT <sub>65–69</sub> = 24.4 <sup>g</sup> DMFT <sub>70–74</sub> = 26.9 FT <sub>65–74</sub> = 2.0 MT <sub>65–74</sub> = 22.4 DMFT <sub>65–74</sub> = 25.7		CPI <sub>healthy</sub> = 0% CPI <sub>bleed</sub> = 1.5% CPI <sub>calculus</sub> = 68% CPI <sub>4–5mm</sub> = 25.5% CPI <sub>&gt;5mm</sub> = 4.6%	No. of teeth: 1–10 = 34% 11–20 = 25% >20 = 16%	25% both 40% maxilla 34% mandible
<b>Mariño et al., 2015</b>	Chile (2011.) <sup>h</sup>	65–74	Regional Oral Health Survey, Maule	328 dentate 231 with gingival recession		MT <sub>65–74</sub> = 19.1 DRS = 0.21 FRS = 0.55	Root caries index = 8.2%	Exposed root surfaces = 70.4% (gingival recession)		
<b>Mariño et al., 2011</b>	Chile (2006–2007)	60+	Regional study, Valparaíso Region	354		DMFT = 21.6 MT = 17.9 FT = 2.9 DT = 1.9		CPI <sub>healthy</sub> = 3.3% CPI <sub>bleed</sub> = 11.4% CPI <sub>calculus</sub> = 42.3% CPI <sub>4+mm</sub> = 43% (57% rural, 18% non-metro, and 47% urban)	No. of teeth: 1–10 = 34% 11–20 = 33% >20 = 13%	20%; geographical variation from 14% in non-metro centers, 21% in urban municipalities and 29% in rural areas

<sup>e</sup> This study in Uruguay reports the prevalence of having one or more sextants for each periodontal income separately, not by the highest score as originally designed by the community periodontal index and the community periodontal index of treatment needs.

<sup>f</sup> This study in Uruguay used a different classification for moderate/severe periodontal disease and a separate one for severe disease using the combination of periodontal pockets and clinical loss of attachment. In the table, the prevalence of severe is defined as having one or more sextants with an index tooth with pockets  $\geq 4$  mm and at least one sextant with clinical attachment loss of  $\geq 6$  mm. The prevalence of moderate/severe disease expanded the clinical attachment loss to  $\geq 4$  mm

<sup>g</sup> This study in Chile reported over a total of 32 permanent teeth. Most studies in the literature do not include the four third molars, so the maximum would be 28 teeth. Also, some data in the table were obtained from a companion publication (Quinteros et al., 2014 [83]). Ninety-eight percent of the Maule population received fluoridated water in the range of 0.6–1.0 mg/L (only 51% in the rural areas).

<sup>h</sup> Same study as reported in Mariño & Giacaman 2014 (32) but focused on root caries and including only 328 dentate participants. According to Katz (35), the root caries index is the ratio of the sum of decayed and filled root surfaces divided by the sum of all root surfaces, including sound, and multiplied by 100 with one or more caries lesions in the roots. Severity was expressed as the mean number of decayed (DRS) or filled (FRS) tooth surfaces. All measures are calculated among those with gingival recession (n = 231).

<b>Massachusetts Department of Public Health, Office of Oral Health, 2010</b>	United States, 2009	60+ long-term care and meal sites	ASTDD protocol <sup>i</sup>	1046	DT >0 35% in meal sites 59% in long-term care			Gingival inflammation 35% in meal sites, 74% in long-term care.		19% in meal sites, of these, 50% of these had no dentures 35% in long-term care, of these 50% had no dentures
<b>Ministério da Saude, 2004</b>	Brazil (2002–2003)	65–74	National survey	5349	P <sub>65–74</sub> = 99.5%	DMFT <sub>65–74</sub> = 27.8		P <sub>pd65–74(4+)</sub> = 6%	≥20 <sup>j</sup> = 10%	58% had upper total dentures and 16% need 34% had lower total dentures and 24% need
<b>Ministério da Saúde, 2012</b>	Brazil (2010.)	65–74	National survey	7509	P <sub>65–74</sub> = 91.9%	DT = 0.62 MT = 25.29 FT = 1.62 DMFT = 27.53	D <sub>Troot</sub> = 0.23 F <sub>Troot</sub> = 0.10	P <sub>bleed</sub> = 18.1% <sup>k</sup> P <sub>hard</sub> = 28.3% P <sub>3–5mm</sub> = 15.8% P <sub>6+mm</sub> = 3.3%		63% use total upper dentures, 18% need 37.5% use total lower dentures, no data on need
<b>Ministry of Health Canada, 2010<sup>l</sup></b>	Canada (2007–2009)	60–79	Canadian Health Measures Survey	1093	P <sub>DMFT&gt;0</sub> = 100% P <sub>DT&gt;0</sub> = 16%	DMFT = 15.7 MT = 5.6 FT = 9.7 DT = 0.4	P <sub>root</sub> = 43% D <sub>Troot</sub> = 0.3 F <sub>Troot</sub> = 1.3 D <sub>FTroot</sub> = 1.6	P <sub>pd(4+)</sub> = 31% P <sub>pd(6+)</sub> = 78% C <sub>PIhealthy</sub> = 3% C <sub>PIbleed</sub> = 23% C <sub>PIcalculus</sub> = 44% C <sub>PI4–5mm</sub> = 24% C <sub>PI&gt;5mm</sub> = 7%		22%
<b>Montero-Aguilar et al., 2012</b>	Puerto Rico (2007.)	70–97	Cohort from San Juan	183				P <sub>pd70–79(5–6)</sub> = 34% P <sub>pd80+(5–6)</sub> = 20% P <sub>pd70–79(7+)</sub> = 13% P <sub>pd80+(7+)</sub> = 5%	Mean = 15.8	
<b>Phipps et al., 2020</b>	United States (2017.)	55+	State telephone survey							P <sub>65–74</sub> = 5.2% P <sub>75+</sub> = 10%
<b>Sánchez-García et al., 2007</b>	Mexico (2007.)	60+	Social security beneficiaries	698	P = 99.7%	DT = 2.4 MT = 12.2 FT = 2.6 DMFT = 17.3	P <sub>root</sub> = 44% D <sub>Troot</sub> = 1.2 F <sub>Troot</sub> = 0.1 D <sub>FTroot</sub> = 1.4			

<sup>i</sup> The ASTDD protocol is the Basic Screening Survey: An approach to monitoring community oral health, 2008. Available at [www.astdd.org](http://www.astdd.org). The target population was adults who are residents of long-term care facilities (nursing homes) and those who participate in congregate meal sites (subsidized meals in senior centers or clubs).

<sup>j</sup> The WHO goal for 2000 was 50% of the population aged 65–75 years having 20 or more teeth present.

<sup>k</sup> This study reports the prevalence of having one or more sextant for each periodontal level separately, not by the highest score as originally designed by the community periodontal index and the community periodontal index of treatment needs.

<sup>l</sup> This study included participants from a cohort of the Puerto Rican elderly from the San Juan area. It used the NHANES criteria but examined four sites on all teeth.

## Coronal dental caries

Dental caries is a chronic disease generated from an imbalance in the process of demineralization and remineralization of the tooth hard tissues (enamel, dentin, and cementum). Demineralization is caused by organic acids from bacterial metabolism in the biofilm surrounding the tooth. Remineralization is the reverse effect and is led by saliva buffering and presence of fluorides. When the balance is broken, a caries lesion is formed. Initial lesions that are not allowed to remineralize may develop a cavity.

The prevalence of dental caries in older adults (DFT >0 or DMFT >0) was consistently over 90% across all countries. Fewer countries reported the prevalence of untreated decay (DT >0). Canada and the United States reported similar estimates for untreated decay, between 15% and 26%. The most recent figure for the United States for adults aged 65 years and older is 16% (26). However, a 2009 study in Massachusetts reported a prevalence of 35% among older adults at meal sites and 59% among residents of long-term care facilities (13). A systematic review (based on six studies) of dental caries among adults aged 60 years and older by the Centers for Disease Control and Prevention (CDC) Division of Oral Health estimated an incidence rate of one coronal carious surface per person per year (27).

Brazil and the United States had data for at least two data points to assess changes in time. In Brazil, the prevalence of dental caries in older adults aged 65–74 years decreased from 99.5% in 2002–2003 (28) to 91.9% in 2010 (29). In the United States, the prevalence of dental caries increased from 93% in 1999–2004 to 96% in 2011–2016. However, the prevalence of untreated decay decreased from 18% to 16% in the same period (26). These findings reflect small population changes in disease occurrence.

The severity of dental caries was reported by either the mean number of decayed, missing, and filled teeth (DMFT) or excluding the “missing” component (DFT). The difference in reporting relies on the difficulty of determining whether the missing tooth was lost due to dental caries or other reasons (24). The lowest mean DMFT was 10 in Canada in 2007–2009 (30), followed by 17 in the United States in 2011–2016 (26). In Canada, the mean number of missing teeth due to caries was 5.6. In comparison, the mean DMFT in Brazil was 27.5 in 2002–2003 (28) and 27.8 in 2010 (29). A similar estimate (24.1) was reported for Uruguay in 2010–2011 (31). Chile published a similar estimate for Maule (25.7) (32) and a slightly lower one for Valparaíso (21.6) (33). The mean DMFT

in Mexico (17.3) corresponded to a selected sample of social security beneficiaries (34).

The main difference across countries is the differential contribution of the DMFT components. While in Canada and the United States filled was the main contributor to DMFT, missing teeth was the main contributor in the other countries. These represent opposite options in dental treatment for dental caries in both groups.

## Root caries

Data for root caries are sparser and less comparable. One study reported using the root caries index (35). Some studies counted the number of decayed and filled teeth or surfaces while others reported just prevalence. A study among social security beneficiaries in Mexico (34) reported a 44% prevalence and an average of 1.4 decayed and filled roots. Similar estimates were reported in Canada at 43% and DFT was 1.6. Even lower values were reported in Brazil (0.33 teeth). Three additional studies in the literature reported a 12% incidence among Hispanic and Latino populations in the United States (36) and Katz's root caries index of 8.2% in Chile (33). Two studies evaluated root caries incidence. In a prospective study in Brazil, Bidinotto et al. reported an incidence density of 4.9 per 100 root-year, with 47% of the cohort developing new lesions after four years (37). In 2004, CDC reported an annual increment of half a surface from a systematic review of 11 studies in Australia, Canada, Sweden, and the United States (38).

## Periodontal diseases

There is still a lack of agreement on a case definition for the epidemiological evaluation of periodontal diseases and what measurements should be included (39). WHO has made efforts to systematize the epidemiological data by using standard instruments such as the community periodontal index (CPI) or its predecessor, the community periodontal index of treatment needs (CPITN), described in the WHO *Oral Health Surveys Basic Methods* (22, 23). In addition, periodontologists have proposed clinical case definitions to grade the severity of the diseases (40, 41). A common approach is to use a periodontal probe to measure periodontal depths in millimeters (pocket depth) or the migration of the periodontal attachment toward the apex of the tooth (clinical attachment loss). The CPI and CPITN simplified the periodontal measurements into categories of pocket depths, i.e., less than 3.5 mm, between 3.5 and 5.5 mm, between 5.5 and 8.5 mm, and more than 8.5 mm using a periodontal probe with



color bands. (22, 23). Most oral epidemiologists would agree that clinical attachments of 3 mm or less would be considered “normal” or within the margin of error (42). In addition, the original CPITN measured hard deposits around the tooth (calculus) and gingival bleeding on probing (22).

There has been extensive discussion on the validity of measurements based on partial examination of the mouth’s periodontal sites, teeth, or anatomical sections based on quadrants or sextants (39). The most recent edition of the *Oral health surveys basic methods* (23) recommends applying the CPI to all teeth and separating bleeding from periodontal probing. A further recommendation is to supplement the pocket depth information with clinical attachment loss measurements using sextants (22). Self-reported measures of periodontal status have been described (43–45) and validated (46) but have not been used extensively in epidemiological surveys. For detailed information on periodontal assessment at the epidemiological level, see Beltrán-Aguilar et al., 2012 (39).

Mariño et al. published two epidemiological studies in Valparaíso and Maule, Chile, using the CPITN protocol (32, 47). In both studies, calculus had the highest prevalence, 42% and 68%, respectively. Probing depths greater than 3 mm were observed in 43% of participants in Valparaíso and 30% in Maule. A subsequent study by the same team in Maule reported 70% of adults with exposed root surfaces (gingival recession) (33).

The most recent data from Brazil (29) reported the prevalence for each category of the CPITN score, not the highest value of all sextants as recommended by WHO (22). The highest prevalence was for calculus (28%), followed by probing depths greater than 3 mm (19%) and bleeding on probing (18%).

In Uruguay, Lorenzo et al. used the same approach as Brazil (48). They reported 35% of adults aged 65–74 years had bleeding, and 29% had one or more teeth with probing depths of 3 mm or more. In addition, they reported that 17% had severe disease, which they defined as having one or more teeth with periodontal probing of 4 mm or more and one with clinical attachment loss of 6 mm or more.

A 2012 study in Puerto Rico reported estimates for participants aged 70–79 and 80+ years. The prevalence for periodontal depths of 5–6 mm was 34% and 20%, respectively. The prevalence of periodontal depths of 7 mm or more was 13% and 5%, respectively (49).

Health Canada used CPITN in their 2007–2009 Canadian Health Measures Survey (30). They reported a prevalence of 23% for bleeding on probing, 44% for calculus, 24% for periodontal depths of 4–5 mm, and 7% for pocket depths of 6 mm or more.

In the United States, the most recent national data were collected in three cycles of the National Health and Nutrition Examination Survey between 2009 and 2014 and measured six sites per tooth on all teeth (50). Among adults aged 65–74 years, the prevalence of periodontal depths of 4+ mm was 51%, and for depths of 6+ mm was 13%. Among adults aged 75 years and older, the prevalence of depths 4+ mm was 45%, and for 6+ mm was 10%. The decrease in prevalence with age reported in this study and the values reported in Puerto Rico (49) could be explained by the increased number of missing teeth, many of which may have been affected by periodontal disease.

A study of long-term care and meal sites in Massachusetts reported that gingival inflammation was twice as prevalent in long-term care residents than in those living independently (13). The inability to consistently clean the teeth may be associated with higher levels of gingival inflammation among long-term care residents.

Some studies have compared prevalence and severity among countries. Gjeramo et al. (51) identified six studies (most carried out in the 1990s) reporting periodontal status in the Latin American region and the Caribbean using the CPITN index. The prevalence of moderate periodontal disease (one or more teeth) ranged from 29% to 40%, while severe periodontal disease ranged from 7% to 71%. Gjeramo et al. (51) assigned Brazil into the low-level category (<10%), Argentina and El Salvador in the medium level (10%–49%), and Chile in the high category ( $\geq$ 50%). Recent national and regional studies in Brazil (29) and Chile (32, 47) support the earlier estimates, indicating little change in periodontal status in these two countries. A recent review by Kossioni et al. (52) included data from Canada and the United States but not from Latin American countries.

In summary, most countries with available data show some level of periodontal sequelae, calculus, or bleeding, the last being a sign of gingival inflammation. More severe levels were observed in older populations and those living in long-term care, but they are less prevalent than milder forms. The lack of an agreed universal case definition makes comparison across countries difficult. However, across all these studies, older adults may experience a higher prevalence and more severe levels of periodontal diseases than younger adults.

## Tooth loss

If teeth with dental caries or periodontal diseases are not treated, the disease progresses and eventually the tooth has to be extracted due to infection. Indeed, the ultimate goal of dental treatment is to prevent tooth loss and maintain functional ability. The dental practitioner has the final decision for extracting a tooth, and in some healthcare systems, extraction may be the only treatment choice. Keeping the natural teeth is also a cultural and generational issue. Baby boomers and later generations expect to keep their natural teeth for as long as possible. The number of natural teeth has been used to indicate inequalities among older adults (53). Another cause for tooth loss is dental injury, but this tends to occur earlier in life.

Missing teeth are replaced using prosthetics and, more recently, implants. Contrary to what was once a standard procedure, not all missing teeth need to be replaced for functional reasons (54). This finding led to the development of the “shortened arch length” concept; that is, most people can chew with a complete set of anterior teeth and premolars (55, 56). The shortened arch length concept led to the international goal of having at least 20 teeth (57) and the Japanese 80/20 goal (20 teeth at age 80) .

The studies in Maule (32) and Valparaíso (47) provide a detailed explanation of tooth retention in Chile. Both populations had similar distributions: 34% having 1-10 teeth; 25% (Maule) and 33% (Valparaíso) having 11-20 teeth, and 16% (Maule) and 13% (Valparaíso) having 20 or more teeth. The researchers concluded that Chile was experiencing an epidemiological transition, with higher rates of tooth retention, which may require more dental care services.

Ten percent of older adults in Brazil had 20 or more teeth (28). However, in that study, remnants and root tips were included in counting the number of teeth, and these would be eventually extracted (52). In a 2007 study in Puerto Rico, the mean number of teeth was 15.8 (49), while the most recent NHANES survey in the United States reported a mean of 21 teeth. In the United States, there was an increase of two teeth on average from 1999-2004 (26). These data suggest variability in the number of teeth present in selected countries. Still, the epidemiological direction—probably an effect of preventive and curative efforts—is toward retaining more natural teeth in the younger generations. Evidence from Australia and Norway supports retaining 20 or more permanent teeth as a workable target for public health interventions (58, 59).

## Edentulism

The clinical decision to extract all remaining teeth is complex and based on the individual needs and available dental services. Some people who retain a handful of teeth seek extraction of the remaining teeth to replace all missing teeth with dentures. Others could be edentulous only in one dental arch for prolonged periods. A common clinical option is to retain some of the roots after endodontic treatment or insert dental implants to increase the retention of dentures and reduce the loss in alveolar bone height with aging. These are known as overdentures. Edentulism can be measured using clinical assessments (26) or by asking study participants whether they have lost all their teeth (60).

Mariño et al. reported that in 2009–2010, 20% of Chileans identified themselves as having no teeth (61). Clinical surveys by the same team reported normative estimates of 20% in Valparaíso in 2006–2007 (47) and 25% in Maule (32). In the former, the investigators reported geographical variations ranging from 14% to 25%.

In Canada, the prevalence in 2007–2009 was 22% among people aged 60–79 years (30). In the United States in 2011–2016, the prevalence was 17% among those aged 65 years and older (26); this represented a 10% point reduction in edentulism since 1999–2004. Reduction in edentulism is a trend that has been reported in the United States (62) and other countries (63). CDC also reported variations in edentulism by race/ethnicity, with the highest prevalence among non-Hispanic blacks (29%) and non-Hispanic Asians (24%) and the lowest among Hispanics (15%) (26, 64). The lower prevalence among Hispanics, primarily of Mexican descent living in the United States, has been described as an epidemiologic paradox (65). In the 1982–1984 Hispanic HANES, 2.6% among Mexican Americans and 6.1% among Cuban Americans were edentulous (66). Researchers at CDC using self-reported data have reported variations at the state level, from 25% in Minnesota to 43% in Arkansas (67). In Massachusetts, the prevalence of edentulism was 19% among adults aged 60 years or older attending meal sites and 35% among adults living in long-term facilities (13). In Washington State, the self-reported prevalence was 5% among adults aged 65–74 years, and 20% among those aged 75 years and older (68). A cohort study of Latinos in the United States reported a prevalence of edentulism of 4% in 2008–2011, but this included all age groups (36).

A Brazilian study reported that 33% of a cohort examined after four years became edentulous over the 2004–2008 period (37). The two most recent national studies in Brazil reported on the use of and need

for dentures: in 2010, 63% of adults aged 65–74 used upper dentures and 16% needed dentures, while 24% had lower dentures and 24% needed dentures. These estimates represent a higher prevalence of edentulism.

Overall, most countries show estimates of edentulism around 20%, and a few reported a historical reduction in prevalence (secular trends). Other countries outside the Region have reported similar results (14). There is also evidence of inequalities in tooth loss and edentulism between and within countries. For example, living in rural areas (69) or families with a poor income (25) are associated with a higher prevalence of edentulism. However, in Valparaíso both rural and metropolitan areas had higher rates of edentulism (47).

Edentulism has been associated with systemic conditions. A CDC study reported that the rate of complete or partial tooth loss was greater among older adults with arthritis, cardiovascular disease, chronic obstructive pulmonary disease, diabetes, and low vision/blindness than among those without these conditions (12). Edentulism has also been associated with loss of body weight and waist circumference in community-dwelling Brazilian adults aged 60 years and older (70). In 2013, edentulism was the 24th ranked leading cause of years lived with disability (YLDs) according to the Global Burden of Disease (71). Edentulism is also included in the Global Burden of Disease assessment of disability-adjusted life years (DALYs). In the most recent update (72), a 64% increase in DALYS was estimated for oral diseases, including tooth loss, worldwide.

## Utilization of dental services

An essential component of oral health is access to dental services for prevention, control, treatment, and evaluation. Access to dental care goes beyond urgent treatment needs or routine dental treatment. A more comprehensive concept is “dental home,” defined as having access to routine, comprehensive, continuous, coordinated, and affordable dental services (73). Having a dental home implies not having barriers to dental care such as availability, proximity, traveling time, identification with the healthcare providers, and paying for the services received. In the United States, the extension of dental homes to older adults has started with calls for the modification and expansion of entitlements such as Medicare to reduce the financial barriers to health (20).

At the health service research level, having a dental visit is the global standard to measure the utilization of health services. It is generally accepted that people should have at least one dental visit per year for a check-up and tooth cleaning, or more often based on the person's needs. These time frames are based on clinical experience rather than on scientific evidence. Indeed, a recent multicenter randomized trial in the United Kingdom showed no difference in dental outcomes between six-month and 24-month recalls (74). These findings may suggest that having limited or alternative dental procedures during long periods of not receiving dental care would be appropriate, for example, during pandemic restrictions (75, 76). This section will provide information regarding the utilization and cost of dental services.

Education (77) and paying for services (78) are significant barriers to dental services. In Brazil, having more education and having higher levels of wealth were directly associated with greater utilization of dental services (79). In general, older adults living in rural areas face greater barriers to accessing dental services than those living in urban areas (80).

The earliest oral health evaluation among older adults in Latin America (81) reported 25% utilization rates across the region. Two decades later, a study in Peru (2015–2017) among adults 60 years and older reported a lower rate (18.5%) but within a six-month time frame (82). In the Peruvian study, a higher level of education was associated with dental visits.

In Brazil, the 2003–2006 national survey reported 6% of adults aged 65–74 had never had a dental visit in their life. Among those who had had a dental visit, 17% had a visit with the previous year. Pain was the main complaint in 48% of those having a dental visit (28). The proportion who had never had a dental visit increased to 15% in a similar national survey in 2012 (29).

A study of social security beneficiaries in Mexico City (34) reported 54% receiving dental health services in the previous 12 months. Of these, 81% sought private dental services, 13% social security services, and 6% public services. Being a woman, having six or fewer years of schooling, having dental caries or sequelae, and having 22 or more teeth were associated with more dental visits among beneficiaries. These may represent people with a higher need for dental care. The main reasons for a dental visit were dental examination (44%), dental restoration (35%), dental cleaning (32%), and dental extractions (23%).

Two studies from Chile reported on the consequences of not having regular dental visits. In the study by Mariño et al. (47) on older adults from the Valparaíso region, 44% of their treatment needs were unmet. Of the 282 participants (20%) who were fully edentulous, 40 (56%) needed both dentures. The Chilean study in Maule reported 41% of restorative needs as unmet (32). Of the 110 (25%) participants who were edentulous, only 14% had both appliances in good condition, and 72 (65%) needed both dentures. These two studies used comparable methods and highlighted regional similarities and differences. They also reported differences in the treatment received within administrative regions. In the Maule study, people in urban areas had more fillings than those living in rural areas, while there were more extracted teeth in the rural areas (83).

In the United States, 45% of adults aged 65 years and older reported a dental visit during 2006 (84) using data from the Medical Expenditure Survey. The most recent data among older adults in the United States (2016) reported by the American Dental Association Health Policy Institute (85) showed no change (43%).

Regarding payment of dental services, data from the Medical Expenditure Survey showed that 38% of adults aged 64 years and older were receiving dental coverage in 2015, an increase of less than 4% since 1996. This means that almost two-thirds of participants did not have dental coverage, public or private (86). Most participants who did hold dental coverage had personal coverage (24% including out-of-pocket expenditure). Among those receiving dental services, 29% were preventive, 42% diagnostic, 7% restorative, 14% prosthetic, and 4% surgical. From the same panel data, mean dental expenditures were lower among persons aged 75 years and older than among those aged 55 to 74 years, suggesting that they may have selected no treatment or lower-cost options such as tooth extraction (12).

Across all countries, at least 50% of older adults did not have a dental visit in the previous year, and in Brazil, 15% had never had a dental visit. Payment for services appears to be an essential factor in having a dental visit and the type of treatment received. For example, social security beneficiaries in Mexico had a wider variety of services than countries where fee-for-services or out-of-pocket payments are required. In the latter, a dental extraction may be the treatment of choice due to cost. Urban versus rural differences were a factor in Chile. Furthermore, a large proportion of edentulous adults require new dentures. In summary, these data suggest that older adults in the Americas do not have adequate access to dental care.

## Discussion

The oral health status of older adults represents a life course event, where social determinants, societal expectations, and government interventions play a role and change throughout the lifetime. Oral diseases impact the functional abilities of older adults and their quality of life, and interfere with their healthy aging. In addition, oral diseases burden families, healthcare systems, communities, and societies and cause inequalities (12). Therefore, immediate actions and long-term upstream changes are required to relieve the current disease burden, improve functional abilities, and reduce inequalities.

This report has focused on the most common oral diseases, their sequelae, and the utilization of dental services. Maintaining a functional masticatory system is an essential aspect of healthy aging. Therefore, oral diseases, tooth loss, and inadequate utilization of dental services preclude healthy aging. Other conditions afflicting older adults but not included in the report are oral and pharyngeal cancers and physical and functional changes consequent to systemic diseases.

Access to care and how older adults seek, receive, and pay for dental services appear to be essential elements in improving the oral health of older adults. Older adults with greater dental needs but not receiving appropriate care exacerbate personal, familial, and societal burdens. Expansion of public health coverage for dental needs and other safety nets such as Medicare (87) in the United States may provide temporary relief. Still, system changes such as universal health coverage are needed for long-term effects. In Chile, the Explicit Health Guarantees (Garantías Explícitas de Salud, GES) guarantee citizens aged 60+ years access to high-quality care for 56 health conditions, including oral health. Also, some countries have integrated dental care into primary health care delivery, but there is a large backlog of accumulated needs to be addressed (88).

The 2021 World Health Assembly resolution ([WHA74.5](#)) about oral health, provides a pivotal starting point for such changes (89). Existing WHO policies for aging populations (5, 90) may guide the development of interventions for older adults. Other aspects include the determination of essential dental procedures (91), better dental practices for older adults (34), a set of global epidemiological indicators (92), and developing medical-dental integration approaches to maintain their oral health and functional abilities.



Some national and regional organizations have taken the lead on improving the oral health of older adults; for example, the Excellence Research Program on Healthy Aging (PIEI-ES) at the University of Talca, Chile (<https://www.otalca.cl/>), and the Michigan Coalition for Oral Health for the Aging (<https://www.micoha.org/>). Government agencies in the United States like CDC (93) and the National Institute of Dental and Craniofacial Research (94) have prepared materials for the public. Professional organizations such as FDI World Dental Federation have also published information for the public (95) and guidelines for dental practitioners (96).

## Gaps in knowledge and methods

This review found a large volume of information on the oral health of older adults, but with gaps in the knowledge and limitations in the methods. For example, there are limited or no data on frail older adults, those living in long-term care, the Caribbean, and Indigenous populations outside Canada and the United States. In terms of surveillance, there was inconsistent reporting by age group despite WHO recommendations.

Most of the data summarized here used clinical or self-reported approaches with a wide range of representation, including local, regional, and national levels. However, except for Canada and the United States, no national objectives delineate the type of data needed to monitor the burden of oral diseases among older adults. Many of the studies in South America followed the WHO *Oral Health Surveys Basic*, which facilitated the comparison between countries. However, there were inconsistencies in applying case definitions, especially for periodontal status.

Regarding newer approaches for disease monitoring, in the United States, the Association of State and Territorial Dental Directors has developed a protocol to complete oral screenings on adults and older people in community-dwelling or long-term care facilities (97). In 2016, FDI World Dental Federation and WHO convened a panel of experts to discuss the oral health challenge of the aging population. One of the topics was identifying practical surveillance approaches for older adults (98). Still, these rely on the traditional approach of measuring disease, not function or health. Thus, new methods are needed to evaluate and monitor oral health and functional abilities among older adults and their progress toward healthy aging.

Tied to these limitations is the need for integration in the reporting, handling, and evaluating of existing and upcoming data. A few years ago, the leading depository of oral health data was the WHO Oral Health Data Bank at Malmö University. The PAHO Oral Health Program made quadrennial reports on oral health status. Both systems provide limited oral data on adults. There is a dire need for integrated system approaches to the oral health of older adults worldwide.

Another missing aspect of oral health for older adults is the skills and competencies of the dental provider. Graduates from dental schools are not yet ready to meet current and future general and oral health needs of older adults despite some positive integrating of competencies into the curriculum in the United States and Europe (52).

## Conclusions

The evidence presented in this report calls for increasing actions to achieve healthy aging in the Region of the Americas by developing policies and interventions, including integrated delivery of dental care with medical and social care. Two frameworks for such initiatives are the 2021 WHO Resolution on Oral Health (99) and the Decade of Healthy Ageing 2020–2030 (100). These efforts should consider the heterogeneity of the older adult population within each country, region, and community, especially those most vulnerable.

A 2012 evaluation of the oral health of older adults in the United States (12) recommended the following actions:

- better integration of oral health care into medical care;
- implementing community programs to promote healthy behaviors and improve access to preventive services;
- developing comprehensive strategies to address the oral health needs of the homebound and long-term care residents;
- assessing the feasibility of a safety net covering all aspects of dental care to eliminate pain and infection.

In terms of care, specialized professionals in geriatric dentistry have proposed a conceptual framework (7) which includes:

- inserting minimal intervention dental procedures in the care of older adults;
- a focus on maintaining oral functionality instead of “treating all”—includes social care, preserving functional ability, and improving quality of life;
- patient-centered care: acknowledging the heterogeneity of needs instead of protocolized treatment;
- interdisciplinary teamwork as a critical aspect of care.

These recommendations underlie the following needs:

- having trained healthcare professionals, including primary care providers, delivering preventive and restorative care;
- establishing dental homes for older adults at community centers and long-term care facilities;
- integrating the economic evaluation of interventions;
- monitoring indicators as new generations move through their life-span.

Responsibility for these recommendations falls on local and national agencies, academic and research institutions, nongovernmental organizations, and international donors, a process that international agencies can facilitate.

## References

1. Ben-Shlomo Y, Kuh D. A life course approach to chronic disease epidemiology: conceptual models, empirical challenges and interdisciplinary perspectives. *Int J Epidemiol.* 2002;31(2):285–293. Available from: <https://pubmed.ncbi.nlm.nih.gov/11980781/>.
2. Sheiham A. Oral health, general health and quality of life. *Bull World Health Organ.* 2005 Sep;83(9):644. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2626333/>.
3. Department of Health and Human Services. Oral health in America: a report of the Surgeon General. Rockville, MD: US Department of Health and Human Services. National Institute of Dental and Craniofacial Research; 2000.
4. Beard JR, Officer A, de Carvalho IA, Sadana R, Pot AM, Michel J-P, et al. The world report on ageing and health: a policy framework for healthy ageing. *Lancet.* 2016;387(10033):2145–54. Available from: [https://doi.org/10.1016/S0140-6736\(15\)00516-4](https://doi.org/10.1016/S0140-6736(15)00516-4).
5. World Health Organization. World report on ageing and health. Geneva: WHO; 2015.
6. Ribeiro de Castro MC, Ramos-e-Silva M. Physiologic alterations of oral mucosa due to aging: a review. *Acta Dermatovenerol Croat.* 2005;13(3):184–8. PMID: 16146622.
7. León S, Giacaman RA. Proposal for a conceptual framework for the development of geriatric dentistry. *J Dent Res.* 2022 Mar;101(3):247–252. Available from: <https://doi.org/10.1177/00220345211042268>.
8. Yaacob M, Worthington HV, Deacon SA, Deery C, Walmsley AD, Robinson PG, et al. Powered versus manual toothbrushing for oral health. *Cochrane Database Syst Rev.* 2014;2014(6):CD002281-CD. Available from: <https://doi.org/10.1002/14651858.CD002281.pub3>.
9. Sarita PT, Witter DJ, Kreulen C, M., Vant't Hof MA, Creugers NH. Chewing abilities of subjects with shortened dental arches. *Community Dent Oral Epidemiol.* 2003 Oct;31(5):328–34. Available from: <https://doi.org/10.1034/j.1600-0528.2003.t01-1-00011.x>.
10. Boscato N, Schuch HS, Grasel CE, Goettens ML. Differences of oral health conditions between adults and older adults: a census in a Southern Brazilian city. *Geriatr Gerontol Int.* 2016 Sep;16(9):1014–20. Available from: <https://doi.org/10.1111/ggi.12588>.

11. Strayer MS. Dental health among homebound elderly. *J Public Health Dent.* 1993 Winter;53(1):12-6. Available from: <https://doi.org/10.1111/j.1752-7325.1993.tb02665.x>.
12. Griffin SO, Jones JA, Brunson D, Griffin PM, Bailey WD. Burden of oral disease among older adults and implications for public health priorities. *Am J Public Health.* 2012 Mar;102(3):411-8. Available from: <https://doi.org/10.2105/AJPH.2011.300362>.
13. Massachusetts Department of Public Health, Office of Oral Health. The Commonwealth's high-risk senior population results and recommendations from 2009 statewide oral health assessment. Boston, MA: Massachusetts Department of Public Health; 2010.
14. Karki A, Monaghan N, Morgan M. Oral health status of older people living in care homes in Wales. *Br Dent J.* 2015 Oct 9;219(7):331-4. Available from: <https://doi.org/10.1038/sj.bdj.2015.756>.
15. Patel J, Wallace J, Doshi M, Gadanya M, Ben Yahya I, Roseman J, et al. Oral health for healthy ageing. *Lancet Healthy Longev.* 2021 Aug;2(8):e521-e527. Available from: [https://doi.org/10.1016/S2666-7568\(21\)00142-2](https://doi.org/10.1016/S2666-7568(21)00142-2).
16. Morley JE, Vellas B, van Kan GA, Anker SD, Bauer JM, Bernabei R, et al. Frailty consensus: a call to action. *J Am Med Dir Assoc.* 2013;14(6):392-7. Available from: <https://doi.org/10.1016/j.jamda.2013.03.022>.
17. Tanaka T, Takahashi K, Hirano H, Kikutani T, Watanabe Y, Ohara Y, et al. Oral frailty as a risk factor for physical frailty and mortality in community-dwelling elderly. *J Gerontol A Biol Sci Med Sci.* 2018 Nov 10;73(12):1661-1667. Available from: <https://doi.org/10.1093/gerona/glx225>.
18. Lauritano D, Moreo G, Della Vella F, Di Stasio D, Carinci F, Lucchese A, et al. Oral health status and need for oral care in an aging population: a systematic review. *Int J Environ Res Public Health.* 2019;16(22):4558. Available from: <https://doi.org/10.3390/ijerph16224558>.
19. Castrejón-Pérez RC, Jiménez-Corona A, Bernabé E, Villa-Romero AR, Arrive E, Dartigues JF, et al. Oral disease and 3-year incidence of frailty in Mexican older adults. *J Gerontol A Biol Sci Med Sci.* 2017 Jul 1;72(7):951-957. Available from: <https://doi.org/10.1093/gerona/glw201>.
20. Chávez EM, Calvo JM, Jones JA. Dental homes for older Americans: The Santa Fe group call for removal of the dental exclusion in Medicare. *Am J Public Health.* 2017 May;107(S1):S41-S43. Available from: <https://doi.org/10.2105/AJPH.2017.303864>.

21. Hoang H, Feike S, Lynden T, Barnett T, Crocombe L. Oral health needs of older migrants with culturally and linguistically diverse backgrounds in developed countries: A systematic review. *Australas J Ageing*. 2020 Sep;39(3):193–208. Available from: <https://doi.org/10.1111/ajag.12759>.
22. World Health Organization. Oral health surveys: basic methods. 4th edition. Geneva: WHO; 1997. Available from: <https://apps.who.int/iris/handle/10665/41905>.
23. World Health Organization. Oral health surveys: basic methods. 5th edition. Geneva: WHO; 2013. Available from: <https://www.who.int/publications/i/item/9789241548649>.
24. Beltrán-Aguilar ED, Barker LK, Canto MT, Dye BA, Gooch BF, Griffin SO, et al. Surveillance for dental caries, dental sealants, tooth retention, edentulism, and enamel fluorosis —United States, 1988–1994 and 1999–2002. *MMWR Surveill Summ*. 2005 Aug 26;54(3):1–43. PMID: 16121123.
25. Dye BA, Tan S, Smith SM, Lewis B, Barker LK, Thornton-Evans G, et al. Trends in oral health status: United States, 1988–1994 and 1999–2004. *Vital Health Stat*. 2007;11(248):1–92. PMID: 17633507.
26. Centers for Disease Control and Prevention. Trends in dental caries and sealants, tooth retention, and edentulism, United States 1999–2004 and 2011–2016. Atlanta: CDC; 2019. Available from: <https://www.cdc.gov/oralhealth/publications/OHSR-2019-index.html>.
27. Griffin SO, Griffin PM, Swann JL, Zlobin N. New coronal caries in older adults: implications for prevention. *J Dent Res*. 2005;84(8):715–20. Available from: <https://doi.org/10.1177/154405910508400806>.
28. Ministério da Saúde Brasil. Projeto SB Brasil 2003: condições de saúde bucal da população brasileira 2002–2003: resultados principais. Brasília: Ministério da Saúde, Secretaria de Atenção à Saúde DdAB; 2004. Available from: [https://bvsmms.saude.gov.br/bvs/publicacoes/projeto\\_sb2004.pdf](https://bvsmms.saude.gov.br/bvs/publicacoes/projeto_sb2004.pdf).
29. Ministério da Saúde Brasil. SB Brasil 2010. Pesquisa Nacional de Saúde Bucal. Resultados principais. Brasília: Ministério da Saúde; 2012. Available from: [https://bvsmms.saude.gov.br/bvs/publicacoes/pesquisa\\_nacional\\_saude\\_bucal.pdf](https://bvsmms.saude.gov.br/bvs/publicacoes/pesquisa_nacional_saude_bucal.pdf).
30. Health Canada. Report of the findings of the oral health component of the Canadian Health Measures Survey 2007–2009. Ottawa: Ministry of Health Canada; 2010. Available from: <https://publications.gc.ca/site/eng/369649/publication.html>.

31. Álvarez L, Liberman J, Abreu S, Mangarelli C, Correa MB, Demarco FF, et al. Dental caries in Uruguayan adults and elders: findings from the first Uruguayan National Oral Health Survey. *Cad Saude Publica*. 2015 Aug;31(8):1663–72. Available from: <https://doi.org/10.1590/0102-311X00132214>.
32. Mariño R, Giacaman RA. Factors related to unmet oral health needs in older adults living in Chile. *Arch Gerontol Geriatr*. 2014 May–Jun;58(3):454–9. Available from: <https://doi.org/10.1016/j.archger.2014.01.003>.
33. Mariño RJ, Fu CS, Giacaman RA. Prevalence of root caries among ambulant older adults living in central Chile. *Gerodontology*. 2015;32(2):107–14. Available from: <https://doi.org/10.1111/ger.12060>.
34. Sánchez-García S, de la Fuente-Hernández J, Juárez-Cedillo T, Mendoza JM, Reyes-Morales H, Solórzano-Santos F, et al. Oral health service utilization by elderly beneficiaries of the Mexican Institute of Social Security in Mexico City. *BMC Health Serv Res*. 2007;7(211). Available from: <https://doi.org/10.1186/1472-6963-7-211>.
35. Katz R. Assessing root caries in populations: the evolution of root caries index. *J Public Health Dent*. 1980 Winter;40(1):7–16. Available from: <https://doi.org/10.1111/j.1752-7325.1980.tb01844.x>.
36. Beck JD, Youngblood M Jr, Atkinson JC, Mauriello S, Kaste LM, Badner VM, et al. The prevalence of caries and tooth loss among participants in the Hispanic Community Health Study/Study of Latinos. *J Am Dent Assoc*. 2014 Jun;145(6):531–40. Available from: <https://doi.org/10.14219/jada.2014.25>.
37. Bidinotto AB, Martins AB, Dos Santos CM, Hugo FN, Hilgert JB, Celeste RK, et al. Four-year incidence rate and predictors of root caries among community-dwelling south Brazilian older adults. *Community Dent Oral Epidemiol*. 2018 Apr;46(2):125–131. Available from: <https://doi.org/10.1111/cdoe.12343>.
38. Griffin SO, Griffin PM, Swann JL, Zlobin N. Estimating rates of new root caries in older adults. *J Dent Res*. 2004 Aug;83(8):634–8. Available from: <https://doi.org/10.1177/154405910408300810>.
39. Beltrán-Aguilar ED, Eke PI, Thornton-Evans G, Petersen PE. Recording and surveillance systems for periodontal diseases. *Periodontol 2000*. 2012;60(1):40–53. Available from: <https://doi.org/10.1111/j.1600-0757.2012.00446.x>.
40. Eke PI, Page RC, Wei L, Thornton-Evans G, Genco RJ. Update of the case definitions for population-based surveillance of periodontitis. *J Periodontol*. 2012;83(12):1449–54. Available from: <https://doi.org/10.1902/jop.2012.110664>.

41. Tonetti MS, Greenwell H, Kornman KS. Staging and grading of periodontitis: Framework and proposal of a new classification and case definition. *J Periodontol*. 2018 Jun;89 Suppl. 1:S159–72. Available from: <https://doi.org/10.1002/JPER.18-0006>.
42. Burt BA, Eklund SA. Periodontal diseases. In: Burt BA, Eklund SA, editors. *Dentistry, dental practice, and the community*, 5th ed. Philadelphia: WB Saunders Company; 1999. p. 237–58.
43. Joshipura KJ, Douglass CW, Garcia RI, Valachovic R, Willett WC. Validity of a self-reported periodontal disease measure. *J Public Health Dent*. 1996;56(4):205–12. Available from: <https://doi.org/10.1111/j.1752-7325.1996.tb02437.x>.
44. Joshipura KJ, Pitiphat W, Douglass CW. Validation of self-reported periodontal measures among health professionals. *J Public Health Dent*. 2002 Spring;62(2):115–21. Available from: <https://doi.org/10.1111/j.1752-7325.2002.tb03431.x>.
45. Blicher B, Joshipura K, Eke P. Validation of self-reported periodontal disease: a systematic review. *J Dent Res*. 2005 Oct;84(10):881–90. Available from: <https://doi.org/10.1177/154405910508401003>.
46. Eke PI, Dye BA, Wei L, Slade GD, Thornton-Evans GO, Beck JD, et al. Self-reported measures for surveillance of periodontitis. *J Dent Res*. 2013 Nov;92(11):1041–7. Available from: <https://doi.org/10.1177/0022034513505621>.
47. Mariño RJ, Cueto A, Badenier O, Acevedo R, Moya R. Oral health status and inequalities among ambulant older adults living in central Chile. *Community Dent Health*. 2011;28(2):143–8. PMID: 21780353.
48. Lorenzo SM, Alvarez R, Andrade E, Piccardo V, Francia A, Massa F, et al. Periodontal conditions and associated factors among adults and the elderly: findings from the first National Oral Health Survey in Uruguay. *Cad Saude Publica*. 2015;31:2425–36. Available from: <https://doi.org/10.1590/0102-311X00012115>.
49. Montero-Aguilar M, Muñoz-Torres F, Elías-Boneta AR, Dye B, Joshipura KJ. High levels of periodontal disease among the older adult population in San Juan, Puerto Rico. *Community Dent Health*. 2012 Sep;29(3):224–8. PMID: 23038939.
50. Eke PI, Thornton-Evans GO, Wei L, Borgnakke WS, Dye BA, Genco RJ. Periodontitis in US Adults: National Health and Nutrition Examination Survey 2009–2014. *J Am Dent Assoc*. 2018 Jul;149(7):576–88.e6. Available from: <https://doi.org/10.1016/j.adaj.2018.04.023>.
51. Gjerme P, Rösing G, Susin C, Oppermann R. Periodontal diseases in Central and South America. *Periodontol 2000*. 2002;29:70–8. Available from: <https://doi.org/10.1034/j.1600-0757.2002.290104.x>.



52. Kossioni AE. Current status and trends in oral health in community-dwelling older adults: a global perspective. *Oral Health Prev Dent*. 2013;11(4):331-40. Available from: <https://doi.org/10.3290/j.ohpd.a30171>.
53. Haugejorden O, Klock KS, Astrom AN, Skaret E, Trovik TA. Socio-economic inequality in the self-reported number of natural teeth among Norwegian adults an analytical study. *Community Dent Oral Epidemiol*. 2008;36:269-78. Available from: <https://doi.org/10.1111/j.1600-0528.2007.00367.x>.
54. World Health Organization. Recent advances in oral health. Report of a WHO Expert Committee. Geneva: WHO; (Tech Rep Ser.) 1992;826:1-37. PMID: 1462607.
55. Kayser AF. Shortened dental arches and oral function. *J Oral Rehabil*. 1981;8:457-62. Available from: <https://doi.org/10.1111/j.1365-2842.1981.tb00519.x>.
56. Witter DJ, van Palenstein Helderma WH, Creugers NHJ, Käyser AF. The shortened dental arch concept and its implications for oral health care. *Community Dent Oral Epidemiol*. 1999;27(4):249-58. Available from: <https://doi.org/10.1111/j.1600-0528.1999.tb02018.x>.
57. Fédération Dentaire Internationale. Global goals for oral health for the year 2000. *Int Dent J*. 1982 Mar;32(1):74-7. PMID: 6951815.
58. Singh KA, Brennan DS. Chewing disability in older adults attributable to tooth loss and other oral conditions. *Gerodontology*. 2012;29(2):106-10. Available from: <https://doi.org/10.1111/j.1741-2358.2010.00412.x>.
59. Hadler-Olsen E, Jönsson B. Oral health and use of dental services in different stages of adulthood in Norway: a cross sectional study. *BMC Oral Health*. 2021;21(257). Available from: <https://doi.org/10.1186/s12903-021-01626-9>.
60. Saman DM, Lemieux A, Arevalo O, Lutfiyya MN. A population-based study of edentulism in the US: does depression and rural residency matter after controlling for potential confounders? *BMC Public Health*. 2014;14:65. Available from: <https://doi.org/10.1186/1471-2458-14-65>.
61. Mariño R, Albala C, Sanchez H, Cea X, Fuentes A. Self-assessed oral-health status and quality of life of older Chilean. *Arch Gerontol Geriatr*. 2013 May-Jun;56(3):513-7. Available from: <https://doi.org/10.1016/j.archger.2012.12.004>.
62. Slade GD, Akinkugbe AA, Sanders AE. Projections of U.S. edentulism prevalence following 5 decades of decline. *J Dent Res*. 2014 Oct;93(10):959-65. Available from: <https://doi.org/10.1177/0022034514546165>.

63. Australian Research Centre for Population Oral Health. National study of adult oral health 2017–18. Adelaide: The University of Adelaide; 2019. Available from: <https://www.adelaide.edu.au/arc poh/national-study/>.
64. Wu B, Liang J, Plassman BL, Remle RC, Bai L. Oral health among white, black, and Mexican–American elders: an examination of edentulism and dental caries. *J Public Health Dent*. 2011 Fall;71(4):308–17. Available from: <https://doi.org/10.1111/j.1752-7325.2011.00273.x>.
65. Spolsky VW, Marcus M, Der-Martirosian C, Coulter ID, Maida CA. Oral health status and the epidemiologic paradox within latino immigrant groups. *BMC Oral Health*. 2012;12(39). Available from: <https://doi.org/10.1186/1472-6831-12-39>.
66. Ismail AI, Szpunar SM. The prevalence of total tooth loss, dental caries, and periodontal disease among Mexican Americans, Cuban Americans, and Puerto Ricans: findings from HHANES 1982–1984. *Am J Public Health*. 1990;80 Suppl:66–70. Available from: <https://doi.org/10.2105/ajph.80.suppl.66>.
67. Griffin SO, Griffin PM, Li C-H, Bailey WD, Brunson D, Jones JA. Changes in older adults’ oral health and disparities: 1999 to 2004 and 2011 to 2016. *J Am Geriatr Soc*. 2019 Jun;67(6):1152–1157. Available from: <https://doi.org/10.1111/jgs.15777>.
68. Phipps KR, Katabi AF, Riter D, Lewis K. Oral health of older adults in Washington State. *J Dent Hyg*. 2020 Feb;94(1):39–48. PMID: 32127428.
69. Slade GD, Spencer AJ, Roberts-Thomson KF. Australia’s dental generations. The National Survey of Adult Oral Health 2004–06. Canberra: Australian Institute of Health and Welfare; 2007. Available from: <https://www.adelaide.edu.au/arc poh/downloads/publications/reports/dental-statistics-research-series/nsaoh-report.pdf>.
70. Andrade FB de, Lebrao ML, Duarte YAdO, Santos JLF. Oral health and changes in weight and waist circumference among community-dwelling older adults in Brazil. *J Am Dent Assoc*. 2014;147:731–6. Available from: <https://doi.org/10.14219/jada.2014.35>.
71. Global Burden of Disease Study 2013 Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2015;386(9995):743–800. Available from: [https://doi.org/10.1016/S0140-6736\(15\)60692-4](https://doi.org/10.1016/S0140-6736(15)60692-4).
72. Kassebaum NJ, Smith AGC, Bernabé E, Fleming TD, Reynolds AE, Vos T, et al. Global, regional, and national prevalence, incidence, and disability-adjusted life years for oral conditions for 195 countries, 1990–2015: a systematic analysis for the Global Burden of Diseases, injuries, and risk factors. *J Dent Res*. 2017 Apr;96(4):380–387. Available from: <https://doi.org/10.1177/0022034517693566>.

73. American Academy of Pediatric Dentistry. Definition of dental home. Chicago: AAPD; 2020. Available from: <https://www.aapd.org/research/oral-health-policies--recommendations/Dental-Home/>.
74. Clarkson JE, Pitts NB, Fee PA, Goulao B, Boyers D, Ramsay CR, et al. Examining the effectiveness of different dental recall strategies on maintenance of optimum oral health: the INTERVAL dental recalls randomised controlled trial. *Br Dent J*. 2021;230(4):236–43. Available from: <https://doi.org/10.1038/s41415-021-2612-0>.
75. Benzian H, Beltrán-Aguilar E, Niederman R. Systemic management of pandemic risks in dental practice: a consolidated framework for COVID-19 Control in dentistry. *Front Med (Lausanne)*. 2021;8:644515. Available from: <https://doi.org/10.3389/fmed.2021.644515>.
76. Benzian H, Niederman R. A dental response to the COVID-19 pandemic—safer aerosol-free emergent (SAFER) dentistry. *Front Med (Lausanne)*. 2020;7:520. Available from: <https://doi.org/10.3389/fmed.2020.00520>.
77. Andrade FB de, Antunes JLF, Andrade FCD, Lima-Costa MFF, Macinko J. Education-related inequalities in dental services use among older adults in 23 countries. *J Dent Res*. 2020 Nov;99(12):1341–1347. Available from: <https://doi.org/10.1177/0022034520935854>.
78. Vujicic M, Buchmueller T, Klein R. Dental care presents the highest level of financial barriers, compared to other types of health care services. *Health Affairs*. 2016;35(12):2176–82. Available from: <https://doi.org/10.1377/hlthaff.2016.0800>.
79. Andrade FB de, Andrade FCD, Noronha K. Measuring socioeconomic inequalities in the use of dental care services among older adults in Brazil. *Community Dent Oral Epidemiol*. 2017 Dec;45(6):559–566. Available from: <https://doi.org/10.1111/cdoe.12323>.
80. Herkrath FJ, Vettore MV, Werneck GL. Utilisation of dental services by Brazilian adults in rural and urban areas: a multi-group structural equation analysis using the Andersen behavioural model. *BMC Public Health*. 2020; 20(1):953. Available from: <https://doi.org/10.1186/s12889-020-09100-x>.
81. Mariño R. Oral health of the elderly; reality, myth, and perspective. *Bull Pan Am Health Organ*. 1994 Sep;28(3):202–10. PMID: 7951363.
82. Azañedo D, Chambergo-Michilot D, Hernández-Vásquez A. Associations between chronic conditions and oral health services utilization in older Peruvian adults: a pooled analysis of the Demographic and Health Survey 2015–2017. *Epidemiol Health*. 2020;42:e2020023. Available from: <https://doi.org/10.4178/epih.e2020023>.

83. Quinteros ME, Cáceres DD, Soto A, Mariño RJ, Giacaman RA. Caries experience and use of dental services in rural and urban adults and older adults from central Chile. *Int Dent J*. 2014 Oct;64(5):260-8. Available from: <https://doi.org/10.1111/idj.12118>.
84. Christian B, Chattopadhyay A, Kingman A, Boroumand S, Adams A, Garcia I. Oral health care services utilisation in the adult US population: Medical Expenditure Panel Survey 2006. *Community Dent Health*. 2013;30(3):161-7. PMID: 24151790.
85. Yarbrough C, Vujicic M. Oral health trends for older Americans. *J Am Dent Assoc*. 2019 Aug;150(8):714-716. Available from: <https://doi.org/10.1016/j.adaj.2019.05.026>.
86. Manski RJ, Rohde F. Dental services: use, expenses, source of payment, coverage and procedure type, 1996-2015: research findings No. 38. Rockville, MD: Agency for Healthcare Research and Quality; 2017. Available from: [https://meps.ahrq.gov/data\\_files/publications/rf38/rf38.pdf](https://meps.ahrq.gov/data_files/publications/rf38/rf38.pdf).
87. Gaskin DJ, Zare H, McCleary R, Kanwar O, Davis AL. Predictors of unmet dental health needs in US adults in 2018: a cross-sectional analysis. *JDR Clin Trans Res*. 2022 Oct;7(4):398-406. Available from: <https://doi.org/10.1177/23800844211035669>.
88. Fonesca FA, Jones KM, Mendes DC, dos Santos Neto PE, Ferreira RC, Pordeus IA, et al. The oral health of seniors in Brazil: addressing the consequences of a historic lack of public health dentistry in an unequal society. *Gerodontology*. 2015;32(1):18-27. Available from: <https://doi.org/10.1111/ger.12046>.
89. Benzian H, Guarnizo-Herreño CC, Kearns C, Muriithi MW, Watt RG. The WHO global strategy for oral health: an opportunity for bold action. *Lancet*. 2021;398(10296):192-4. Available from: [https://doi.org/10.1016/S0140-6736\(21\)01404-5](https://doi.org/10.1016/S0140-6736(21)01404-5).
90. World Health Organization. Oral health in ageing societies: integration of oral health and general health. Geneva: WHO; 2006.
91. Benzian H, Beltrán-Aguilar E, Mathur MR, Niederman R. Pandemic considerations on essential oral health care. *J Dent Res*. 2021 Mar;100(3):221-225. Available from: <https://doi.org/10.1177/0022034520979830>.
92. Fukai K, Ogawa H, Hescot P. Oral health for healthy longevity in an ageing society: maintaining momentum and moving forward. *Int Dent J*. 2017 Sep;67(Suppl. 2):3-6. Available from: <https://doi.org/10.1111/idj.12347>.
93. Centers for Disease Control and Prevention. Older adult oral health: facts about older adult oral health. Atlanta, GA: CDC; 2021. Available from: [https://www.cdc.gov/oralhealth/basics/adult-oral-health/adult\\_older.htm](https://www.cdc.gov/oralhealth/basics/adult-oral-health/adult_older.htm).

94. National Institute of Dental and Craniofacial Research. Oral health & older adults. Bethesda, MD: NIDCR; 2019. Available from: <https://www.nidcr.nih.gov/health-info/for-older-adults>.
95. FDI Dental Dental Federation. Good oral health through life. Geneva-Cointrin: FDI; [date unknown]. Available from: <https://www.fdiworlddental.org/good-oral-health-through-life>.
96. FDI World Dental Federation. Managing older adults: chairside guide. Geneva-Cointrin: FDI; 2019. Available from: <https://fdiworlddental.org/managing-older-adults-chairside-guide>.
97. Association of State and Territorial Dental Directors. ASTDD basic screening surveys. Reno, NV: ASTDD; 2022. Available from: <https://www.astdd.org/basic-screening-survey-tool#adults>.
98. Miyazaki H, Jones JA, Beltrán-Aguilar ED. Surveillance and monitoring of oral health in elderly people. *Int Dent J*. 2017 Sep;67(Suppl. 2):34-41. Available from: <https://doi.org/10.1111/idj.12348>.
99. World Health Organization. World Health Assembly resolution paves the way for better oral health care. Geneva: WHO; 2021. Available from: <https://www.who.int/news/item/27-05-2021-world-health-assembly-resolution-paves-the-way-for-better-oral-health-care>.
100. World Health Organization. Decade of Healthy Ageing 2020-2030. Geneva: WHO; 2020. Available from: [https://cdn.who.int/media/docs/default-source/decade-of-healthy-ageing/final-decade-proposal/decade-proposal-final-apr2020-en.pdf?sfvrsn=b4b75ebc\\_25&download=true](https://cdn.who.int/media/docs/default-source/decade-of-healthy-ageing/final-decade-proposal/decade-proposal-final-apr2020-en.pdf?sfvrsn=b4b75ebc_25&download=true).

*Oral Health Status of Older Adults in the Americas* is part of a series of publications entitled “The Decade of Healthy Aging in the Americas: Situation and Challenges.” In order to outline the current knowledge available on the situation of health and well-being of older persons in the Americas at the beginning of the United Nations Decade of Healthy Aging (2021–2023), this document presents data and existing evidence on oral health in the population of the Americas, especially older people, and calls for increasing actions to achieve healthy aging in the Region of the Americas by developing policies and interventions, including integrated delivery of dental care with medical and social care. Oral health is related to different aspects of health and directly influences older adults’ intrinsic capacity, especially because of the close relationship with the vitality domain. Maintaining and improving intrinsic capacity is key to achieving healthy aging. The accelerated population aging of Latin America and the Caribbean will shape the capacities of countries and health systems to respond to the specific demands of the population. Therefore, in order to design inclusive and sustainable systems, it is necessary to have updated and effective accurate information. The Decade of Healthy Aging 2021–2030 is a period to guide action toward data generation and monitoring, and this document contributes to that strategy.

